

On the number of constituents of products of characters

by

Maria Loukaki
Department of Applied Mathematics
University of Crete
Knosou Av. 74 GR-71409
Heraklion-Crete GREECE
E-mail: loukaki@tem.uoc.gr

and

Alexander Moretó
Departament d'Àlgebra
Universitat de València
46100 Burjassot. València SPAIN
E-mail: Alexander.Moreto@uv.es@

The second author was supported by the FEDER, the Spanish Ministerio de Ciencia y Educación, grants MTM2004-04665 and MTM2004-06067-C02-01, the Generalitat Valenciana and the Programa Ramón y Cajal.

In this note, p will stand for a prime number. Let φ and ψ be faithful characters of a finite p -group P . What can be said about the number of different irreducible constituents of the product $\varphi\psi$? At first sight, it does not seem reasonable to expect strong restrictions for the possible values of this number. However, in [1] it was proved that if the number of constituents of this product is bigger than one, then it is at least $(p + 1)/2$.

It seems reasonable to ask what further restrictions can be found. In p. 237 of [1] it was conjectured that if the number of constituents of the product of two faithful characters of a finite p -group, for $p \geq 5$, is bigger than $(p + 1)/2$, then it is at least p . The goal of this note is to give a counterexample to this conjecture.

Theorem. *Let $P = C_p \wr C_p$ for $p \geq 5$. There exist $\varphi, \psi \in \text{Irr}(P)$ faithful such that $\varphi\psi$ has exactly $p - 1$ distinct irreducible constituents.*

Proof. Write $P = CA$, where A is the base group, which is elementary abelian of order p^p . It is clear that the non-linear characters of P are induced from characters of A . In particular, they have degree p . Fix any non-principal character $\lambda \in \text{Irr}(C_p)$. Then any character of A can be written in the form $\nu = \lambda^{i_1} \times \cdots \times \lambda^{i_p}$ for some integers $i_j = 0, \dots, p - 1$. Thus, we can identify the character ν with the p -tuple (i_1, \dots, i_p) . It is clear that $\nu^G \in \text{Irr}(G)$ if and only if not all the i_j 's are equal.

We have that $Z(P) = \{(x, \dots, x) \mid x \in C_p\}$ is the unique minimal normal subgroup of P . Also, if $\nu^G \in \text{Irr}(G)$, it follows from Lemma 5.11 of [2] that ν^G is faithful if and only if $(x, \dots, x) \notin \text{Ker } \nu^G$. Notice that if $\lambda(x) = \varepsilon$ for a primitive p th root of unity, then

$$\nu^G(x, \dots, x) = p\varepsilon^{i_1 + \cdots + i_p}.$$

Thus ν^G is faithful if and only if $\sum_{j=1}^p i_j \not\equiv 0 \pmod{p}$.

Consider the characters of A associated to the p -tuples $(1, 0, 0, \dots, 0)$ and $(1, 1, 0, \dots, 0)$. They induce faithful irreducible characters of P , φ and ψ respectively. We claim that $\varphi\psi$ has $p - 1$ distinct irreducible constituents.

The character φ_A decomposes as the sum of the characters associated to $(1, 0, 0, \dots, 0), (0, 1, 0, \dots, 0), (0, 0, 1, \dots, 0), \dots, (0, 0, 0, \dots, 1)$. We can argue similarly with ψ_A . The product of two characters of A corresponds to the componentwise sum of the associated p -tuples and two characters of A are P -conjugate if and only if we can go from the p -tuple associated to one of the characters to the other by a cyclic permutation of the components. Now, it is easy to see that the number of constituents of the character $\varphi\psi$ is the number of characters of P lying over the characters of A corresponding to the p -tuples $(2, 1, 0, 0, \dots, 0), (1, 2, 0, 0, \dots, 0), (1, 1, 1, 0, \dots, 0), (1, 1, 0, 1, \dots, 0), \dots,$

$(1, 1, 0, 0, \dots, 1)$. Here we have p different p -tuples. The third of these p -tuples and the last one correspond to P -conjugate characters of A , so they induce the same character of P . It is easy to see that no other pair of p -tuples are conjugate. The claim follows. \square

We have been unable to find any example where the number of constituents of the product of two faithful characters of a p -group has more than $(p+1)/2$ distinct irreducible constituents but less than $p-1$. So the following modification of the conjecture could still be true.

Question. *Let φ and ψ be faithful irreducible characters of a finite p -group P . Assume that $\varphi\psi$ has more than $(p+1)/2$ distinct irreducible constituents. Does it necessarily have at least $p-1$ irreducible constituents?*

References

- [1] E. Adan-Bante, Products of characters and finite p -groups, *J. Algebra* **277** (2004), 236–255.
- [2] M. Isaacs, *Character Theory of Finite Groups*, Dover, New York, 1994.