

---

**Michael Papadimitrakis**

# **Notes on Complex Analysis**

---

Department of Mathematics

University of Crete



# Contents

<b>1</b>	<b>The complex plane and the sphere of Riemann.</b>	<b>1</b>
1.1	The complex plane. . . . .	1
1.2	Argument and polar representation. . . . .	3
1.3	Stereographic projection and the sphere of Riemann. . . . .	5
<b>2</b>	<b>Series and curvilinear integrals.</b>	<b>12</b>
2.1	Series of numbers. . . . .	12
2.2	Sequences and series of functions. . . . .	16
2.3	Curvilinear integrals. . . . .	17
<b>3</b>	<b>Holomorphic functions.</b>	<b>25</b>
3.1	Differentiability and holomorphy. . . . .	25
3.2	The Cauchy-Riemann equations. . . . .	28
3.3	Conformality. . . . .	33
<b>4</b>	<b>Examples of holomorphic functions.</b>	<b>35</b>
4.1	Linear fractional transformations. . . . .	35
4.2	The exponential function. . . . .	39
4.3	Branches of the logarithmic function. . . . .	42
4.4	Powers and branches of roots. . . . .	47
4.5	Functions defined by curvilinear integrals. . . . .	52
4.5.1	Indefinite integrals. . . . .	52
4.5.2	Integrals with parameter. . . . .	55
4.6	Functions defined by power series. . . . .	56
<b>5</b>	<b>Local behaviour and basic properties of holomorphic functions.</b>	<b>62</b>
5.1	The theorem of Cauchy for triangles. . . . .	62
5.2	Primitives and the theorem of Cauchy in convex regions. . . . .	63
5.3	Cauchy's formulas for circles and infinite differentiability. . . . .	65
5.4	Morera's theorem. . . . .	67
5.5	Liouville's theorem. The fundamental theorem of algebra. . . . .	67
5.6	Maximum principle. . . . .	68
5.7	Taylor series and Laurent series. . . . .	70
5.8	Roots and the principle of identity. . . . .	75
5.9	Isolated singularities. . . . .	78
5.10	The open mapping theorem. . . . .	83
5.11	Local mapping properties. . . . .	84
<b>6</b>	<b>Global behaviour of holomorphic functions.</b>	<b>86</b>
6.1	Index of a closed curve with respect to a point. . . . .	86
6.2	Homotopy. . . . .	91
6.3	Combinatorial results for curves and square nets. . . . .	94

6.4	The theorem of Cauchy in general open sets. . . . .	96
6.5	The residue theorem. . . . .	99
6.6	The argument principle. The theorem of Rouché. . . . .	108

# Chapter 1

## The complex plane and the sphere of Riemann.

### 1.1 The complex plane.

In  $\mathbb{R}^2$ , besides the usual vector space *addition*, which is defined by

$$(x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2),$$

there is the operation of *multiplication*, defined by

$$(x_1, y_1)(x_2, y_2) = (x_1x_2 - y_1y_2, x_1y_2 + y_1x_2).$$

We can easily prove that  $\mathbb{R}^2$  equipped with these two binary operations is an algebraic *field*. The neutral element of multiplication is  $(1, 0)$  and the inverse of  $(x, y) \neq (0, 0)$  is  $(\frac{x}{x^2+y^2}, -\frac{y}{x^2+y^2})$ .

We denote  $\mathbb{C}$  the set  $\mathbb{R}^2$  equipped with the above addition and multiplication.

It is easy to prove that the function  $\mathbb{R} \ni x \mapsto (x, 0) \in \mathbb{C}$  is a one-to-one *field homomorphism* from  $\mathbb{R}$  into  $\mathbb{C}$ . This permits the *identification* of  $\mathbb{R}$  with the subset  $\{(x, 0) \mid x \in \mathbb{R}\}$  of  $\mathbb{C}$ . In other words, we may identify  $x \in \mathbb{R}$  with the corresponding  $(x, 0) \in \mathbb{C}$  and consider  $\mathbb{R}$  as a subset of  $\mathbb{C}$ . This is exactly the same as the identification we make when we want to view  $\mathbb{R}$  as the real line, the  $x$ -axis, in the two-dimensional plane identified with  $\mathbb{R}^2$ . From now on we do not distinguish between  $x$  and  $(x, 0)$ , i.e.  $(x, 0) = x$ .

We define  $i$ , the *imaginary unit*, to be the element  $(0, 1)$  and then we have

$$(x, 0) + i(y, 0) = (x, 0) + (0, 1)(y, 0) = (x, 0) + (0, y) = (x, y).$$

If we replace  $(x, 0)$  and  $(y, 0)$  with the corresponding  $x$  and  $y$ , we get

$$(x, y) = x + iy.$$

From now on we shall write the elements of  $\mathbb{C} = \mathbb{R}^2$  in both forms:  $(x, y)$  and  $x + iy$ . We shall prefer the second,  $x + iy$ , the *complex form* of the elements of  $\mathbb{C}$ . We say that  $x + iy$  is a **complex number** and that  $\mathbb{C}$  is the set of complex numbers.

Now the definitions of addition and multiplication take the forms:

$$(x_1 + iy_1) + (x_2 + iy_2) = (x_1 + x_2) + i(y_1 + y_2),$$

$$(x_1 + iy_1)(x_2 + iy_2) = (x_1x_2 - y_1y_2) + i(x_1y_2 + y_1x_2).$$

In particular we have  $(\pm i)^2 = -1$ . We shall prove later that, besides the polynomial equation  $z^2 + 1$  which has as solutions the complex numbers  $\pm i$ , every polynomial equation with coefficients in  $\mathbb{C}$  is solvable in  $\mathbb{C}$ . In other words, we shall prove that  $\mathbb{C}$  is an *algebraically closed field*.

The usual *order relation*  $<$ , which makes  $\mathbb{R}$  an ordered field, cannot be extended in  $\mathbb{C}$ . In fact,  $\mathbb{C}$  cannot be equipped with any order relation so that it becomes an ordered field (with the addition and multiplication already defined in  $\mathbb{C}$ ). Indeed, no matter what the order relation is, we must have that an element of the form  $z^2 = z z$  is “positive” if  $z \neq 0$ , and then we end up with the contradiction:  $1 = 1^2$  is “positive” and  $-1 = i^2$  is also “positive”. Therefore, when we write inequalities like  $z \leq w$  or  $z < w$  we always accept that  $z, w$  are real numbers.

It is customary to use symbols like  $x, y, u, v, t, \xi, \eta$  for real numbers and symbols like  $z, w, \zeta$  for complex numbers. For instance, we write  $z = x + iy, w = u + iv, \zeta = \xi + i\eta$ .

For every  $z = (x, y) = x + iy$  we introduce the symbols

$$\operatorname{Re} z = x, \quad \operatorname{Im} z = y, \quad \bar{z} = (x, -y) = x - iy, \quad |z| = \sqrt{x^2 + y^2}$$

These are called **real part**, **imaginary part**, **conjugate** and **absolute value** (or **modulus**) of  $z$ , respectively.

The useful identities

$$\operatorname{Re} z = \frac{1}{2}(z + \bar{z}), \quad \operatorname{Im} z = \frac{1}{2i}(z - \bar{z}), \quad z \bar{z} = |z|^2$$

are trivial to prove.

The geometrical model for  $\mathbb{C}$  is the same as for  $\mathbb{R}^2$ , i.e. the cartesian plane with two perpendicular axes: every  $z = (x, y) = x + iy$  corresponds to the point of the plane with abscissa  $x$  and ordinate  $y$ . The horizontal axis of all points  $(x, 0) = x$  is the *real axis*. The vertical axis of all points  $(0, y) = iy$  is the *imaginary axis*. In this framework, the cartesian plane is also called **complex plane**.

We recall that the cartesian equation of the general line in the plane is

$$ax + by = c,$$

where  $a, b, c \in \mathbb{R}, a^2 + b^2 \neq 0$ . If we set  $z = x + iy$  and  $w = a + ib \neq 0$ , then the above equation takes the form

$$\operatorname{Re}(\bar{w}z) = c.$$

Similarly, the defining inequalities  $ax + by < c$  and  $ax + by > c$  of the two halfplanes on the two sides of the line with equation  $ax + by = c$  become  $\operatorname{Re}(\bar{w}z) < c$  and  $\operatorname{Re}(\bar{w}z) > c$ , respectively.

We shall denote

$$[z_1, z_2] = \{(1-t)z_1 + tz_2 \mid 0 \leq t \leq 1\}$$

the *linear segment* joining the points  $z_1, z_2$ . When we say *interval* we mean a linear segment on the real line:  $[a, b] \subseteq \mathbb{R}$ .

The *euclidean distance* between the points  $z_1 = (x_1, y_1)$  and  $z_2 = (x_2, y_2)$  is

$$\sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2} = |z_1 - z_2|.$$

Therefore, the *circle*, the *open disc* and the *closed disc* with center  $z = (x, y)$  and radius  $r > 0$  take the form

$$C_z(r) = \{w \mid |w - z| = r\}, \quad D_z(r) = \{w \mid |w - z| < r\}, \quad \bar{D}_z(r) = \{w \mid |w - z| \leq r\}.$$

We recall the special symbols  $\mathbb{T}$  for the unit circle  $C_0(1)$ ,  $\mathbb{D}$  for the unit disc  $D_0(1)$  and  $\bar{\mathbb{D}}$  for the closed unit disc  $\bar{D}_0(1)$ .

The *real part* and the *imaginary part* of a complex function  $f : A \rightarrow \mathbb{C}$ , where  $A$  is any nonempty set, are the functions  $u = \operatorname{Re} f : A \rightarrow \mathbb{R}$  and  $v = \operatorname{Im} f : A \rightarrow \mathbb{R}$ , respectively, defined by

$$u(a) = \operatorname{Re} f(a) = \frac{1}{2}(f(a) + \overline{f(a)}), \quad v(a) = \operatorname{Im} f(a) = \frac{1}{2i}(f(a) - \overline{f(a)}).$$

Of course, we have  $f(a) = u(a) + iv(a) = (u(a), v(a))$  for  $a \in A$ .

Now,  $\mathbb{C} = \mathbb{R}^2$  has the familiar euclidean metric space structure: we have the notions of interior point, boundary point, limit point and accumulation point of a set, interior  $A^\circ$ , boundary  $\partial A$  and closure  $\bar{A}$  of a set  $A$ , open set, closed set, compact set and connected set and the related properties. We also have the notions of convergence of sequences of complex numbers and limits and continuity of functions defined in  $\mathbb{C}$  or taking values in  $\mathbb{C}$ .

We only recall the following very simple properties of limits. The variable points  $z, w$  may represent the terms of a sequence or the values of a function and so we get the familiar algebraic properties of limits of sequences and of functions.

Of course, the convergence  $z \rightarrow z_0$  is equivalent to  $|z - z_0| \rightarrow 0$ . If  $z = (x, y)$  and  $z_0 = (x_0, y_0)$ , then the equivalence between  $z \rightarrow z_0$  and  $x \rightarrow x_0, y \rightarrow y_0$  now takes the form of the equivalence between  $z \rightarrow z_0$  and  $\operatorname{Re} z \rightarrow \operatorname{Re} z_0, \operatorname{Im} z \rightarrow \operatorname{Im} z_0$ . Also, if  $z \rightarrow z_0$  and  $w \rightarrow w_0$ , then  $z + w \rightarrow z_0 + w_0$  and  $zw \rightarrow z_0 w_0$ . Both can be proved either by reducing them to convergence of real and imaginary parts or -preferably- by using the triangle inequality:

$$|(z + w) - (z_0 + w_0)| \leq |z - z_0| + |w - w_0|$$

and

$$\begin{aligned} |zw - z_0 w_0| &= |(z - z_0)(w - w_0) + (z - z_0)w_0 + (w - w_0)z_0| \\ &\leq |z - z_0||w - w_0| + |z - z_0||w_0| + |w - w_0||z_0|. \end{aligned}$$

If  $z \rightarrow z_0 \neq 0$ , we can prove that  $\frac{1}{z} \rightarrow \frac{1}{z_0}$  using the equality  $|\frac{1}{z} - \frac{1}{z_0}| = \frac{|z - z_0|}{|z||z_0|}$ . We use the equality  $|\bar{z} - \bar{z}_0| = |z - z_0|$  to prove that  $z \rightarrow z_0$  implies  $\bar{z} \rightarrow \bar{z}_0$ . Similarly, we use the triangle inequality  $||z| - |z_0|| \leq |z - z_0|$  to prove that  $z \rightarrow z_0$  implies  $|z| \rightarrow |z_0|$ .

We shall consider the limit  $z \rightarrow \infty$  in section 1.3 where the point  $\infty$  will be introduced.

We also mention the standard examples of polynomial functions

$$p(z) = a_n z^n + \cdots + a_1 z + a_0$$

and rational functions

$$r(z) = \frac{p(z)}{q(z)} = \frac{a_n z^n + \cdots + a_1 z + a_0}{b_m z^m + \cdots + b_1 z + b_0}.$$

A polynomial function is continuous in  $\mathbb{C}$  and a rational function is also continuous in  $\mathbb{C}$  except at the roots of the polynomial in its denominator. Again, we shall consider the limits of  $p(z)$  and  $r(z)$  at infinity and the limits of  $r(z)$  at the roots of its denominator in section 1.3 where the point  $\infty$  will be introduced.

## 1.2 Argument and polar representation.

The trigonometric functions  $\sin$  and  $\cos$  are defined and their properties are studied in the theory of functions of a real variable. In particular, we know that  $\sin$  and  $\cos$  are periodic with smallest positive period  $2\pi$ , i.e.  $\sin(\theta + 2\pi) = \sin \theta$  and  $\cos(\theta + 2\pi) = \cos \theta$ .

Let  $I$  be any interval of length  $2\pi$  which contains only one of its endpoints, e.g.  $[0, 2\pi)$  or  $(-\pi, \pi]$ . Then we know that for every  $a, b \in \mathbb{R}$  with  $a^2 + b^2 = 1$  there exists a unique  $\theta \in I$  so that  $\cos \theta = a$  and  $\sin \theta = b$ . Equivalently, for every  $\zeta \in \mathbb{C}$  with  $|\zeta| = 1$  there exists a unique  $\theta \in I$  so that  $\zeta = \cos \theta + i \sin \theta$ . Therefore, the function

$$\cos + i \sin : \mathbb{R} \rightarrow \mathbb{T}$$

is periodic with  $2\pi$  as its smallest positive period and its restriction  $\cos + i \sin : I \rightarrow \mathbb{T}$  to any interval  $I$  of length  $2\pi$  which contains only one of its endpoints is one-to-one and onto  $\mathbb{T}$ . Thus, for every  $\zeta \in \mathbb{T}$  the equation  $\cos \theta + i \sin \theta = \zeta$  has infinitely many solutions in  $\mathbb{R}$  and exactly one solution in each interval  $I$  of length  $2\pi$  which contains only one of its endpoints.

Now, for every  $z \in \mathbb{C}$ ,  $z \neq 0$ , we have  $\frac{z}{|z|} \in \mathbb{T}$  and so the equation  $\cos \theta + i \sin \theta = \frac{z}{|z|}$  has infinitely many solutions in  $\mathbb{R}$  and exactly one solution in each interval  $I$  of length  $2\pi$  which contains only one of its endpoints. The set of all these solutions is called **argument** or **angle** of  $z$  and it is denoted  $\arg z$ , i.e.

$$\arg z = \left\{ \theta \mid \cos \theta + i \sin \theta = \frac{z}{|z|} \right\}.$$

So we have the equivalence:

$$\theta \in \arg z \iff \cos \theta + i \sin \theta = \frac{z}{|z|}.$$

Thus,  $\arg z$  has infinitely many elements and it is clear, by the  $2\pi$ -periodicity of  $\sin$  and  $\cos$ , that these elements form a (two-sided) arithmetical progression of step  $2\pi$ . In other words, if  $\theta$  is an arbitrary element of  $\arg z$ , then all elements of  $\arg z$  are described by  $\theta + k2\pi$ ,  $k \in \mathbb{Z}$ .

On the other hand, the unique solution of the equation  $\cos \theta + i \sin \theta = \frac{z}{|z|}$  in the interval  $(-\pi, \pi]$  is called **principal argument** or **principal angle** of  $z$  and it is denoted  $\text{Arg } z$ :

$$\theta = \text{Arg } z \iff \cos \theta + i \sin \theta = \frac{z}{|z|} \text{ and } -\pi < \theta \leq \pi.$$

Thus,  $\text{Arg } z$  is one of the elements of  $\arg z$ , the one which is contained in  $(-\pi, \pi]$ .

**Examples.** (i)  $\text{Arg } 3 = 0$  and  $\arg 3 = \{k2\pi \mid k \in \mathbb{Z}\}$ .

(ii)  $\text{Arg}(4i) = \frac{\pi}{2}$  and  $\arg(4i) = \{\frac{\pi}{2} + k2\pi \mid k \in \mathbb{Z}\}$ .

(iii)  $\text{Arg}(-2) = \pi$  and  $\arg(-2) = \{\pi + k2\pi \mid k \in \mathbb{Z}\}$ .

(iv)  $\text{Arg}(1+i) = \frac{\pi}{4}$  and  $\arg(1+i) = \{\frac{\pi}{4} + k2\pi \mid k \in \mathbb{Z}\}$ .

(v)  $\text{Arg}(-1-i\sqrt{3}) = \frac{4\pi}{3}$  and  $\arg(-1-i\sqrt{3}) = \{\frac{4\pi}{3} + k2\pi \mid k \in \mathbb{Z}\}$ .

We remark that *we do not define argument or angle for the number 0*.

Since the elements of  $\arg z$  form an arithmetical progression of step  $2\pi$ , it is obvious that, if  $z_1, z_2 \neq 0$ , then either  $\arg z_1 = \arg z_2$  or  $\arg z_1 \cap \arg z_2 = \emptyset$ . More precisely,  $\arg z_1 = \arg z_2$  if and only if  $\frac{z_1}{z_2} > 0$  or, equivalently, if and only if  $z_1, z_2$  belong to the same halfline with vertex 0.

Comparing real and imaginary parts of the two sides of the following identity, we see that it is equivalent to the well-known addition formulas of  $\sin$  and  $\cos$ :

$$\cos(\theta_1 + \theta_2) + i \sin(\theta_1 + \theta_2) = (\cos \theta_1 + i \sin \theta_1)(\cos \theta_2 + i \sin \theta_2).$$

A direct consequence by induction is the familiar *formula of de Moivre*:

$$\cos(n\theta) + i \sin(n\theta) = (\cos \theta + i \sin \theta)^n \quad \text{for every } n \in \mathbb{Z}.$$

**Proposition 1.1.** *For every nonzero  $z_1, z_2$  we have*

$$\arg(z_1 z_2) = \arg z_1 + \arg z_2.$$

*By this we mean that the sum of any element of  $\arg z_1$  and any element of  $\arg z_2$  is an element of  $\arg(z_1 z_2)$  and, conversely, any element of  $\arg(z_1 z_2)$  is the sum of an element of  $\arg z_1$  and an element of  $\arg z_2$ .*

*Proof.* We take any  $\theta_1 \in \arg z_1$  and any  $\theta_2 \in \arg z_2$  and  $\theta = \theta_1 + \theta_2$ . Then by the addition formulas,  $\cos \theta + i \sin \theta = (\cos \theta_1 + i \sin \theta_1)(\cos \theta_2 + i \sin \theta_2) = \frac{z_1}{|z_1|} \frac{z_2}{|z_2|} = \frac{z_1 z_2}{|z_1 z_2|}$ . Therefore,  $\theta \in \arg(z_1 z_2)$ .

Conversely, we take any  $\theta \in \arg(z_1 z_2)$ . We consider  $\theta_1 \in \arg z_1$  and we define  $\theta_2 = \theta - \theta_1$ . Then  $\cos \theta_2 + i \sin \theta_2 = \frac{\cos \theta + i \sin \theta}{\cos \theta_1 + i \sin \theta_1} = \frac{z_1 z_2}{|z_1 z_2|} / \frac{z_1}{|z_1|} = \frac{z_2}{|z_2|}$ . Therefore,  $\theta_2 \in \arg z_2$  and  $\theta = \theta_1 + \theta_2$ .  $\square$

We note that the equality  $\text{Arg}(z_1 z_2) = \text{Arg } z_1 + \text{Arg } z_2$  is not true in general.



**Example 1.2.1.**  $\text{Arg}(-1) + \text{Arg}(-1) = \pi + \pi = 2\pi$ , while  $\text{Arg}((-1)(-1)) = \text{Arg} 1 = 0$ .

The equalities  $|z_1 z_2| = |z_1| |z_2|$  and  $\arg(z_1 z_2) = \arg z_1 + \arg z_2$  express the well-known geometric rule: *when two complex numbers are multiplied, their distances from 0 are multiplied and their angles are added.*

It is clear by now that for every  $z \neq 0$  we may write

$$z = r(\cos \theta + i \sin \theta),$$

where  $r = |z|$  and  $\theta \in \arg z$ . This is called a **polar representation** of  $z$ . There are infinitely many polar representations of  $z$ , one for each  $\theta \in \arg z$ . The polar representation with  $\theta = \text{Arg} z$  is called **principal polar representation** of  $z$ .

As in the case of the argument, *we do not define polar representation for the number 0.*

### Exercises.

**1.2.1.** Which are all the possible values of  $\text{Arg}(z_1 z_2) - \text{Arg} z_1 - \text{Arg} z_2$ ?

**1.2.2.** Prove that  $\arg \frac{1}{z} = \arg \bar{z} = -\arg z$  and  $\arg(-z) = \pi + \arg z$ , after you assign the proper meaning to these equalities.

**1.2.3.** Prove the following statement for any nonzero  $z, z_1$  and  $z_2$ . It is true that  $z = z_1 z_2$  if and only if the triangle  $T(0, 1, z_1)$  with vertices  $0, 1, z_1$  is *similar* to the triangle  $T(0, z_2, z)$  with vertices  $0, z_2, z$  ( $0$  corresponding to  $0$ ,  $1$  corresponding to  $z_2$  and  $z_1$  corresponding to  $z$ ). This expresses the geometric visualization of the operation of multiplication in  $\mathbb{C}$ .

## 1.3 Stereographic projection and the sphere of Riemann.

Let  $\mathbb{S}^2 = \{(\xi, \eta, \zeta) \in \mathbb{R}^3 \mid \xi^2 + \eta^2 + \zeta^2 = 1\}$  be the unit sphere in  $\mathbb{R}^3$ . Through the usual identifications, we consider  $\mathbb{C} = \mathbb{R}^2$  as the set of points  $z = x + iy = (x, y) = (x, y, 0)$  of  $\mathbb{R}^3$ .

A characteristic point of  $\mathbb{S}^2$  is the north pole  $N = (0, 0, 1)$ . We take any  $z = x + iy \in \mathbb{C}$  and the line  $Nz$  in  $\mathbb{R}^3$ , which contains  $N$  and  $z$ . Clearly, this line intersects  $\mathbb{S}^2$  at  $N$ . We shall see that there is a second point of intersection  $A = (\xi, \eta, \zeta)$  of  $Nz$  and  $\mathbb{S}^2$ . That  $A = (\xi, \eta, \zeta)$  belongs to  $Nz$  is equivalent to  $\overrightarrow{NA} = t \overrightarrow{Nz}$  for some  $t \in \mathbb{R}$ . This is equivalent to

$$\begin{aligned} \xi - 0 &= t(x - 0) \\ \eta - 0 &= t(y - 0) \\ \zeta - 1 &= t(0 - 1) \end{aligned} \tag{1.1}$$

On the other hand, that  $A = (\xi, \eta, \zeta)$  belongs to  $\mathbb{S}^2$  is equivalent to

$$\xi^2 + \eta^2 + \zeta^2 = 1. \tag{1.2}$$

That  $A = (\xi, \eta, \zeta)$  is a common point of  $Nz$  and  $\mathbb{S}^2$  is equivalent to  $(\xi, \eta, \zeta, t)$  being a solution of the system of the four equations (1.1) and (1.2). We easily solve this system and we find two distinct solutions: the point  $N = (0, 0, 1)$ , which we already know, and the point

$$A = \left( \frac{2x}{x^2+y^2+1}, \frac{2y}{x^2+y^2+1}, \frac{x^2+y^2-1}{x^2+y^2+1} \right).$$

Now we consider the mapping

$$\mathbb{C} \ni z = x + iy \longmapsto A = (\xi, \eta, \zeta) = \left( \frac{2x}{x^2+y^2+1}, \frac{2y}{x^2+y^2+1}, \frac{x^2+y^2-1}{x^2+y^2+1} \right) \in \mathbb{S}^2 \setminus \{N\}$$

from  $\mathbb{C}$  to  $\mathbb{S}^2 \setminus \{N\}$ . We check easily that this mapping is one-to-one and onto  $\mathbb{S}^2 \setminus \{N\}$  and that the inverse mapping is

$$\mathbb{S}^2 \setminus \{N\} \ni A = (\xi, \eta, \zeta) \longmapsto z = x + iy = \frac{\xi}{1-\zeta} + i \frac{\eta}{1-\zeta} \in \mathbb{C}.$$

The two mutually inverse mappings just defined between  $\mathbb{C}$  and  $\mathbb{S}^2 \setminus \{N\}$  are called **stereographic projections**. We write  $\mathbb{C} \leftrightarrow \mathbb{S}^2 \setminus \{N\}$  to denote the action of the two stereographic projections.

We shall see now that both stereographic projections are continuous. We take two points  $z = x + iy$  and  $z_0 = x_0 + iy_0$  in  $\mathbb{C}$ . Let their images, through stereographic projection, be the points  $A = (\xi, \eta, \zeta)$  and  $A_0 = (\xi_0, \eta_0, \zeta_0)$  in  $\mathbb{S}^2 \setminus \{N\}$ . Using the formulas of stereographic projection, we can prove that the euclidean distance in  $\mathbb{R}^3$  between  $A$  and  $A_0$  equals

$$\begin{aligned} |A - A_0| &= \sqrt{(\xi - \xi_0)^2 + (\eta - \eta_0)^2 + (\zeta - \zeta_0)^2} \\ &= \dots = \frac{2\sqrt{(x-x_0)^2 + (y-y_0)^2}}{\sqrt{x^2+y^2+1}\sqrt{x_0^2+y_0^2+1}} = \frac{2|z-z_0|}{\sqrt{|z|^2+1}\sqrt{|z_0|^2+1}}. \end{aligned} \quad (1.3)$$

We also take  $z = x + iy$  in  $\mathbb{C}$  and let its image, through stereographic projection, be  $A = (\xi, \eta, \zeta)$  in  $\mathbb{S}^2 \setminus \{N\}$ . We find that the euclidean distance in  $\mathbb{R}^3$  between  $A$  and  $N$  equals

$$|A - N| = \sqrt{(\xi - 0)^2 + (\eta - 0)^2 + (\zeta - 1)^2} = \dots = \frac{2}{\sqrt{x^2+y^2+1}} = \frac{2}{\sqrt{|z|^2+1}}. \quad (1.4)$$

Now, if  $z \rightarrow z_0$ , then (1.3) implies that  $A \rightarrow A_0$ . Conversely, assume that  $A \rightarrow A_0$ . Then  $A \not\rightarrow N$  and (1.4) shows that  $|z|$  stays bounded. Hence (1.3) implies that  $z \rightarrow z_0$ . We conclude that both *stereographic projections are homeomorphisms between the metric spaces  $\mathbb{C}$  and  $\mathbb{S}^2 \setminus \{N\}$* .

We can continue the previous argument and examine the behaviour of  $z$  in  $\mathbb{C}$  when its image  $A$  in  $\mathbb{S}^2 \setminus \{N\}$  tends to the north pole  $N$ . Indeed, (1.4) shows that  $A \rightarrow N$  if and only if  $|z| \rightarrow +\infty$ . In other words,  $A \rightarrow N$  if and only if the euclidean distance of  $z$  from 0 becomes arbitrarily large.

Now, it is natural to introduce and attach to  $\mathbb{C}$  an ‘‘ideal point’’, denoted  $\infty$  and called **infinity**, whose euclidean distance from 0 is  $+\infty$ . We define the **extended complex plane** or the **sphere of Riemann** to be

$$\widehat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}.$$

We also extend the previously defined stereographic projections  $\mathbb{C} \leftrightarrow \mathbb{S}^2 \setminus \{N\}$  to be the **stereographic projections**  $\widehat{\mathbb{C}} \leftrightarrow \mathbb{S}^2$  which map each of  $\infty \in \widehat{\mathbb{C}}$  and  $N \in \mathbb{S}^2$  onto the other.

Thus, both stereographic projections  $\widehat{\mathbb{C}} \leftrightarrow \mathbb{S}^2$  are bijective mappings between  $\widehat{\mathbb{C}}$  and  $\mathbb{S}^2$ . We have seen that their restrictions  $\mathbb{C} \leftrightarrow \mathbb{S}^2 \setminus \{N\}$  are homeomorphisms between the metric spaces  $\mathbb{C}$  and  $\mathbb{S}^2 \setminus \{N\}$ . In order to examine the continuity properties of the extended stereographic projections, we have to equip the sets  $\widehat{\mathbb{C}}$  and  $\mathbb{S}^2$  with corresponding metrics. The metric on  $\mathbb{S}^2 \setminus \{N\}$ , i.e. the euclidean distance on  $\mathbb{R}^3$ , is also a metric on  $\mathbb{S}^2$ . But it is clear that the euclidean metric on  $\mathbb{C}$  cannot be extended to become a metric on  $\widehat{\mathbb{C}}$ . The problem can be solved if we use the equalities (1.3) and (1.4) to transfer the metric on  $\mathbb{S}^2$  to a metric on  $\widehat{\mathbb{C}}$ . If  $z, z_0 \in \mathbb{C}$ , we consider their images  $A, A_0 \in \mathbb{S}^2 \setminus \{N\}$  and we define the new distance between  $z, z_0$  to be equal to the euclidean distance in  $\mathbb{R}^3$  between  $A, A_0$  given by (1.3) in terms of  $z, z_0$ . If  $z \in \mathbb{C}$  and  $z_0 = \infty$ , we consider their images  $A \in \mathbb{S}^2 \setminus \{N\}$  and  $A_0 = N$  and we define the new distance between  $z, z_0$  to be equal to the euclidean distance in  $\mathbb{R}^3$  between  $A, A_0$  given by (1.4) in terms of  $z$ . The new distance between two points of  $\widehat{\mathbb{C}}$  or between a point of  $\widehat{\mathbb{C}}$  and  $\infty$  is called *chordal distance*. In other words, we define the **chordal distance**  $\chi(z_1, z_2)$  between  $z_1, z_2$  in  $\widehat{\mathbb{C}}$  to be the euclidean distance in  $\mathbb{R}^3$  between their images, through stereographic projection, in  $\mathbb{S}^2$ . I.e.

$$\chi(z_1, z_2) = \begin{cases} \frac{2|z_1 - z_2|}{\sqrt{|z_1|^2 + 1}\sqrt{|z_2|^2 + 1}}, & \text{if } z_1, z_2 \in \mathbb{C} \\ \frac{2}{\sqrt{|z|^2 + 1}}, & \text{if } z_1 = z \in \mathbb{C}, z_2 = \infty \text{ or } z_1 = \infty, z_2 = z \in \mathbb{C} \\ 0, & \text{if } z_1 = z_2 = \infty \end{cases}$$

**Proposition 1.2.** *The function  $\chi : \widehat{\mathbb{C}} \times \widehat{\mathbb{C}} \rightarrow \mathbb{R}$  is a metric on  $\widehat{\mathbb{C}}$ .*

*Proof.* We must prove that chordal distance has the following basic properties:

- (i)  $\chi(z_1, z_2) \geq 0$  for every  $z_1, z_2 \in \widehat{\mathbb{C}}$ .
- (ii) If  $z_1, z_2 \in \widehat{\mathbb{C}}$ , then:  $\chi(z_1, z_2) = 0$  if and only if  $z_1 = z_2$ .
- (iii)  $\chi(z_1, z_2) = \chi(z_2, z_1)$  for every  $z_1, z_2 \in \widehat{\mathbb{C}}$ .
- (iv)  $\chi(z_1, z_3) \leq \chi(z_1, z_2) + \chi(z_2, z_3)$  for every  $z_1, z_2, z_3 \in \widehat{\mathbb{C}}$ .

The first three properties are obvious. The fourth, the triangle inequality, can be proved after many calculations using the formula of the chordal distance. But there is a better way. If we take the stereographic projections  $A_1, A_2, A_3$  in  $\mathbb{S}^2$  of  $z_1, z_2, z_3$ , then from the definition of the chordal distance we have  $\chi(z_i, z_j) = |A_i - A_j|$  and, since euclidean distance in  $\mathbb{R}^3$  satisfies the triangle inequality, we get  $\chi(z_1, z_3) = |A_1 - A_3| \leq |A_1 - A_2| + |A_2 - A_3| = \chi(z_1, z_2) + \chi(z_2, z_3)$ .  $\square$

The metric  $\chi$  on  $\widehat{\mathbb{C}}$  is called **chordal metric**.

We thus have a second way to measure distances in the complex plane. Besides the euclidean distance  $|z_1 - z_2|$  we also have the chordal distance  $\chi(z_1, z_2) = \frac{2|z_1 - z_2|}{\sqrt{|z_1|^2 + 1}\sqrt{|z_2|^2 + 1}}$ .

**Proposition 1.3.**  $\widehat{\mathbb{C}}$  with the chordal metric and  $\mathbb{S}^2$  with the euclidean metric of  $\mathbb{R}^3$  are homeomorphic metric spaces.

*Proof.* Stereographic projections are homeomorphisms between the two metric spaces. In fact they are more than that: they are **isometries**. Indeed, if  $z_1, z_2 \in \widehat{\mathbb{C}}$  correspond to  $A_1, A_2 \in \mathbb{S}^2$ , then by the definition of the chordal metric we have  $\chi(z_1, z_2) = |A_1 - A_2|$ . I.e. stereographic projections preserve distances and hence they are both continuous.  $\square$

Proposition 1.4 describes the relation between the chordal metric and the euclidean metric in their common domain.

**Proposition 1.4.** The chordal metric on  $\mathbb{C}$  and the euclidean metric on  $\mathbb{C}$  are equivalent.

*Proof.* Indeed, if  $z, z_0 \in \mathbb{C}$ , then  $z \rightarrow z_0$  with respect to the euclidean distance if and only if  $z \rightarrow z_0$  with respect to the chordal distance. To see this we consider the images  $A, A_0 \in \mathbb{S}^2 \setminus \{N\}$  of  $z, z_0$  under stereographic projection. We have proved already that  $z \rightarrow z_0$  with respect to the euclidean distance in  $\mathbb{C}$  if and only if  $A \rightarrow A_0$  with respect to the euclidean distance in  $\mathbb{R}^3$ . But the euclidean distance between  $A, A_0$  is equal to the chordal distance between  $z, z_0$ . Therefore,  $|z - z_0| \rightarrow 0 \Leftrightarrow |A - A_0| \rightarrow 0 \Leftrightarrow \chi(z, z_0) \rightarrow 0$ .  $\square$

**Proposition 1.5.** Let  $z \in \mathbb{C}$ . Then  $z \rightarrow \infty$  in  $\widehat{\mathbb{C}}$  if and only if  $|z| \rightarrow +\infty$ .

*Proof.* This is obvious from  $\chi(z, \infty) = 2/\sqrt{|z|^2 + 1}$ .  $\square$

We have introduced  $\infty$  as the ideal point towards which a variable point  $z$  on the complex plane moves when its euclidean distance from 0 becomes arbitrarily large. *It is time to mention the difference with the ideal points  $\pm\infty$  we attach to  $\mathbb{R}$ .* A variable point  $x$  on the real line moves away from 0 in exactly two specific directions: either to the left or to the right and then we say, respectively, that it moves *towards*  $-\infty$  or *towards*  $+\infty$ . On the plane though there are no two uniquely specified directions. A point can move away from 0 either on arbitrary halflines (i.e. in infinitely many directions) or making an arbitrary “spiral-like movement” or in a completely arbitrary manner. Therefore, we may only say that the point moves *towards infinity*.

Now let us say a few things about neighborhoods of points in  $\widehat{\mathbb{C}}$  with respect to the chordal metric. We start with the neighborhoods of  $\infty$ . If we denote  $N_x(r)$  the  $r$ -neighborhood of a point  $x$  in the general metric space, then the  $r$ -neighborhood of  $\infty$  in the metric space  $(\widehat{\mathbb{C}}, \chi)$  is the set

$$\begin{aligned} N_\infty(r) &= \{z \in \widehat{\mathbb{C}} \mid \chi(z, \infty) < r\} = \{z \in \mathbb{C} \mid 2/\sqrt{|z|^2 + 1} < r\} \cup \{\infty\} \\ &= \begin{cases} \{z \in \mathbb{C} \mid |z| > \sqrt{(4/r^2) - 1}\} \cup \{\infty\}, & \text{if } 0 < r \leq 2 \\ \widehat{\mathbb{C}}, & \text{if } r > 2 \end{cases} \end{aligned}$$

We observe that the “small” neighborhoods of  $\infty$ , i.e. the neighborhoods  $N_\infty(r)$  with  $0 < r < 2$ , are the complements of closed discs in  $\mathbb{C}$  with center 0, together with  $\infty$ . To simplify notation we make the change of variable:  $\frac{1}{s} = \sqrt{\frac{4}{r^2} - 1}$ . When  $r$  increases in  $(0, 2)$ ,  $s$  increases in  $(0, +\infty)$  and conversely. We call  $s$ -**neighborhood** of  $\infty$  in  $\widehat{\mathbb{C}}$  the set

$$D_\infty(s) = \{z \mid |z| > \frac{1}{s}\} \cup \{\infty\},$$

i.e. the complement of the closed disc with center 0 and radius  $\frac{1}{s}$ , together with  $\infty$ .

We see that the neighborhoods of  $\infty$  in  $\widehat{\mathbb{C}}$  with respect to the chordal metric are of three kinds: the sets  $D_\infty(s)$  with  $s > 0$ , the set  $\widehat{\mathbb{C}} \setminus \{0\}$  (the case  $r = 2$  or, equivalently,  $s = +\infty$ ) and the whole set  $\widehat{\mathbb{C}}$ . Since in any metric space the “small” neighborhoods are those which characterize interior points, boundary points, limit points, limits of functions or sequences etc., in the case of  $\widehat{\mathbb{C}}$  and its point  $\infty$  we shall pay attention only to the neighborhoods of the form  $D_\infty(s)$ .

Now the following should be clear.

- (i) The point  $\infty$  is an interior point of  $A \subseteq \widehat{\mathbb{C}}$  with respect to the chordal metric if and only if  $A$  contains, besides  $\infty$ , the complement of a closed disc in  $\mathbb{C}$  with center 0.
- (ii) The point  $\infty$  is not a limit point of  $A \subseteq \widehat{\mathbb{C}}$  with respect to the chordal metric if and only if  $A$  is contained in a closed disc with center 0 or, equivalently,  $A$  is a bounded set in  $\mathbb{C}$  with respect to the euclidean metric.
- (iii) If  $\infty \notin A$ , i.e. if  $A \subseteq \mathbb{C}$ , then the following four statements are equivalent:  $\infty$  is a boundary point of  $A$  with respect to the chordal metric,  $\infty$  is a limit point of  $A$  with respect to the chordal metric,  $\infty$  is an accumulation point of  $A$  with respect to the chordal metric,  $A$  is not bounded in  $\mathbb{C}$  with respect to the euclidean metric.

Now we continue with the neighborhoods with respect to the chordal metric of a point  $z_0 \in \mathbb{C}$ . The  $r$ -neighborhood of  $z_0 \in \mathbb{C}$  in  $\widehat{\mathbb{C}}$  with respect to the chordal metric is the set

$$N_{z_0}(r) = \{z \in \widehat{\mathbb{C}} \mid \chi(z, z_0) < r\}.$$

This set does not have a simple form. Depending on the exact values of  $z_0$  and  $r$ , it is an open disc or an open halfplane or the complement of a closed disc (together with  $\infty$ ). Even when  $N_{z_0}(r)$  is an open disc,  $z_0$  is *not* its euclidean center. Look at exercise 1.3.2 for details. Since the chordal metric and the euclidean metric are equivalent in  $\mathbb{C}$ , we have the following relation between neighborhoods  $N_{z_0}(r)$  with respect to the chordal metric and neighborhoods (i.e. the familiar discs)  $D_{z_0}(r)$  with respect to the euclidean metric: for every  $\epsilon > 0$  there is  $\delta > 0$  so that  $D_{z_0}(\delta) \subseteq N_{z_0}(\epsilon)$  and, conversely, for every  $\epsilon > 0$  there is  $\delta > 0$  so that  $N_{z_0}(\delta) \subseteq D_{z_0}(\epsilon)$ . From this we conclude easily that  $z_0 \in \mathbb{C}$  is an interior point or a boundary point or a limit point of a set  $A \subseteq \widehat{\mathbb{C}}$  with respect to the chordal metric if and only if it is, respectively, an interior point or a boundary point or a limit point of  $A$  with respect to the euclidean metric.

If  $A \subseteq \mathbb{C}$  and we write  $A^\circ$ ,  $\partial A$  and  $\overline{A}$  for the interior, the boundary and the closure of  $A$  with respect to the euclidean metric and  $A^{\circ, \chi}$ ,  $\partial_\chi A$  and  $\overline{A}^\chi$  for the interior, the boundary and the closure of  $A$  with respect to the chordal metric, then we easily see that

$$A^{\circ, \chi} = A^\circ, \quad \partial_\chi A = \partial A, \quad \overline{A}^\chi = \overline{A} \quad \text{if } A \text{ is a bounded } \subseteq \mathbb{C}$$

and

$$A^{\circ, \chi} = A^\circ, \quad \partial_\chi A = \partial A \cup \{\infty\}, \quad \overline{A}^\chi = \overline{A} \cup \{\infty\} \quad \text{if } A \text{ is an unbounded } \subseteq \mathbb{C}.$$

Of course, when we say bounded or unbounded we mean with respect to the euclidean metric.

For instance, if  $A \subseteq \mathbb{C}$  is bounded, then it is open with respect to the chordal metric if and only if it is open with respect to the euclidean metric, and it is closed with respect to the chordal metric if and only if it is closed with respect to the euclidean metric. If  $A \subseteq \mathbb{C}$  is *not bounded*, then again it is open with respect to the chordal metric if and only if it is open with respect to the euclidean

metric, but, even if it is closed with respect to the euclidean metric, we have to attach  $\infty$  to  $A$  to make it closed with respect to the chordal metric.

Regarding compactness, we know that  $\mathbb{C}$  is not compact either with respect to the euclidean metric or with respect to the chordal metric. Indeed,  $\mathbb{C}$  is not compact with respect to the euclidean metric, because it is not bounded. And then it is not compact with respect to the chordal metric, because the two metrics are equivalent in  $\mathbb{C}$ . But  $\widehat{\mathbb{C}}$  is compact (with respect to the chordal metric, of course). Indeed,  $\widehat{\mathbb{C}}$  is homeomorphic to  $\mathbb{S}^2$ , which is compact since it is a closed and bounded set in  $\mathbb{R}^3$ . Now,  $\widehat{\mathbb{C}}$  is produced from  $\mathbb{C}$  by the attachment to  $\mathbb{C}$  of the single point  $\infty$ . This situation has a name in topology: we say that  $\widehat{\mathbb{C}}$  is a **one-point compactification** of  $\mathbb{C}$ .

Based on the usual algebraic rules of limits, we may extend in the standard way the algebraic operations in the set  $\widehat{\mathbb{C}}$ :

$$\begin{aligned} z + \infty = \infty + z = \infty, \quad -\infty = \infty, \quad z - \infty = \infty - z = \infty, \\ z \infty = \infty z = \infty \quad \text{if } z \neq 0, \quad \infty \infty = \infty, \\ \frac{1}{\infty} = 0, \quad \frac{1}{0} = \infty, \quad \frac{z}{\infty} = 0, \quad \frac{\infty}{z} = \infty, \\ \overline{\infty} = \infty, \quad |\infty| = +\infty. \end{aligned}$$

For example, the rule  $z_0 + \infty = \infty$  (when  $z_0 \in \mathbb{C}$ ) can be based on the following argument. If  $z \rightarrow z_0$  and  $w \rightarrow \infty$  in  $\widehat{\mathbb{C}}$ , then  $|z - z_0| \rightarrow 0$  and  $|w| \rightarrow +\infty$  and then, by the triangle inequality,  $|z + w| \geq |w| - |z - z_0| \rightarrow +\infty$ . Hence  $z + w \rightarrow \infty$  in  $\widehat{\mathbb{C}}$ . All other rules can be based on similar arguments.

The following are not defined:

$$\infty + \infty, \quad \infty - \infty, \quad 0 \infty, \quad \infty 0, \quad \frac{\infty}{\infty}, \quad \frac{0}{0}.$$

They are called **indeterminate forms**.

For instance, regarding the case of  $\infty + \infty$ , one can easily find examples of points  $z, w$  such that  $z \rightarrow \infty$  and  $w \rightarrow \infty$  but such that  $z + w$  has either no limit or any preassigned limit. The same is true in all other cases.

Observe the case of  $\frac{1}{0} = \infty$ . In  $\mathbb{R}$  the expression  $\frac{1}{0}$  is an indeterminate form, since when the real number  $x$  is small and  $> 0$  then  $\frac{1}{x}$  is large and  $> 0$  and hence  $\frac{1}{x}$  moves towards  $+\infty$ , and when  $x$  is small and  $< 0$  then  $\frac{1}{x}$  is large and  $< 0$  and hence  $\frac{1}{x}$  moves towards  $-\infty$ . But in  $\mathbb{C}$ , when  $z$  is small, i.e. when  $|z|$  is small (and necessarily  $> 0$ ), then the distance  $|\frac{1}{z}| = \frac{1}{|z|}$  of  $\frac{1}{z}$  from 0 is large and hence  $\frac{1}{z}$  moves towards  $\infty$ . So we define  $\frac{1}{0} = \infty$ .

**Example 1.3.1.** Let us consider any polynomial function  $p(z) = a_n z^n + a_{n-1} z^{n-1} + \dots + a_1 z + a_0$  with  $a_n \neq 0$ . The domain of definition of  $p$  is  $\mathbb{C}$ .

For every  $z_0 \in \mathbb{C}$  we have

$$\lim_{z \rightarrow z_0} p(z) = p(z_0),$$

using the algebraic rules of limits and the trivial limits:  $\lim_{z \rightarrow z_0} c = c$  and  $\lim_{z \rightarrow z_0} z = z_0$ . Therefore,  $p$  is continuous in  $\mathbb{C}$ .

If the degree of  $p$  is  $\geq 1$ , i.e.  $n \geq 1$  and  $a_n \neq 0$ , then

$$\lim_{z \rightarrow \infty} p(z) = \infty$$

since  $p(z) = z^n(a_n + a_{n-1}\frac{1}{z} + \dots + a_0\frac{1}{z^n}) \rightarrow \infty a_n = \infty$ .

Thus, if the degree of  $p$  is  $\geq 1$ , we may define  $p(\infty) = \infty$  and then  $p : \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$  is continuous in  $\widehat{\mathbb{C}}$ . If the degree of  $p$  is 0, then the function is constant:  $p(z) = a_0$  for all  $z$ . Hence

$$\lim_{z \rightarrow \infty} p(z) = a_0.$$

In this case we may define  $p(\infty) = a_0$  and again  $p : \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$  is continuous in  $\widehat{\mathbb{C}}$ .

**Example 1.3.2.** Now we take a rational function  $r(z) = \frac{p(z)}{q(z)} = \frac{a_n z^n + \dots + a_1 z + a_0}{b_m z^m + \dots + b_1 z + b_0}$  with  $a_n, b_m \neq 0$ . The domain of definition of  $r$  is  $\mathbb{C} \setminus \{z_1, \dots, z_s\}$ , where  $z_1, \dots, z_s$  are the roots of  $q$ . If  $z_0 \in \mathbb{C}$  and  $q(z_0) \neq 0$ , then using the algebraic rules of limits, we get:

$$\lim_{z \rightarrow z_0} r(z) = r(z_0).$$

Therefore  $r$  is continuous in its domain of definition.

Writing  $r$  in the form  $r(z) = z^{n-m}(a_n + a_{n-1}\frac{1}{z} + \dots + a_0\frac{1}{z^n})/(b_m + b_{m-1}\frac{1}{z} + \dots + b_0\frac{1}{z^m})$ , we can prove that

$$\lim_{z \rightarrow \infty} r(z) = \begin{cases} \infty, & \text{if } n > m \\ \frac{a_n}{b_m}, & \text{if } n = m \\ 0, & \text{if } n < m \end{cases}$$

Finally, let  $z_0 \in \mathbb{C}$  and  $q(z_0) = 0$ . Thus  $z_0$  is any of the roots  $z_1, \dots, z_s$  of  $q$ . Then  $z - z_0$  divides  $q(z)$ , and there is  $k \geq 1$  and a polynomial  $q_1(z)$  so that  $q(z) = (z - z_0)^k q_1(z)$  and  $q_1(z_0) \neq 0$ . This means that the *multiplicity* of the root  $z_0$  of  $q(z)$  is  $k$ . There is also  $l \geq 0$  and a polynomial  $p_1(z)$  so that  $p(z) = (z - z_0)^l p_1(z)$  and  $p_1(z_0) \neq 0$ . Indeed, if  $p(z_0) = 0$ , then  $l \geq 1$  is the multiplicity of  $z_0$  as a root of  $p(z)$  and, if  $p(z_0) \neq 0$ , we take  $l = 0$  (and say that the multiplicity of  $z_0$  as a root of  $p(z)$  is zero) and  $p_1(z) = p(z)$ . So for every  $z$  different from the roots of  $q(z)$  we have  $r(z) = (z - z_0)^{l-k} \frac{p_1(z)}{q_1(z)}$  and  $p_1(z_0) \neq 0, q_1(z_0) \neq 0$ . Now  $\frac{p_1(z_0)}{q_1(z_0)}$  is neither  $\infty$  nor 0, and hence

$$\lim_{z \rightarrow z_0} r(z) = \begin{cases} \infty, & \text{if } k > l \\ \frac{p_1(z_0)}{q_1(z_0)}, & \text{if } k = l \\ 0, & \text{if } k < l \end{cases}$$

Exactly as in the polynomial case, a rational function can be considered to be a function  $r : \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$  continuous in  $\widehat{\mathbb{C}}$ . Indeed, at every  $z_0 \in \widehat{\mathbb{C}}$  a rational function  $r$  has a specific limit. If  $z_0$  is in the usual domain of definition of  $r$ , then the limit of  $r$  at  $z_0$  coincides with  $r(z_0)$ . If  $z_0$  is either  $\infty$  or a root of the denominator of  $r$ , then we define  $r(z_0)$  to be the limit of  $r$  at  $z_0$ .

**Example 1.3.3.** The sequence  $((-2)^n)$  does not have a limit as a real sequence since its subsequences of the odd and the even indices have the different limits  $-\infty$  and  $+\infty$ . But as a complex sequence  $((-2)^n)$  tends to  $\infty$ , because  $|(-2)^n| = 2^n \rightarrow +\infty$ .

**Example 1.3.4.** Let us consider the geometric progression  $(z^n)$ .

If  $|z| < 1$ , then  $|z^n - 0| = |z|^n \rightarrow 0$  and hence  $z^n \rightarrow 0$ .

If  $|z| > 1$ , then  $|z^n| = |z|^n \rightarrow +\infty$  and hence  $z^n \rightarrow \infty$ .

If  $z = 1$ , then  $z^n = 1 \rightarrow 1$ .

Finally, let  $|z| = 1, z \neq 1$  and assume that  $z^n \rightarrow w$ . Since  $|z^n| = |z|^n = 1$  for every  $n$ , we find that  $|w| = 1$ . From  $z^n \rightarrow w$  we have  $z = \frac{z^{n+1}}{z^n} \rightarrow \frac{w}{w} = 1$  and we arrive at a contradiction.

Thus:

$$z^n \begin{cases} \rightarrow 0, & \text{if } |z| < 1 \\ \rightarrow 1, & \text{if } z = 1 \\ \rightarrow \infty, & \text{if } |z| > 1 \\ \text{has no limit,} & \text{if } |z| = 1, z \neq 1 \end{cases}$$

### Exercises.

**1.3.1.** Prove that  $\chi(z_1, z_2) \leq 2$  for every  $z_1, z_2 \in \widehat{\mathbb{C}}$ . When does  $\chi(z_1, z_2) = 2$  happen?

**1.3.2.** (i) Let  $l$  be any line in  $\mathbb{C}$ . We define  $\widehat{l} = l \cup \{\infty\}$  and call it *line in  $\widehat{\mathbb{C}}$* . We call *circle in  $\widehat{\mathbb{C}}$*  every circle in  $\mathbb{C}$ . Prove that stereographic projection maps circles in  $\widehat{\mathbb{C}}$  onto circles in  $\mathbb{S}^2$  which do

- not contain  $N$  (and conversely) and lines in  $\widehat{\mathbb{C}}$  onto circles in  $\mathbb{S}^2$  which contain  $N$  (and conversely).
- (ii) Find the images in  $\mathbb{S}^2$  through stereographic projection of the following subsets (or collections of subsets) of  $\widehat{\mathbb{C}}$ :  $\{z \mid |z| < 1\}$ ,  $\{z \mid |z| = 1\}$ ,  $\{z \mid |z| > 1\} \cup \{\infty\}$ ,  $\{z \mid \operatorname{Re} z > 0\}$ ,  $\{z \mid \operatorname{Re} z = 0\}$ ,  $\{z \mid \operatorname{Re} z < 0\}$ , the collection of lines containing a fixed point  $\neq \infty$ , the collection of circles with a fixed center, the collection of lines parallel to a fixed line, the collection of circles tangent to a fixed circle at a fixed point, the collection of circles containing two fixed points.
- (iii) Let  $z, w \in \widehat{\mathbb{C}}$  and let  $A, B \in \mathbb{S}^2$  be their images through stereographic projection. If  $z, w$  are symmetric with respect to a line  $\widehat{l}$  in  $\widehat{\mathbb{C}}$  which contains 0, which is the relative position of  $A, B$  with respect to the image of  $\widehat{l}$  in  $\mathbb{S}^2$ ? If  $w = \frac{1}{\bar{z}}$ , which is the relative position of  $A, B$  in  $\mathbb{S}^2$ ?
- (iv) Consider a set of the form  $P = \{z \in \widehat{\mathbb{C}} \mid \chi(z, z_0) = r\}$ , where  $z_0 \in \widehat{\mathbb{C}}$  and  $r > 0$ , i.e. a “circle” with respect to the chordal metric. If  $z_0 = \infty$ , prove that  $P$  is a circle in  $\widehat{\mathbb{C}}$ , i.e. in  $\mathbb{C}$ , and find its euclidean center and its euclidean radius. If  $z_0 \in \mathbb{C}$ , prove that  $P$  is either a circle in  $\widehat{\mathbb{C}}$ , i.e. in  $\mathbb{C}$ , and in this case find its euclidean center and its euclidean radius, or a line in  $\widehat{\mathbb{C}}$ .
- (v) If the lines  $\widehat{l}_1, \widehat{l}_2$  have angle  $\theta$  at their common point  $z \in \mathbb{C}$ , prove that their images through stereographic projection, i.e. two circles in  $\mathbb{S}^2$  containing the image  $A$  of  $z$  and the north pole  $N$ , have the same angle  $\theta$  at both  $A$  and  $N$ .

## Chapter 2

# Series and curvilinear integrals.

### 2.1 Series of numbers.

A **series of complex numbers** or, simply, **complex series** is an expression

$$z_1 + z_2 + \cdots + z_n + \cdots \quad \text{or} \quad \sum_{n=1}^{+\infty} z_n.$$

If all  $z_n$  are real, we talk about a **series of real numbers** or **real series**. The  $s_n = z_1 + \cdots + z_n$  are the **partial sums** of the series. We say that the series **converges** if the sequence  $(s_n)$  converges and then the limit  $s$  of  $(s_n)$  is called **sum** of the series and we write  $\sum_{n=1}^{+\infty} z_n = s$ . We say that the series **diverges** if  $(s_n)$  diverges. If  $(s_n)$  diverges to  $\infty$ , then we say that the series **diverges to  $\infty$**  and that  $\infty$  is the **sum** of the series and we write  $\sum_{n=1}^{+\infty} z_n = \infty$ .

We note that the sum of a complex series can be either a complex number or  $\infty$ . Only a real series can have sum equal to  $+\infty$  or  $-\infty$ . Therefore, when we write  $\sum_{n=1}^{+\infty} z_n = +\infty$  or  $-\infty$ , we accept that all  $z_n$  are real and that the series diverges to  $+\infty$  or  $-\infty$  as a real series. Of course, if a real series diverges to  $+\infty$  or  $-\infty$ , then as a complex series it diverges to  $\infty$ .

**Example 2.1.1.** We have  $\sum_{n=1}^{+\infty} c = 0$ , if  $c = 0$ , and  $\sum_{n=1}^{+\infty} c = \infty$ , if  $c \neq 0$ .

**Example 2.1.2.** To examine the geometric series  $\sum_{n=0}^{+\infty} z^n$ , we use the formula  $1 + z + \cdots + z^n = \frac{1-z^{n+1}}{1-z}$  for its partial sums, and we find that its sum is

$$\sum_{n=0}^{+\infty} z^n \begin{cases} = \frac{1}{1-z}, & \text{if } |z| < 1 \\ = \infty, & \text{if } |z| > 1 \text{ or } z = 1 \\ \text{it does not exist,} & \text{if } |z| = 1, z \neq 1 \end{cases}$$

The usual simple algebraic rules, which hold for real series, hold also for complex series. We mention them without proofs. The proofs for the complex case are identical with the proofs in the real case.

**Proposition 2.1.** *If  $\sum_{n=1}^{+\infty} z_n$  converges, then  $z_n \rightarrow 0$ .*

**Proposition 2.2.** *Provided that the right sides of the following formulas exist and that they are not indeterminate forms, we have  $\sum_{n=1}^{+\infty} (z_n + w_n) = \sum_{n=1}^{+\infty} z_n + \sum_{n=1}^{+\infty} w_n$ ,  $\sum_{n=1}^{+\infty} \lambda z_n = \lambda \sum_{n=1}^{+\infty} z_n$  and  $\sum_{n=1}^{+\infty} \overline{z_n} = \overline{\sum_{n=1}^{+\infty} z_n}$ .*

*Moreover, if  $z_n = x_n + iy_n$ , then  $\sum_{n=1}^{+\infty} z_n$  converges if and only if  $\sum_{n=1}^{+\infty} x_n$  and  $\sum_{n=1}^{+\infty} y_n$  converge, and  $\sum_{n=1}^{+\infty} z_n = \sum_{n=1}^{+\infty} x_n + i \sum_{n=1}^{+\infty} y_n$ .*

Regarding the *comparison theorems*, we may say that, since these are based on order relations which can be expressed only between real numbers, when we write  $\sum_{n=1}^{+\infty} z_n \leq \sum_{n=1}^{+\infty} w_n$  as a consequence of  $z_n \leq w_n$ , we accept that all  $z_n, w_n$  are real and then we just apply the well-known comparison theorems for real series.



**Cauchy criterion.** The series  $\sum_{n=1}^{+\infty} z_n$  converges if and only if for every  $\epsilon > 0$  there is  $n_0$  so that  $|\sum_{k=m+1}^n z_k| = |z_{m+1} + \cdots + z_n| < \epsilon$  for every  $m, n$  with  $n > m \geq n_0$ .

*Proof.* Let  $s_n = z_1 + \cdots + z_n$ . The series converges if and only if  $(s_n)$  converges or, equivalently, if and only if  $(s_n)$  is a Cauchy sequence. That  $(s_n)$  is a Cauchy sequence means that for every  $\epsilon > 0$  there is  $n_0$  so that  $|z_{m+1} + \cdots + z_n| = |s_n - s_m| < \epsilon$  for every  $n, m$  with  $n > m \geq n_0$ .  $\square$

We say that  $\sum_{n=1}^{+\infty} z_n$  **converges absolutely** if the (real) series  $\sum_{n=1}^{+\infty} |z_n|$  converges, i.e. if  $\sum_{n=1}^{+\infty} |z_n| < +\infty$ .

**Criterion of absolute convergence.** If  $\sum_{n=1}^{+\infty} z_n$  converges absolutely, then it converges and we have  $|\sum_{n=1}^{+\infty} z_n| \leq \sum_{n=1}^{+\infty} |z_n|$ .

*Proof.* Let  $\sum_{n=1}^{+\infty} |z_n|$  converge and take any  $\epsilon > 0$ . From the Cauchy criterion we have that there is  $n_0$  so that  $|z_{m+1}| + \cdots + |z_n| < \epsilon$  and hence  $|z_{m+1} + \cdots + z_n| < \epsilon$  for every  $m, n$  with  $n > m \geq n_0$ . The Cauchy criterion, again, implies that  $\sum_{n=1}^{+\infty} z_n$  converges.

Now we take the partial sums  $s_n = z_1 + \cdots + z_n$  and  $S_n = |z_1| + \cdots + |z_n|$ . We have  $|s_n| \leq S_n$  for all  $n$  and, taking the limit of this as  $n \rightarrow +\infty$ , we finish the proof.  $\square$

**Ratio test of d' Alembert.** Let  $z_n \neq 0$  for all  $n$ .

(i) If  $\overline{\lim} \left| \frac{z_{n+1}}{z_n} \right| < 1$ , then  $\sum_{n=1}^{+\infty} z_n$  converges absolutely.

(ii) If  $\underline{\lim} \left| \frac{z_{n+1}}{z_n} \right| > 1$ , then  $\sum_{n=1}^{+\infty} z_n$  diverges.

(iii) If  $\underline{\lim} \left| \frac{z_{n+1}}{z_n} \right| \leq 1 \leq \overline{\lim} \left| \frac{z_{n+1}}{z_n} \right|$ , then there is no general conclusion.

*Proof.* (i) Take  $a$  so that  $\overline{\lim} \left| \frac{z_{n+1}}{z_n} \right| < a < 1$ . Then there is  $n_0$  so that  $\left| \frac{z_{n+1}}{z_n} \right| \leq a$  for every  $n \geq n_0$ . Now, for every  $n \geq n_0$  we get  $|z_n| = \left| \frac{z_n}{z_{n-1}} \right| \left| \frac{z_{n-1}}{z_{n-2}} \right| \cdots \left| \frac{z_{n_0+1}}{z_{n_0}} \right| |z_{n_0}| \leq a^{n-n_0} |z_{n_0}| = c a^n$ , where  $c = |z_{n_0}|/a^{n_0}$ . Since  $0 \leq a < 1$ , the geometric series  $\sum_{n=1}^{+\infty} a^n$  converges and, by comparison,  $\sum_{n=1}^{+\infty} |z_n|$  also converges.

(ii) There is  $n_0$  so that  $\left| \frac{z_{n+1}}{z_n} \right| \geq 1$  for every  $n \geq n_0$ . Now, for every  $n \geq n_0 + 1$  we have  $|z_n| \geq |z_{n-1}| \geq \cdots \geq |z_{n_0}| > 0$ . This implies that  $z_n \not\rightarrow 0$  and  $\sum_{n=1}^{+\infty} z_n$  diverges.

(iii) For the series  $\sum_{n=1}^{+\infty} \frac{1}{n}$  and  $\sum_{n=1}^{+\infty} \frac{1}{n^2}$  we have that  $\left| \frac{1/(n+1)}{1/n} \right| \rightarrow 1$  and  $\left| \frac{1/(n+1)^2}{1/n^2} \right| \rightarrow 1$ . The first series diverges and the second converges.  $\square$

**Root test of Cauchy.** (i) If  $\overline{\lim} \sqrt[n]{|z_n|} < 1$ , then  $\sum_{n=1}^{+\infty} z_n$  converges absolutely.

(ii) If  $\overline{\lim} \sqrt[n]{|z_n|} > 1$ , then  $\sum_{n=1}^{+\infty} z_n$  diverges.

(iii) If  $\overline{\lim} \sqrt[n]{|z_n|} = 1$ , then there is no general conclusion.

*Proof.* (i) We consider any  $a$  such that  $\overline{\lim} \sqrt[n]{|z_n|} < a < 1$ . Then there is  $n_0$  so that  $\sqrt[n]{|z_n|} \leq a$  and hence  $|z_n| \leq a^n$  for every  $n \geq n_0$ . Since  $0 \leq a < 1$ , the geometric series  $\sum_{n=1}^{+\infty} a^n$  converges and, by comparison,  $\sum_{n=1}^{+\infty} |z_n|$  also converges.

(ii) We have  $\sqrt[n]{|z_n|} \geq 1$  for infinitely many  $n$ . Therefore,  $|z_n| \geq 1$  for infinitely many  $n$  and hence  $z_n \not\rightarrow 0$ . Thus,  $\sum_{n=1}^{+\infty} z_n$  diverges.

(iii) For the series  $\sum_{n=1}^{+\infty} \frac{1}{n}$  and  $\sum_{n=1}^{+\infty} \frac{1}{n^2}$  we have  $\sqrt[n]{|1/n|} \rightarrow 1$  and  $\sqrt[n]{|1/n^2|} \rightarrow 1$ . The first series diverges and the second converges.  $\square$

Applying the ratio test and the root test to specific series  $\sum_{n=1}^{+\infty} z_n$ , we find very often that the limits  $\lim_{n \rightarrow +\infty} \left| \frac{z_{n+1}}{z_n} \right|$  and  $\lim_{n \rightarrow +\infty} \sqrt[n]{|z_n|}$  exist. We know (and we used it in the proofs of parts (iii) of both tests) that in this case:  $\underline{\lim} = \overline{\lim} = \lim$ .

**Example 2.1.3.** To the series  $\sum_{n=1}^{+\infty} \frac{z^n}{n!}$  we apply the ratio test. If  $z = 0$ , the series obviously converges absolutely. If  $z \neq 0$ , then  $\left| \frac{z^{n+1}/(n+1)!}{z^n/n!} \right| = |z|/(n+1) \rightarrow 0 < 1$ . Hence the series converges absolutely for every  $z$ .

Now we apply the root test. We have  $\sqrt[n]{|z^n/n!|} = |z|/\sqrt[n]{n!} \rightarrow 0 < 1$  and we arrive at the same conclusion as before.

**Example 2.1.4.** We consider  $\sum_{n=1}^{+\infty} \frac{z^n}{n^2}$  and we apply the ratio test. If  $z = 0$ , the series obviously converges absolutely. If  $z \neq 0$ , then  $\left| \frac{z^{n+1}/(n+1)^2}{z^n/n^2} \right| \rightarrow |z|$ . Hence, if  $0 < |z| < 1$ , the series converges absolutely and, if  $|z| > 1$ , the series diverges.

Now we apply the root test. We have  $\sqrt[n]{|z^n/n^2|} \rightarrow |z|$ . Therefore, if  $|z| < 1$ , the series converges absolutely and, if  $|z| > 1$ , the series diverges.

If  $|z| = 1$ , none of the two tests applies. But we observe that  $\sum_{n=1}^{+\infty} \left| \frac{z^n}{n^2} \right| = \sum_{n=1}^{+\infty} \frac{1}{n^2} < +\infty$  in this case, and  $\sum_{n=1}^{+\infty} \frac{z^n}{n^2}$  converges absolutely.

Conclusion:  $\sum_{n=1}^{+\infty} \frac{z^n}{n^2}$  converges absolutely if  $|z| \leq 1$ , and diverges if  $|z| > 1$ .

**Lemma 2.1.** Let  $(a_n)$  and  $(z_n)$  be two sequences and  $s_n = z_1 + \dots + z_n$  for every  $n$ . Then we have

$$\sum_{k=m+1}^n a_k z_k = \sum_{k=m+1}^n (a_k - a_{k+1}) s_k + a_{n+1} s_n - a_{m+1} s_m$$

for every  $n, m$  with  $n > m$ . This is the **summation by parts formula due to Abel**.

*Proof.* We have

$$\begin{aligned} \sum_{k=m+1}^n a_k z_k &= \sum_{k=m+1}^n a_k (s_k - s_{k-1}) = \sum_{k=m+1}^n a_k s_k - \sum_{k=m}^{n-1} a_{k+1} s_k \\ &= \sum_{k=m+1}^n (a_k - a_{k+1}) s_k + a_{n+1} s_n - a_{m+1} s_m \end{aligned}$$

and the proof is complete.  $\square$

**Dirichlet test.** Let  $(a_n)$  and  $(z_n)$  be two sequences and  $s_n = z_1 + \dots + z_n$  for every  $n$ . If  $(a_n)$  is real and decreasing and  $a_n \rightarrow 0$  and if  $(s_n)$  is bounded, then  $\sum_{n=1}^{+\infty} a_n z_n$  converges.

*Proof.* There is  $M$  so that  $|s_n| \leq M$  for every  $n$ . Now, let  $\epsilon > 0$ . Since  $a_n \rightarrow 0$ , there is  $n_0$  so that  $0 \leq a_n < \frac{\epsilon}{2M+1}$  for every  $n \geq n_0$ . Then lemma 2.1 implies that, if  $n_0 \leq m < n$ ,

$$\begin{aligned} \left| \sum_{k=m+1}^n a_k z_k \right| &\leq \sum_{k=m+1}^n (a_k - a_{k+1}) |s_k| + a_{n+1} |s_n| + a_{m+1} |s_m| \\ &\leq \sum_{k=m+1}^n (a_k - a_{k+1}) M + a_{n+1} M + a_{m+1} M = 2a_{m+1} M < \epsilon. \end{aligned}$$

The criterion of Cauchy implies that  $\sum_{n=1}^{+\infty} a_n z_n$  converges.  $\square$

**Abel test.** Let  $(a_n)$ ,  $(z_n)$  be two sequences and  $s_n = z_1 + \dots + z_n$  for every  $n$ . If  $(a_n)$  is real and decreasing and bounded below and if  $(s_n)$  converges, i.e. if  $\sum_{n=1}^{+\infty} z_n$  converges, then  $\sum_{n=1}^{+\infty} a_n z_n$  converges.

*Proof.* Since  $(a_n)$  is real and decreasing and bounded below, there is  $a$  so that  $a_n \rightarrow a$ . We set  $a'_n = a_n - a$  and then  $(a'_n)$  is real and decreasing and  $a'_n \rightarrow 0$ . We also have that  $(s_n)$  is bounded and hence Dirichlet's test implies that the series  $\sum_{n=1}^{+\infty} a'_n z_n$  converges. Now, since  $\sum_{n=1}^{+\infty} z_n$  also converges, we find that  $\sum_{n=1}^{+\infty} a_n z_n = \sum_{n=1}^{+\infty} a'_n z_n + a \sum_{n=1}^{+\infty} z_n$  and hence  $\sum_{n=1}^{+\infty} a_n z_n$  converges.  $\square$

**Example 2.1.5.** If  $(a_n)$  is real and decreasing and  $a_n \rightarrow 0$ , then  $\sum_{n=0}^{+\infty} a_n z^n$  converges for every  $z$  with  $|z| \leq 1$ ,  $z \neq 1$ .

If  $|z| < 1$ , the result is immediate from the criterion of absolute convergence. Indeed, we have  $\sum_{n=0}^{+\infty} |a_n z^n| \leq a_1 \sum_{n=0}^{+\infty} |z|^n < +\infty$ . If  $|z| = 1$  and  $z \neq 1$ , we apply the Dirichlet test. To do this we prove that the partial sums  $s_n = 1 + z + z^2 + \dots + z^n$  are bounded:  $|s_n| = \frac{|1 - z^{n+1}|}{|1 - z|} \leq \frac{2}{|1 - z|}$  and  $\frac{2}{|1 - z|}$  is finite since  $z \neq 1$ .

**Example 2.1.6.** We consider  $\sum_{n=1}^{+\infty} \frac{z^n}{n}$ . As in example 2.1.4, the application of either the ratio test or the root test gives that the series converges absolutely if  $|z| < 1$  and diverges if  $|z| > 1$ . If  $|z| = 1$ , none of the two tests applies. If  $z = 1$ , the series becomes  $\sum_{n=1}^{+\infty} \frac{1}{n}$  and diverges. If  $|z| = 1$ ,  $z \neq 1$ , then  $\sum_{n=1}^{+\infty} \left| \frac{z^n}{n} \right| = \sum_{n=1}^{+\infty} \frac{1}{n} = +\infty$ , and  $\sum_{n=1}^{+\infty} \frac{z^n}{n}$  does not converge absolutely.

But the series is a particular instance of the series in the previous example and hence converges if  $|z| = 1, z \neq 1$ . In general, when a series is convergent but not absolutely convergent we say that it is **conditionally convergent**.

Conclusion:  $\sum_{n=1}^{+\infty} \frac{z^n}{n}$  converges absolutely if  $|z| < 1$ , diverges if  $|z| > 1$  or  $z = 1$ , and converges conditionally if  $|z| = 1, z \neq 1$ .

Let  $\sum_{n=0}^{+\infty} a_n$  and  $\sum_{n=0}^{+\infty} b_n$  be two series. If  $c_n = a_0 b_n + a_1 b_{n-1} + \cdots + a_{n-1} b_1 + a_n b_0$  for every  $n \geq 0$ , then the series

$$\sum_{n=0}^{+\infty} c_n = \sum_{n=0}^{+\infty} (a_0 b_n + a_1 b_{n-1} + \cdots + a_{n-1} b_1 + a_n b_0)$$

is called **Cauchy product** of the two series.

**Proposition 2.3.** *If the series  $\sum_{n=0}^{+\infty} a_n$  and  $\sum_{n=0}^{+\infty} b_n$  converge absolutely, then their Cauchy product  $\sum_{n=0}^{+\infty} c_n$  converges absolutely. Moreover, we have  $\sum_{n=0}^{+\infty} c_n = \sum_{n=0}^{+\infty} a_n \sum_{n=0}^{+\infty} b_n$  for the sums of the three series.*

*Proof.* We have  $|c_n| \leq |a_0||b_n| + |a_1||b_{n-1}| + \cdots + |a_{n-1}||b_1| + |a_n||b_0|$ . Hence, if  $S = \sum_{n=0}^{+\infty} |a_n| < +\infty$  and  $T = \sum_{n=0}^{+\infty} |b_n| < +\infty$ , then

$$\sum_{n=0}^N |c_n| \leq \sum_{n=0}^N (\sum_{k=0}^n |a_k||b_{n-k}|) = \sum_{k=0}^N |a_k| (\sum_{n=k}^N |b_{n-k}|) \leq \sum_{k=0}^N |a_k| T \leq ST$$

for every  $N$ . Thus,  $\sum_{n=0}^{+\infty} |c_n| \leq ST < +\infty$  and  $\sum_{n=0}^{+\infty} c_n$  converges absolutely.

Now, let  $s = \sum_{n=0}^{+\infty} a_n$ ,  $t = \sum_{n=0}^{+\infty} b_n$  and  $u = \sum_{n=0}^{+\infty} c_n$ . Moreover, let  $s_n = a_0 + \cdots + a_n$ ,  $t_n = b_0 + \cdots + b_n$  and  $u_n = c_0 + \cdots + c_n$  be the partial sums of the three series and also  $S_n = |a_0| + \cdots + |a_n|$ ,  $T_n = |b_0| + \cdots + |b_n|$ . Then

$$\begin{aligned} u_N &= \sum_{n=0}^N c_n = \sum_{n=0}^N (\sum_{k=0}^n a_k b_{n-k}) = \sum_{k=0}^N a_k (\sum_{n=k}^N b_{n-k}) = \sum_{k=0}^N a_k (\sum_{m=0}^{N-k} b_m) \\ &= \sum_{k=0}^N a_k t_{N-k} \end{aligned}$$

and hence  $s_N t_N - u_N = \sum_{k=0}^N a_k (t_N - t_{N-k})$ . We take  $p = [\frac{N}{2}]$  and we get

$$s_N t_N - u_N = \sum_{k=0}^p a_k (t_N - t_{N-k}) + \sum_{k=p+1}^N a_k (t_N - t_{N-k}). \quad (2.1)$$

If  $0 \leq k \leq p$ , then  $N - k \geq N - p \geq p$  and hence

$$|t_N - t_{N-k}| = |\sum_{m=N-k+1}^N b_m| \leq \sum_{m=N-k+1}^{+\infty} |b_m| \leq \sum_{m=p+1}^{+\infty} |b_m| = T - T_p. \quad (2.2)$$

If  $p+1 \leq k \leq N$ , then

$$|t_N - t_{N-k}| = |\sum_{m=N-k+1}^N b_m| \leq \sum_{m=N-k+1}^{+\infty} |b_m| \leq T. \quad (2.3)$$

Now, (2.1), (2.2) and (2.3) imply

$$\begin{aligned} |s_N t_N - u_N| &\leq \sum_{k=0}^p |a_k| |t_N - t_{N-k}| + \sum_{k=p+1}^N |a_k| |t_N - t_{N-k}| \\ &\leq (T - T_p) \sum_{k=0}^p |a_k| + T \sum_{k=p+1}^N |a_k| \leq (T - T_p) S + T(S - S_p). \end{aligned}$$

Now,  $N \rightarrow +\infty$  implies  $p \rightarrow +\infty$  and hence  $S_p \rightarrow S$  and  $T_p \rightarrow T$ . Therefore,  $s_N t_N - u_N \rightarrow 0$  when  $N \rightarrow +\infty$  and we conclude that  $u = st$ .  $\square$

### Exercises.

**2.1.1.** Which of the series  $\sum_{n=1}^{+\infty} (\frac{1}{n} + \frac{i}{n^2})$ ,  $\sum_{n=1}^{+\infty} (\frac{n}{2^n} + \frac{i}{n^3})$ ,  $\sum_{n=1}^{+\infty} \frac{1+i^n}{n^2}$ ,  $\sum_{n=1}^{+\infty} \frac{1}{2+i^n}$ ,  $\sum_{n=1}^{+\infty} \frac{1}{n+i}$ ,  $\sum_{n=1}^{+\infty} \frac{1}{n^2+in}$  converge?

**2.1.2.** Find the sum of the series  $\sum_{n=1}^{+\infty} n(-1)^{n-1}$  if we consider it as a complex series and also if we consider it as a real series.

**2.1.3.** Apply the ratio test whenever possible:  $\sum_{n=1}^{+\infty} n^3 i^n$ ,  $\sum_{n=1}^{+\infty} \frac{n!}{i^n}$ ,  $\sum_{n=1}^{+\infty} \frac{(1+i)^n}{n!}$ ,  $\sum_{n=1}^{+\infty} \frac{(2i)^n n!}{n^n}$ ,  $\sum_{n=1}^{+\infty} \frac{(2+i)^n n!}{n^n}$ ,  $\sum_{n=1}^{+\infty} \frac{e^n n!}{n^n}$ ,  $\sum_{n=1}^{+\infty} \frac{(n!)^2}{(2n)!}$ ,  $\sum_{n=1}^{+\infty} \frac{(4i)^n (n!)^2}{(2n)!}$ ,  $\sum_{n=1}^{+\infty} \frac{(3+i)(6+i)(9+i)\cdots(3n+i)}{(3+4i)(3+8i)(3+12i)\cdots(3+4ni)}$ .

Apply the root test whenever possible:  $\sum_{n=1}^{+\infty} n^n i^n$ ,  $\sum_{n=1}^{+\infty} (\frac{n+i}{2n-i})^n$ ,  $\sum_{n=1}^{+\infty} (\frac{n+i}{n-i})^{2n}$ ,  $\sum_{n=1}^{+\infty} \frac{n^3}{(1+2i)^n}$ ,  $\sum_{n=1}^{+\infty} n^3(1-i)^n$ ,  $\sum_{n=1}^{+\infty} \frac{(2+3i)^n}{n^n}$ ,  $\sum_{n=1}^{+\infty} \frac{n+i}{(\sqrt[n]{n+i})^n}$ .

**2.1.4.** If  $\sum_{n=1}^{+\infty} |z_n| < +\infty$ , prove that  $\sum_{n=1}^{+\infty} z_n(\cos n\theta + i \sin n\theta)$  converges.

**2.1.5.** Let  $z_n = x_n + iy_n$  for all  $n$ . Prove that  $\sum_{n=1}^{+\infty} z_n$  converges absolutely if and only if  $\sum_{n=1}^{+\infty} x_n, \sum_{n=1}^{+\infty} y_n$  converge absolutely.

**2.1.6.** Let  $|a_n|r^n \leq Mn^k$  for all  $n$ . Prove that  $\sum_{n=1}^{+\infty} a_n z^n$  converges for every  $z$  with  $|z| < r$ .

**2.1.7.** Find all  $z$  for which  $\sum_{n=1}^{+\infty} \frac{z^n}{2+z^n}$  converges.

**2.1.8.** Let  $0 \leq \theta_0 < \frac{\pi}{2}$  and  $\text{Arg } z_n \in [-\theta_0, \theta_0]$  for every  $n$ . Prove that  $\sum_{n=1}^{+\infty} z_n$  converges if and only if it converges absolutely. Prove that  $\sum_{n=1}^{+\infty} z_n = \infty$  if and only if  $\sum_{n=1}^{+\infty} |z_n| = +\infty$ .

**2.1.9.** Find a series  $\sum_{n=1}^{+\infty} z_n$  which converges and is such that  $\sum_{n=1}^{+\infty} z_n^2$  diverges.

**2.1.10.** Check the conditional convergence and the absolute convergence of the series:  $\sum_{n=1}^{+\infty} \frac{i^n}{n}$ ,  $\sum_{n=2}^{+\infty} \frac{i^n}{n \log n}$ ,  $\sum_{n=2}^{+\infty} \frac{i^n}{n(\log n)^2}$ ,  $\sum_{n=1}^{+\infty} i^{n-1} \sin \frac{1}{n}$ ,  $\sum_{n=1}^{+\infty} i^{n-1} (1 - \cos \frac{1}{n})$ .

**2.1.11.** Let  $s_n = z_1 + \cdots + z_n$  for all  $n$ . If  $(a_{n+1}s_n)$  converges and if  $\sum_{n=1}^{+\infty} (a_n - a_{n+1})s_n$  converges, prove that  $\sum_{n=1}^{+\infty} a_n z_n$  converges. In particular: if  $(s_n)$  is bounded, if  $a_n \rightarrow 0$  and if  $\sum_{n=1}^{+\infty} |a_n - a_{n+1}| < +\infty$ , prove that  $\sum_{n=1}^{+\infty} a_n z_n$  converges.

What is the relation of all these with the tests of Dirichlet and Abel?

**2.1.12.** (i) If  $\sum_{n=0}^{+\infty} a_n$  converges and  $\sum_{n=0}^{+\infty} b_n$  converges absolutely, prove that their Cauchy product  $\sum_{n=0}^{+\infty} c_n$  converges and that  $\sum_{n=0}^{+\infty} c_n = \sum_{n=0}^{+\infty} a_n \sum_{n=0}^{+\infty} b_n$ .

(ii) Prove that the series  $\sum_{n=1}^{+\infty} \frac{(-1)^{n-1}}{\sqrt{n}}$  converges but that the Cauchy product of this series with itself does not converge.

## 2.2 Sequences and series of functions.

Let  $A$  be any nonempty set (not necessarily a subset of  $\mathbb{C}$ ),  $(f_n)$  be a sequence of bounded complex functions defined in  $A$  and  $f$  be a bounded complex function defined also in  $A$ . We say that the sequence  $(f_n)$  **converges** to  $f$  **uniformly** in  $A$  if

$$\sup_{a \in A} |f_n(a) - f(a)| \rightarrow 0.$$

In other words,  $f_n \rightarrow f$  uniformly in  $A$  if for every  $\epsilon > 0$  there is  $n_0$  so that  $|f_n(a) - f(a)| \leq \epsilon$  for every  $n \geq n_0$  and every  $a \in A$ .

It is easy to see that, if  $f_n \rightarrow f$  uniformly in  $A$ , then  $f_n(a) \rightarrow f(a)$  for every  $a \in A$ , i.e.  $(f_n)$  converges to  $f$  **pointwise** in  $A$ . Indeed, for every  $a' \in A$  we have  $0 \leq |f_n(a') - f(a')| \leq \sup_{a \in A} |f_n(a) - f(a)| \rightarrow 0$ .

From the notion of uniform convergence of a sequence of functions we move to the notion of uniform convergence of a series of functions in the obvious way, i.e. through the sequence of partial sums of the series. We write  $s_n = f_1 + \cdots + f_n$  and we say that the series of functions  $\sum_{n=1}^{+\infty} f_n$  **converges** to its **sum**  $s$  **uniformly** in  $A$  if  $s_n \rightarrow s$  uniformly in  $A$ . In this case we write  $\sum_{n=1}^{+\infty} f_n = s$  uniformly in  $A$ .

As in the case of a sequence of functions, we have that, if  $\sum_{n=1}^{+\infty} f_n = s$  uniformly in  $A$ , then  $\sum_{n=1}^{+\infty} f_n(a) = s(a)$  for every  $a \in A$ , i.e.  $\sum_{n=1}^{+\infty} f_n = s$  **pointwise** in  $A$ .

For the next two results we assume that  $A$  is a subset of a metric space.

**Proposition 2.4.** Let  $f_n \rightarrow f$  uniformly in  $A$  and let  $a_0 \in A$ . If every  $f_n$  is continuous at  $a_0$ , then  $f$  is continuous at  $a_0$ . In particular, if every  $f_n$  is continuous in  $A$ , then  $f$  is continuous in  $A$ .

*Proof.* Let  $\epsilon > 0$ . Then there is  $n_0$  so that  $|f_n(a) - f(a)| \leq \frac{\epsilon}{3}$  for every  $n \geq n_0$  and every  $a \in A$ . In particular, we have  $|f_{n_0}(a) - f(a)| \leq \frac{\epsilon}{3}$  for every  $a \in A$ . Now,  $f_{n_0}$  is continuous at  $a_0$  and so, if  $a \in A$  is close to  $a_0$ , then we have  $|f_{n_0}(a) - f_{n_0}(a_0)| \leq \frac{\epsilon}{3}$  and hence

$$|f(a) - f(a_0)| \leq |f(a) - f_{n_0}(a)| + |f_{n_0}(a) - f_{n_0}(a_0)| + |f_{n_0}(a_0) - f(a_0)| \leq \frac{\epsilon}{3} + \frac{\epsilon}{3} + \frac{\epsilon}{3} = \epsilon.$$

Therefore,  $f$  is continuous at  $a_0$ .  $\square$

**Proposition 2.5.** Let  $\sum_{n=1}^{+\infty} f_n = s$  uniformly in  $A$  and let  $a_0 \in A$ . If every  $f_n$  is continuous at  $a_0$ , then  $s$  is continuous at  $a_0$ . In particular, if every  $f_n$  is continuous in  $A$ , then  $s$  is continuous in  $A$ .

*Proof.* We consider the partial sums  $s_n = f_1 + \cdots + f_n$ . Then every  $s_n$  is continuous at  $a_0$  and proposition 2.4 implies that  $s$  is continuous at  $a_0$ .  $\square$

Finally, we have a basic criterion for uniform convergence of a series of functions.

**Weierstrass test.** Let  $|f_n(a)| \leq M_n$  for every  $n$  and every  $a \in A$ . If the series (of non-negative terms)  $\sum_{n=1}^{+\infty} M_n$  converges, i.e. if  $\sum_{n=1}^{+\infty} M_n < +\infty$ , then  $\sum_{n=1}^{+\infty} f_n$  converges uniformly in  $A$ .

*Proof.* For every  $a \in A$  we have  $\sum_{n=1}^{+\infty} |f_n(a)| \leq \sum_{n=1}^{+\infty} M_n < +\infty$  and hence  $\sum_{n=1}^{+\infty} f_n(a)$  converges (as a series of complex numbers). We define  $s(a) = \sum_{n=1}^{+\infty} f_n(a)$  for every  $a \in A$ . Now we consider the partial sums  $s_n = f_1 + \cdots + f_n$  and then for every  $a \in A$  we have

$$\begin{aligned} |s_n(a) - s(a)| &= \left| \sum_{k=1}^n f_k(a) - \sum_{k=1}^{+\infty} f_k(a) \right| = \left| \sum_{k=n+1}^{+\infty} f_k(a) \right| \leq \sum_{k=n+1}^{+\infty} |f_k(a)| \\ &\leq \sum_{k=n+1}^{+\infty} M_k. \end{aligned}$$

This implies that  $\sup_{a \in A} |s_n(a) - s(a)| \leq \sum_{k=n+1}^{+\infty} M_k \rightarrow 0$  when  $n \rightarrow +\infty$  and hence  $s_n \rightarrow s$  uniformly in  $A$ .  $\square$

### Exercises.

**2.2.1.** Prove that  $\sum_{n=-\infty}^{+\infty} \frac{1}{(z+n)^2}$  converges for every  $z \in \mathbb{C} \setminus \mathbb{Z}$  and uniformly in every compact set  $K \subseteq \mathbb{C} \setminus \mathbb{Z}$ .

**2.2.2.** (i) If  $K \subseteq \mathbb{C} \setminus \mathbb{T}$  is compact, prove that there is  $r$  with  $0 < r < 1$  ( $r$  depends on  $K$ ) so that for every  $z \in K$  either  $|z| \leq r$  or  $|z| \geq \frac{1}{r}$  holds.

(ii) Prove that  $\sum_{n=0}^{+\infty} \frac{z^n}{z^{2n+1}}$  converges uniformly in every compact  $K \subseteq \mathbb{C} \setminus \mathbb{T}$ .

**2.2.3.** (i) If  $\operatorname{Re} z > -\frac{1}{2}$ , prove that  $\left| \frac{z}{z+1} \right| < 1$ . If  $K \subseteq \{z \mid \operatorname{Re} z > -\frac{1}{2}\}$  is compact, prove that there is  $r$  with  $0 < r < 1$  ( $r$  depends on  $K$ ) so that  $\left| \frac{z}{z+1} \right| \leq r$  for every  $z \in K$ .

(ii) Prove that  $\sum_{n=0}^{+\infty} \left(\frac{z}{z+1}\right)^n$  converges for every  $z$  in the halfplane  $\{z \mid \operatorname{Re} z > -\frac{1}{2}\}$  and uniformly in every compact subset of this halfplane.

## 2.3 Curvilinear integrals.

We shall first extend the notion of integral of a *real* function over an interval to the notion of integral of a *complex* function over an interval.

Let  $f$  be a complex function defined in the interval  $[a, b]$  and let  $u = \operatorname{Re} f$  and  $v = \operatorname{Im} f$  be the real and imaginary parts of  $f$ . We say that  $f$  is **(Riemann) integrable** over  $[a, b]$  if  $u, v$  are both (Riemann) integrable over  $[a, b]$  and in this case we define the **(Riemann) integral** of  $f$  over  $[a, b]$  to be

$$\int_a^b f(t) dt = \int_a^b u(t) dt + i \int_a^b v(t) dt. \quad (2.4)$$

Since the numbers  $\int_a^b u(t) dt$  and  $\int_a^b v(t) dt$  are real, we see that

$$\operatorname{Re} \int_a^b f(t) dt = \int_a^b \operatorname{Re} f(t) dt, \quad \operatorname{Im} \int_a^b f(t) dt = \int_a^b \operatorname{Im} f(t) dt.$$

Now let us take any subdivision  $\Delta = \{t_0, \dots, t_n\}$  of  $[a, b]$  and any choice  $\Xi = \{\xi_1, \dots, \xi_n\}$  of intermediate points  $\xi_k \in [t_{k-1}, t_k]$  and the corresponding Riemann sum  $\sum_{k=1}^n f(\xi_k)(t_k - t_{k-1})$ . If  $w(\Delta) = \max_{1 \leq k \leq n} (t_k - t_{k-1})$  is the width of the subdivision  $\Delta$ , then we know that

$$\lim_{w(\Delta) \rightarrow 0} \sum_{k=1}^n u(\xi_k)(t_k - t_{k-1}) = \int_a^b u(t) dt$$

$$\lim_{w(\Delta) \rightarrow 0} \sum_{k=1}^n v(\xi_k)(t_k - t_{k-1}) = \int_a^b v(t) dt.$$

Multiplying the second relation with  $i$ , adding and using (2.4), we find

$$\lim_{w(\Delta) \rightarrow 0} \sum_{k=1}^n f(\xi_k)(t_k - t_{k-1}) = \int_a^b f(t) dt.$$

**Example 2.3.1.** If  $f$  is piecewise-continuous in  $[a, b]$ , then  $u = \operatorname{Re} f$  and  $v = \operatorname{Im} f$  are also piecewise-continuous in  $[a, b]$ . Hence  $u, v$  are integrable, and so  $f$  is also integrable over  $[a, b]$ .

The following propositions are analogous to similar well-known propositions about integrals of real functions and can be proved easily by the reader. One should decompose every complex function into its real and imaginary parts and use the analogous properties for real functions together with (2.4).

**Proposition 2.6.** Let  $f_1, f_2$  be integrable over  $[a, b]$  and  $\lambda_1, \lambda_2 \in \mathbb{C}$ . Then  $\lambda_1 f_1 + \lambda_2 f_2$  is integrable over  $[a, b]$  and  $\int_a^b (\lambda_1 f_1(t) + \lambda_2 f_2(t)) dt = \lambda_1 \int_a^b f_1(t) dt + \lambda_2 \int_a^b f_2(t) dt$ .

**Proposition 2.7.** Let  $a < b < c$ . If  $f$  is integrable over  $[a, b]$  and over  $[b, c]$ , then  $f$  is integrable over  $[a, c]$  and  $\int_a^c f(t) dt = \int_a^b f(t) dt + \int_b^c f(t) dt$ .

**Proposition 2.8.** If  $f_1, f_2$  are integrable over  $[a, b]$ , then  $f_1 f_2$  is integrable over  $[a, b]$ .

The proof of the next proposition is not entirely trivial.

**Proposition 2.9.** If  $f$  is integrable over  $[a, b]$ , then  $|f|$  is integrable over  $[a, b]$  and  $|\int_a^b f(t) dt| \leq \int_a^b |f(t)| dt$ .

*Proof.* Let  $u = \operatorname{Re} f$ ,  $v = \operatorname{Im} f$ . Then  $u, v$  are integrable over  $[a, b]$  hence  $|f| = \sqrt{u^2 + v^2}$  is integrable over  $[a, b]$ . Now we have two cases.

(i) Let  $\int_a^b f(t) dt = 0$ . Then  $|\int_a^b f(t) dt| \leq \int_a^b |f(t)| dt$  is clearly true.

(ii) Let  $\int_a^b f(t) dt \neq 0$ . We take any element  $\theta$  of the argument of the number  $\int_a^b f(t) dt$  and we set  $z = \cos \theta + i \sin \theta$ . Now,  $|\int_a^b f(t) dt| = \bar{z} \int_a^b f(t) dt = \int_a^b (\bar{z} f(t)) dt$ . The left side of this equality is real and hence its right side is also real and thus equal to its real part! Hence

$$|\int_a^b f(t) dt| = \operatorname{Re} \int_a^b (\bar{z} f(t)) dt = \int_a^b \operatorname{Re}(\bar{z} f(t)) dt \leq \int_a^b |\bar{z} f(t)| dt = \int_a^b |f(t)| dt$$

since  $\operatorname{Re}(\bar{z} f(t)) \leq |\bar{z} f(t)|$  for every  $t \in [a, b]$ . □

We recall that every *continuous* complex function  $\gamma : [a, b] \rightarrow \mathbb{C}$ , where  $[a, b]$  is any interval, is called *curve* in the complex plane.

The set of the values of a curve  $\gamma$ , i.e. the set  $\gamma^* = \{\gamma(t) \mid t \in [a, b]\} \subseteq \mathbb{C}$  is the *trajectory* of the curve and it is a compact and connected subset of  $\mathbb{C}$ , since  $\gamma$  is continuous and  $[a, b]$  is compact and connected. The points  $\gamma(a)$  and  $\gamma(b)$  are the endpoints, the initial and the final endpoint, respectively, of the curve.

The variable  $t \in [a, b]$  is the *parameter* and  $[a, b]$  is the *parametric interval* of the curve. When the parameter  $t$  increases in  $[a, b]$ , the variable point  $\gamma(t)$  moves on the trajectory  $\gamma^*$  in a definite

direction (from the initial to the final endpoint) which is the so-called *direction* of the curve. To be more precise, the sense of *direction* is realized as follows: when  $a \leq t_1 < t_2 \leq b$ , then we say that  $\gamma(t_1)$  is *before*  $\gamma(t_2)$  and that  $\gamma(t_2)$  is *after*  $\gamma(t_1)$  in the trajectory.

Finally,

$$z = \gamma(t), \quad t \in [a, b],$$

is the *parametric equation* of the curve  $\gamma$ .

If the endpoints of the curve  $\gamma$  coincide, i.e.  $\gamma(a) = \gamma(b)$ , then we say that the curve is *closed*.

If  $\gamma(t) \in A$  for all  $t \in [a, b]$ , i.e. if  $\gamma^* \subseteq A$ , then we say that *the curve is in A*.

The term *curve* for the continuous function  $\gamma$  is justified by the fact that the shape of its trajectory  $\gamma^*$  is, usually, what in everyday language we call *curve in the plane*. Sometimes we use the term *curve* for the trajectory  $\gamma^*$  even though this is not typically correct. The reason is that there are cases of different curves  $\gamma_1, \gamma_2$  with the same trajectory  $\gamma_1^* = \gamma_2^*$ .

**Example 2.3.2.** If  $z_0, z_1 \in \mathbb{C}$ , then the parametric equation  $z = \gamma(t) = \frac{t-a}{b-a}z_1 + \frac{b-t}{b-a}z_0, t \in [a, b]$ , defines a curve  $\gamma$  whose trajectory  $\gamma^*$  is the linear segment  $[z_0, z_1]$ . Its initial and final endpoints are  $z_0$  and  $z_1$ , respectively, and its direction is from  $z_0$  to  $z_1$ . The *same* linear segment  $[z_0, z_1]$  is the trajectory of another curve  $\gamma$  with parametric equation  $z = \gamma(t) = tz_1 + (1-t)z_0, t \in [0, 1]$ .

**Example 2.3.3.** If  $r > 0$ , then the parametric equation  $z = \gamma(t) = z_0 + r(\cos t + i \sin t), t \in [0, 2\pi]$ , defines a closed curve  $\gamma$  whose trajectory  $\gamma^*$  is the circle  $C_{z_0}(r)$ . The direction of this curve is the so-called *positive direction of rotation* around  $z_0$ : the counterclockwise rotation.

If we consider the curve  $\gamma$  with parametric equation  $z = \gamma(t) = z_0 + r(\cos(2t) + i \sin(2t)), t \in [0, 2\pi]$ , then we get a different curve. But the trajectories of the two curves coincide: the circle  $C_{z_0}(r)$ . The direction of the two curves is the same: the positive direction of rotation around  $z_0$ . But the first curve goes around  $z_0$  only once, while the second curve goes around  $z_0$  twice.

Let  $\gamma : [a, b] \rightarrow \mathbb{C}$  be a curve and let  $x = \operatorname{Re} \gamma$  and  $y = \operatorname{Im} \gamma$  be the real and imaginary parts of  $\gamma$ , i.e.  $\gamma(t) = x(t) + iy(t) = (x(t), y(t))$  for all  $t \in [a, b]$ . If  $\gamma$  is differentiable at  $t_0 \in [a, b]$  or, equivalently, if  $x, y$  are differentiable at  $t_0$ , then  $\gamma'(t_0) = x'(t_0) + iy'(t_0) = (x'(t_0), y'(t_0))$  is the *tangent vector* of the trajectory  $\gamma^*$  at its point  $\gamma(t_0)$ . If  $\gamma'(t_0) \neq 0$ , then the vector  $\gamma'(t_0)$  determines the *tangent line* of the trajectory  $\gamma^*$  at its point  $\gamma(t_0)$  and its direction is the same as the direction of the curve. Strictly speaking, at its endpoints,  $\gamma(a), \gamma(b)$ , the curve can only have *tangent halflines*; not tangent lines. If  $t_0 = a$  and  $\gamma'(a) \neq 0$ , then the vector  $\gamma'(a)$  determines the *tangent halfline* of the trajectory at the endpoint  $\gamma(a)$  with direction coinciding with the direction of the curve. If  $t_0 = b$  and  $\gamma'(b) \neq 0$ , then the vector  $-\gamma'(b)$  determines the *tangent halfline* of the trajectory at the endpoint  $\gamma(b)$  with direction opposite to the direction of the curve. If at some  $t_0 \in (a, b)$  the one-sided derivatives  $\gamma'_-(t_0) \neq 0$  and  $\gamma'_+(t_0) \neq 0$  exist but they are not equal, then the tangent halflines of the trajectory at its point  $\gamma(t_0)$  may not be opposite and so there may be no tangent line of the trajectory at this point: one of the halflines is determined by  $\gamma'_+(t_0)$  and the other by  $-\gamma'_-(t_0)$ .

We know that, if the curve  $\gamma : [a, b] \rightarrow \mathbb{C}$  is *continuously differentiable* or *smooth*, i.e. if  $\gamma' : [a, b] \rightarrow \mathbb{C}$  is continuous in  $[a, b]$ , then the *length* of the curve, denoted  $l(\gamma)$ , is equal to

$$l(\gamma) = \int_a^b |\gamma'(t)| dt. \quad (2.5)$$

**Example 2.3.4.** The curve  $\gamma$  with parametric equation  $z = \gamma(t) = \frac{b-t}{b-a}z_0 + \frac{t-a}{b-a}z_1, t \in [a, b]$ , has length  $l(\gamma) = \int_a^b |\gamma'(t)| dt = \int_a^b \left| \frac{z_1 - z_0}{b-a} \right| dt = \left| \frac{z_1 - z_0}{b-a} \right| \int_a^b dt = |z_1 - z_0|$ .

**Example 2.3.5.** If  $r > 0$  the curve  $\gamma$  with parametric equation  $z = \gamma(t) = z_0 + r(\cos t + i \sin t), t \in [0, 2\pi]$ , has length  $l(\gamma) = \int_0^{2\pi} |\gamma'(t)| dt = \int_0^{2\pi} |r(-\sin t + i \cos t)| dt = \int_0^{2\pi} r dt = 2\pi r$ .

The same formula (2.5) gives the length of the curve  $\gamma$  if this is *piecewise continuously differentiable* or *piecewise smooth*. This means that there is a subdivision  $a = t_0 < t_1 < \dots < t_{n-1} <$

$t_n = b$  of the parametric interval  $[a, b]$  so that the restriction of  $\gamma$  in every  $[t_{k-1}, t_k]$  is continuously differentiable. (Strictly speaking, at the division points  $t_k$  the derivative of  $\gamma$  may not exist; the two one-sided derivatives should exist and be finite at these points.)

Now let  $\gamma_1 : [a, b] \rightarrow \mathbb{C}$  be a curve. We consider any  $\sigma : [c, d] \rightarrow [a, b]$  which is one-to-one in the interval  $[c, d]$  and onto  $[a, b]$ , has continuous derivative in  $[c, d]$  and has  $\sigma'(s) > 0$  for every  $s \in [c, d]$ . Thus,  $\sigma$  is strictly increasing in  $[c, d]$  and  $\sigma(c) = a, \sigma(d) = b$ . Every such  $\sigma$  is called *change of parameter*. Then  $\gamma_2 = \gamma_1 \circ \sigma : [c, d] \rightarrow \mathbb{C}$  is continuous in  $[c, d]$  and hence it is a new curve. We say that  $\gamma_2$  is a *reparametrization* of  $\gamma_1$ : the parameter of  $\gamma_1$  is  $t \in [a, b]$  and the parameter of  $\gamma_2$  is  $s \in [c, d]$ . The curves  $\gamma_1, \gamma_2$  have the same trajectory, the same endpoints and the same direction. Since  $\sigma'$  is continuous and  $> 0$ , the two curves are simultaneously (piecewise) smooth and, in this case, their lengths are equal:

$$\begin{aligned} l(\gamma_2) &= \int_c^d |\gamma_2'(s)| ds = \int_c^d |\gamma_1'(\sigma(s))| |\sigma'(s)| ds = \int_c^d |\gamma_1'(\sigma(s))| \sigma'(s) ds \\ &= \int_a^b |\gamma_1'(t)| dt = l(\gamma_1). \end{aligned}$$

We may define the following relation between curves:  $\gamma_1 \sim \gamma_2$  if  $\gamma_2$  is a reparametrization of  $\gamma_1$ . It is not difficult to prove that this relation between curves is an *equivalence relation*, i.e. it satisfies the following three properties:

- (i)  $\gamma \sim \gamma$ .
- (ii)  $\gamma_1 \sim \gamma_2 \Rightarrow \gamma_2 \sim \gamma_1$ .
- (iii)  $\gamma_1 \sim \gamma_2, \gamma_2 \sim \gamma_3 \Rightarrow \gamma_1 \sim \gamma_3$ .

Indeed: (i) Let  $\gamma : [a, b] \rightarrow \mathbb{C}$  be any curve. We consider the change of parameter  $id : [a, b] \rightarrow [a, b]$ , defined by  $id(t) = t$ , and then  $\gamma = \gamma \circ id : [a, b] \rightarrow \mathbb{C}$ . Thus,  $\gamma \sim \gamma$ . (ii) Let  $\gamma_1 \sim \gamma_2$ . Then  $\gamma_2 = \gamma_1 \circ \sigma$  where  $\sigma : [c, d] \rightarrow [a, b]$  is a change of parameter. But then  $\sigma^{-1} : [a, b] \rightarrow [c, d]$  is also a change of parameter and  $\gamma_1 = \gamma_2 \circ \sigma^{-1}$ . Therefore  $\gamma_2 \sim \gamma_1$ . (iii) Let  $\gamma_1 \sim \gamma_2$  and  $\gamma_2 \sim \gamma_3$ . Then  $\gamma_2 = \gamma_1 \circ \sigma$  and  $\gamma_3 = \gamma_2 \circ \tau$ , where  $\sigma : [c, d] \rightarrow [a, b]$  and  $\tau : [e, f] \rightarrow [c, d]$  are changes of parameter. But then  $\chi = \sigma \circ \tau : [e, f] \rightarrow [a, b]$  is a change of parameter and  $\gamma_3 = \gamma_2 \circ \tau = (\gamma_1 \circ \sigma) \circ \tau = \gamma_1 \circ \chi$ . Therefore  $\gamma_1 \sim \gamma_3$ .

It is of some value to note that if we have a curve  $\gamma$  with parametric interval  $[a, b]$  and if we are given an arbitrary interval  $[c, d]$ , then there is a reparametrization of  $\gamma$  with parametric interval  $[c, d]$  instead of  $[a, b]$ . We can do this if we can find an appropriate change of parameter  $\sigma : [c, d] \rightarrow [a, b]$ . There are many such  $\sigma$ , but a simple one is  $t = \sigma(s) = \frac{d-s}{d-c}a + \frac{s-c}{d-c}b$ ,  $s \in [c, d]$ . Therefore, if for some reason (and we shall presently see that there is such a reason) we do not distinguish between curves which are reparametrizations of each other, then the parametric interval of a curve is of no particular importance: we may consider a reparametrization of a given curve changing the given parametric interval to any other which we might prefer.

For every curve  $\gamma : [a, b] \rightarrow \mathbb{C}$  we consider the curve  $\neg\gamma : [a, b] \rightarrow \mathbb{C}$  given by  $(\neg\gamma)(t) = \gamma(a + b - t)$ ,  $t \in [a, b]$ . Then  $\neg\gamma$  is called *opposite* of  $\gamma$ . The curves  $\gamma$  and  $\neg\gamma$  have the same trajectory but opposite directions. Also, the two curves are simultaneously (piecewise) smooth and, in this case, their lengths are equal:

$$l(\neg\gamma) = \int_a^b |(\neg\gamma)'(t)| dt = \int_a^b |\gamma'(a + b - t)| dt = - \int_b^a |\gamma'(s)| ds = \int_a^b |\gamma'(s)| ds = l(\gamma).$$

If the curves  $\gamma_1 : [a, b] \rightarrow \mathbb{C}$  and  $\gamma_2 : [b, c] \rightarrow \mathbb{C}$  have  $\gamma_1(b) = \gamma_2(b)$ , then we say that  $\gamma_1, \gamma_2$  (in this order) are *successive* and then we may define the curve  $\gamma_1 \dot{+} \gamma_2 : [a, c] \rightarrow \mathbb{C}$  by

$$(\gamma_1 \dot{+} \gamma_2)(t) = \begin{cases} \gamma_1(t), & \text{if } a \leq t \leq b \\ \gamma_2(t), & \text{if } b \leq t \leq c \end{cases}$$

The curve  $\gamma_1 \dot{+} \gamma_2$  is called *sum* of  $\gamma_1$  and  $\gamma_2$ . If  $\gamma_1$  and  $\gamma_2$  are (piecewise) smooth,  $\gamma_1 \dot{+} \gamma_2$  is also piecewise smooth. The trajectory  $(\gamma_1 \dot{+} \gamma_2)^*$  is the union of the trajectories  $\gamma_1^*$  and  $\gamma_2^*$ .

Of course, the sum of two curves can be generalized to the sum of more than two curves provided that these are successive.



**Example 2.3.6.** Every polygonal line can be considered as the trajectory of a piecewise regular curve. This curve is the sum of successive curves each of which has as its trajectory a corresponding linear segment of the polygonal line.

Through the operation of summation of successive curves, we may consider successive curves as one curve and, conversely, we may consider one curve as a sum of successive curves.

The length of the sum of successive piecewise smooth curves equals the sum of their lengths:

$$\begin{aligned} l(\gamma_1 + \gamma_2) &= \int_a^c |(\gamma_1 + \gamma_2)'(t)| dt = \int_a^b |(\gamma_1 + \gamma_2)'(t)| dt + \int_b^c |(\gamma_1 + \gamma_2)'(t)| dt \\ &= \int_a^b |\gamma_1'(t)| dt + \int_b^c |\gamma_2'(t)| dt = l(\gamma_1) + l(\gamma_2). \end{aligned}$$

Now we shall extend the notion of integral of a complex function over an *interval* to the notion of integral of a complex function over a *curve*. Let  $\gamma : [a, b] \rightarrow \mathbb{C}$  be a piecewise smooth curve and let  $f : \gamma^* \rightarrow \mathbb{C}$  be continuous in the trajectory  $\gamma^* = \{\gamma(t) \mid t \in [a, b]\}$ . Then  $f \circ \gamma : [a, b] \rightarrow \mathbb{C}$  is continuous in  $[a, b]$ . Thus,  $(f \circ \gamma)\gamma'$  is piecewise continuous in  $[a, b]$  and hence integrable over  $[a, b]$ . We define the **curvilinear integral** of  $f$  over  $\gamma$  by

$$\int_{\gamma} f(z) dz = \int_a^b (f \circ \gamma)(t) \gamma'(t) dt = \int_a^b f(\gamma(t)) \gamma'(t) dt.$$

We shall usually write

$$\oint_{\gamma} f(z) dz$$

when  $\gamma$  is closed.

We remark that *whenever a curve  $\gamma$  is mentioned with respect either to its length  $l(\gamma)$  or to a curvilinear integral of a function over  $\gamma$  we shall always assume that  $\gamma$  is piecewise smooth.*

**Example 2.3.7.** Let  $\gamma$  be the curve with parametric equation  $z = \gamma(t) = (1-t)z_0 + tz_1$ ,  $t \in [0, 1]$ . The trajectory of  $\gamma$  is the linear segment  $[z_0, z_1]$  having direction from  $z_0$  to  $z_1$ . If  $f$  is continuous in  $[z_0, z_1]$ , then the curvilinear integral  $\int_{\gamma} f(z) dz$  is denoted  $\int_{[z_0, z_1]} f(z) dz$ . I.e.

$$\int_{[z_0, z_1]} f(z) dz = \int_{\gamma} f(z) dz = (z_1 - z_0) \int_0^1 f((1-t)z_0 + tz_1) dt.$$

This is the *curvilinear integral of  $f$  over the linear segment  $[z_0, z_1]$  from  $z_0$  to  $z_1$ .*

**Example 2.3.8.** Let  $r > 0$  and  $\gamma$  be the curve with parametric equation  $z = \gamma(t) = z_0 + r(\cos t + i \sin t)$ ,  $t \in [0, 2\pi]$ . The trajectory of  $\gamma$  is the circle  $C_{z_0}(r)$  with the positive direction of rotation around  $z_0$ . If  $f$  is continuous in the circle  $C_{z_0}(r)$ , then the curvilinear integral  $\oint_{\gamma} f(z) dz$  is denoted  $\oint_{C_{z_0}(r)} f(z) dz$ . I.e.,

$$\oint_{C_{z_0}(r)} f(z) dz = \oint_{\gamma} f(z) dz = \int_0^{2\pi} f(z_0 + r(\cos t + i \sin t)) r(-\sin t + i \cos t) dt.$$

This is the *curvilinear integral of  $f$  over the circle  $C_{z_0}(r)$  with the positive direction of rotation.*

An important concrete instance of the previous example is the following.

**Example 2.3.9.** If  $n \in \mathbb{Z}$ , we know that  $\int_0^{2\pi} \sin(nt) dt = 0$ . Also,  $\int_0^{2\pi} \cos(nt) dt = 2\pi$ , if  $n = 0$ , and  $\int_0^{2\pi} \cos(nt) dt = 0$ , if  $n \neq 0$ . Therefore, if  $n \in \mathbb{Z}$ , we get

$$\begin{aligned} \oint_{C_{z_0}(r)} (z - z_0)^n dz &= \int_0^{2\pi} r^n (\cos t + i \sin t)^n r(-\sin t + i \cos t) dt \\ &= ir^{n+1} \int_0^{2\pi} (\cos t + i \sin t)^n (\cos t + i \sin t) dt \\ &= ir^{n+1} \int_0^{2\pi} (\cos((n+1)t) + i \sin((n+1)t)) dt \\ &= \begin{cases} 2\pi i, & \text{if } n = -1 \\ 0, & \text{if } n \neq -1 \end{cases} \end{aligned}$$

The following propositions are easy to prove.

**Proposition 2.10.** *If  $\gamma$  is a piecewise smooth curve,  $f_1, f_2$  are continuous in  $\gamma^*$  and  $\lambda_1, \lambda_2 \in \mathbb{C}$ , then  $\int_{\gamma} (\lambda_1 f_1(z) + \lambda_2 f_2(z)) dz = \lambda_1 \int_{\gamma} f_1(z) dz + \lambda_2 \int_{\gamma} f_2(z) dz$ .*

*Proof.* An application of proposition 2.6 and of the definition of the curvilinear integral.  $\square$

**Proposition 2.11.** *If  $\gamma$  is a piecewise smooth curve and  $f$  is continuous in  $\gamma^*$ , then  $|\int_{\gamma} f(z) dz| \leq \sup_{z \in \gamma^*} |f(z)| l(\gamma)$ .*

*Proof.* If  $\gamma : [a, b] \rightarrow \mathbb{C}$ , then

$$\begin{aligned} |\int_{\gamma} f(z) dz| &= |\int_a^b f(\gamma(t)) \gamma'(t) dt| \leq \int_a^b |f(\gamma(t))| |\gamma'(t)| dt \leq \sup_{z \in \gamma^*} |f(z)| \int_a^b |\gamma'(t)| dt \\ &= \sup_{z \in \gamma^*} |f(z)| l(\gamma). \end{aligned}$$

The first inequality uses proposition 2.9.  $\square$

**Proposition 2.12.** *If  $\gamma$  is a piecewise smooth curve,  $f_n, \phi$  are continuous in  $\gamma^*$  and  $f_n \rightarrow f$  uniformly in  $\gamma^*$ , then  $\int_{\gamma} f_n(z) \phi(z) dz \rightarrow \int_{\gamma} f(z) \phi(z) dz$ .*

*Proof.* Because of uniform convergence,  $f$  is continuous in  $\gamma^*$ . Therefore, the existence of the integrals  $\int_{\gamma} f_n(z) \phi(z) dz$  and  $\int_{\gamma} f(z) \phi(z) dz$  is assured. Now, proposition 2.11 implies

$$\begin{aligned} |\int_{\gamma} f_n(z) \phi(z) dz - \int_{\gamma} f(z) \phi(z) dz| &= |\int_{\gamma} (f_n(z) - f(z)) \phi(z) dz| \\ &\leq \sup_{z \in \gamma^*} |f_n(z) - f(z)| \sup_{z \in \gamma^*} |\phi(z)| l(\gamma). \end{aligned}$$

Since  $\sup_{z \in \gamma^*} |f_n(z) - f(z)| \rightarrow 0$ , we get that  $\int_{\gamma} f_n(z) \phi(z) dz \rightarrow \int_{\gamma} f(z) \phi(z) dz$ .  $\square$

We may rewrite the result of proposition 2.12 in the form

$$\lim_{n \rightarrow +\infty} \int_{\gamma} f_n(z) \phi(z) dz = \int_{\gamma} \lim_{n \rightarrow +\infty} f_n(z) \phi(z) dz$$

of an interchange between the symbols  $\lim_{n \rightarrow +\infty}$  and  $\int_{\gamma}$ . This interchange under the assumption of uniform convergence is the content of proposition 2.12.

**Proposition 2.13.** *If  $\gamma$  is a piecewise smooth curve,  $f_n, \phi$  are continuous in  $\gamma^*$  and  $\sum_{n=1}^{+\infty} f_n = s$  uniformly in  $\gamma^*$ , then  $\sum_{n=1}^{+\infty} \int_{\gamma} f_n(z) \phi(z) dz = \int_{\gamma} s(z) \phi(z) dz$ .*

*Proof.* We consider the partial sums  $s_n = f_1 + \dots + f_n$  and we apply proposition 2.12 to them. Then

$$\sum_{k=1}^n \int_{\gamma} f_k(z) \phi(z) dz = \int_{\gamma} \sum_{k=1}^n f_k(z) \phi(z) dz = \int_{\gamma} s_n(z) \phi(z) dz \rightarrow \int_{\gamma} s(z) \phi(z) dz.$$

I.e. the series (of numbers)  $\sum_{n=1}^{+\infty} \int_{\gamma} f_n(z) \phi(z) dz$  converges to (the number)  $\int_{\gamma} s(z) \phi(z) dz$ .  $\square$

As in the case of proposition 2.12, we may rewrite the result of proposition 2.13 in the form

$$\sum_{n=1}^{+\infty} \int_{\gamma} f_n(z) \phi(z) dz = \int_{\gamma} \sum_{n=1}^{+\infty} f_n(z) \phi(z) dz,$$

since  $\sum_{n=1}^{+\infty} f_n(z) = s(z)$  for every  $z \in \gamma^*$ . Again, this interchange between the symbols  $\sum_{n=1}^{+\infty}$  and  $\int_{\gamma}$  under the assumption of uniform convergence is the content of proposition 2.13.

**Proposition 2.14.** *If each of the piecewise smooth curves  $\gamma_1, \gamma_2$  is a reparametrization of the other and  $f$  is continuous in  $\gamma_1^* = \gamma_2^*$ , then  $\int_{\gamma_2} f(z) dz = \int_{\gamma_1} f(z) dz$ .*

*Proof.* If  $\gamma_1 : [a, b] \rightarrow \mathbb{C}$  and  $\gamma_2 : [c, d] \rightarrow \mathbb{C}$ , then there is a change of parameter  $\sigma : [c, d] \rightarrow [a, b]$  so that  $\gamma_2(s) = \gamma_1(\sigma(s))$  for all  $s \in [c, d]$ . Then

$$\begin{aligned} \int_{\gamma_2} f(z) dz &= \int_c^d f(\gamma_2(s))\gamma_2'(s) ds = \int_c^d f(\gamma_1(\sigma(s)))\gamma_1'(\sigma(s))\sigma'(s) ds \\ &= \int_a^b f(\gamma_1(t))\gamma_1'(t) dt = \int_{\gamma_1} f(z) dz \end{aligned}$$

after a change of parameter in the third integral.  $\square$

At this point we observe that replacing a curve  $\gamma_1$  with a reparametrization  $\gamma_2$  of it does not alter certain objects related to the curve: its trajectory, its endpoints, its direction, its length, the number of times it covers its trajectory and, more important, the curvilinear integrals of continuous functions defined over its trajectory. Since in this course we shall use curves mostly to evaluate curvilinear integrals, we conclude that there is no reason to distinguish between a curve and its reparametrizations. Therefore, when we have a geometric object  $C$  which we would call, in everyday language, *curve in the plane*, e.g. a linear segment or a circle or a polygonal line, and a continuous function  $f : C \rightarrow \mathbb{C}$ , we can give a meaning to

$$\int_C f(z) dz$$

by specifying a piecewise continuously differentiable  $\gamma : [a, b] \rightarrow \mathbb{C}$ , i.e. a piecewise smooth curve, with trajectory  $\gamma^*$  coinciding with  $C$ , with endpoints coinciding with the endpoints of  $C$  and a specific assigned direction. The use of different curves, which are reparametrizations of the particular  $\gamma$  we have chosen, will not alter the value of the integral. In fact we have already seen two examples of this situation. One is the curvilinear integral  $\int_{[z_0, z_1]} f(z) dz$  for which we use any parametric equation with trajectory equal to the linear segment  $[z_0, z_1]$  and direction from  $z_0$  to  $z_1$ . The simplest such parametric equation is  $z = \gamma(t) = (1 - t)z_0 + tz_1$ ,  $t \in [0, 1]$ . The second example is the curvilinear integral  $\oint_{C_{z_0}(r)} f(z) dz$  for which we use any parametric equation with trajectory equal to the circle  $C_{z_0}(r)$  and which covers this circle once and in the positive direction of rotation around  $z_0$ . The simplest such parametric equation is  $z = \gamma(t) = z_0 + r(\cos t + i \sin t)$ ,  $t \in [0, 2\pi]$ .

**Proposition 2.15.** *Let  $\gamma_1, \gamma_2$  be two successive piecewise smooth curves and let  $f$  be continuous in  $\gamma_1^* \cup \gamma_2^*$ . Then  $\int_{\gamma_1 + \gamma_2} f(z) dz = \int_{\gamma_1} f(z) dz + \int_{\gamma_2} f(z) dz$ .*

*Proof.* Let  $\gamma_1 : [a, b] \rightarrow \mathbb{C}$  and  $\gamma_2 : [b, c] \rightarrow \mathbb{C}$  with  $\gamma_1(b) = \gamma_2(b)$ . Then

$$\begin{aligned} \int_{\gamma_1 + \gamma_2} f(z) dz &= \int_a^c f((\gamma_1 + \gamma_2)(t))(\gamma_1 + \gamma_2)'(t) dt \\ &= \int_a^b f(\gamma_1(t))\gamma_1'(t) dt + \int_b^c f(\gamma_2(t))\gamma_2'(t) dt = \int_{\gamma_1} f(z) dz + \int_{\gamma_2} f(z) dz. \end{aligned}$$

The second equality uses proposition 2.7.  $\square$

**Proposition 2.16.** *If  $\gamma$  is a piecewise smooth curve and  $f$  is continuous in  $\gamma^*$ , then  $\int_{-\gamma} f(z) dz = -\int_{\gamma} f(z) dz$ .*

*Proof.* If  $\gamma : [a, b] \rightarrow \mathbb{C}$ , then

$$\begin{aligned} \int_{-\gamma} f(z) dz &= \int_a^b f((-\gamma)(t))(-\gamma)'(t) dt = -\int_a^b f(\gamma(a + b - t))\gamma'(a + b - t) dt \\ &= \int_b^a f(\gamma(s))\gamma'(s) ds = -\int_a^b f(\gamma(s))\gamma'(s) ds = -\int_{\gamma} f(z) dz. \end{aligned}$$

after a simple change of parameter in the third integral.  $\square$

**Example 2.3.10.** Let  $\gamma$  be the curve describing the linear segment  $[z_0, z_1]$  from  $z_0$  to  $z_1$ . Then  $-\gamma$  describes the same segment from  $z_1$  to  $z_0$ . Therefore,  $\int_{[z_0, z_1]} f(z) dz = \int_{\gamma} f(z) dz$  and  $\int_{[z_1, z_0]} f(z) dz = \int_{-\gamma} f(z) dz$ . Hence  $\int_{[z_1, z_0]} f(z) dz = -\int_{[z_0, z_1]} f(z) dz$ .

Before we leave this section, we should mention three variants of the notion of the curvilinear integral. Let  $\gamma : [a, b] \rightarrow \mathbb{C}$  be a piecewise smooth curve and let  $f : \gamma^* \rightarrow \mathbb{C}$  be continuous in the trajectory  $\gamma^* = \{\gamma(t) \mid t \in [a, b]\}$ . If  $\gamma(t) = (x(t), y(t)) = x(t) + iy(t)$  for every  $t \in [a, b]$ , we define

$$\int_{\gamma} f(z) dx = \int_a^b f(\gamma(t))x'(t) dt, \quad \int_{\gamma} f(z) dy = \int_a^b f(\gamma(t))y'(t) dt,$$

$$\int_{\gamma} f(z) |dz| = \int_a^b f(\gamma(t))|\gamma'(t)| dt.$$

Trivially, we have  $\int_{\gamma} f(z) dz = \int_{\gamma} f(z) dx + i \int_{\gamma} f(z) dy$ .

We leave to the reader the easy task to show that each of the three new integrals satisfies all properties of the original  $\int_{\gamma} f(z) dz$ , expressed in propositions 2.10 - 2.16. The only difference is with the integral  $\int_{\gamma} f(z) |dz|$  which, regarding proposition 2.16, does *not* change its sign when we replace  $\gamma$  with  $\neg\gamma$ . Moreover, the basic inequality in proposition 2.11 takes the more precise form:

$$\left| \int_{\gamma} f(z) dz \right| \leq \int_{\gamma} |f(z)| |dz| \leq \sup_{z \in \gamma^*} |f(z)| l(\gamma).$$

Indeed, observing the string of equalities/inequalities in the proof of proposition 2.11, we recognize  $\int_{\gamma} |f(z)| |dz|$  as the third integral from the left. It is very common with beginning students to make the mistake:  $\left| \int_{\gamma} f(z) dz \right| \leq \int_{\gamma} |f(z)| dz$ .

We should also say that  $\int_{\gamma} |dz| = l(\gamma)$ . In calculus texts one usually sees the symbol  $ds$  instead of  $|dz|$  for the infinitesimal length  $|\gamma'(t)| dt$  over the curve.

### Exercises.

**2.3.1.** Calculate  $\int_{\gamma} |z| dz$ , where  $\gamma$  is each of the following curves with initial endpoint  $-i$  and final endpoint  $i$ . (i)  $\gamma(t) = it$  for  $t \in [-1, 1]$ . (ii)  $\gamma(t) = \cos t + i \sin t$  for  $t \in [-\frac{\pi}{2}, \frac{\pi}{2}]$ . (iii)  $\gamma(t) = -\cos t + i \sin t$  for  $t \in [-\frac{\pi}{2}, \frac{\pi}{2}]$ .

**2.3.2.** (i) If  $n \in \mathbb{Z}$ ,  $n \geq 0$ , prove that  $\int_{\gamma} z^n dz = \frac{z_1^{n+1} - z_0^{n+1}}{n+1}$ , where  $z_0, z_1$  are the initial and the final endpoint of the piecewise smooth  $\gamma$ .

(ii) Are there polynomials  $p_n(z)$  so that  $p_n(z) \rightarrow \frac{1}{z}$  uniformly in the circle  $C_0(1)$ ? Think in terms of curvilinear integrals over the circle  $C_0(1)$ .

**2.3.3.** Let  $f$  be continuous in the ring  $\{z \mid 0 < |z| < r_0\}$  or in the ring  $\{z \mid r_0 < |z| < +\infty\}$ . We define  $M(r) = \max_{z \in C_0(r)} |f(z)|$  and assume that  $rM(r) \rightarrow 0$  when  $r \rightarrow 0$  or  $r \rightarrow +\infty$ , respectively. If  $\gamma_r(t) = r(\cos t + i \sin t)$  for  $t_1 \leq t \leq t_2$ , then prove that  $\int_{\gamma_r} f(z) dz \rightarrow 0$  when  $r \rightarrow 0$  or  $r \rightarrow +\infty$ , respectively.

**2.3.4.** Let  $f : D_{z_0}(R) \rightarrow \mathbb{C}$  be continuous. Prove that  $\lim_{r \rightarrow 0} \oint_{C_{z_0}(r)} \frac{f(z)}{z-z_0} dz = 2\pi i f(z_0)$ .

**2.3.5.** Let  $f : \Omega \rightarrow \mathbb{C}$  be continuous in the open set  $\Omega$  and let  $[a_n, b_n], [a, b] \subseteq \Omega$  for every  $n$ . If  $a_n \rightarrow a$  and  $b_n \rightarrow b$ , prove that  $\int_{[a_n, b_n]} f(z) dz \rightarrow \int_{[a, b]} f(z) dz$ .

**2.3.6.** Let  $f : \Omega \rightarrow \mathbb{C}$  be continuous in the open set  $\Omega$  and  $\gamma$  be a piecewise smooth curve in  $\Omega$ . Prove that for every  $\epsilon > 0$  there is a polygonal curve  $\sigma$  in  $\Omega$  so that  $|\int_{\sigma} f(z) dz - \int_{\gamma} f(z) dz| < \epsilon$ .

**2.3.7.** Prove that  $|\int_a^b f(t) dt| = \int_a^b |f(t)| dt$  if and only if there is some halfline  $l$  with vertex 0 so that  $f(t) \in l$  for every continuity point  $t$  of  $f$ .

**2.3.8.** Let  $\gamma : [a, b] \rightarrow \mathbb{C}$  be a piecewise smooth curve and  $f : \gamma^* \rightarrow \mathbb{C}$  be continuous in  $\gamma^*$ . Consider any subdivision  $\Delta = \{t_0, \dots, t_n\}$  of  $[a, b]$  and any choice  $\Xi = \{\xi_1, \dots, \xi_n\}$  of intermediate points  $\xi_k \in [t_{k-1}, t_k]$ . Then we say that  $\Delta^* = \{z_0, \dots, z_n\}$ , where  $z_k = \gamma(t_k)$ , is a subdivision of the trajectory  $\gamma^*$  and that  $\Xi^* = \{\eta_1, \dots, \eta_n\}$ , where  $\eta_k = \gamma(\xi_k)$ , is a choice of intermediate points on the trajectory:  $\eta_k$  is between  $z_{k-1}$  and  $z_k$  on the trajectory. We say that  $\sum_{k=1}^n f(z_k)(\eta_k - \eta_{k-1})$  is the corresponding Riemann sum. If  $w(\Delta^*) = \max_{1 \leq k \leq n} |z_k - z_{k-1}|$  is the width of the subdivision  $\Delta^*$ , then prove that  $\lim_{w(\Delta^*) \rightarrow 0} \sum_{k=1}^n f(z_k)(\eta_k - \eta_{k-1}) = \int_{\gamma} f(z) dz$ .

## Chapter 3

# Holomorphic functions.

### 3.1 Differentiability and holomorphy.

Let  $f$  be a complex function defined in  $A \subseteq \mathbb{C}$  and  $z_0$  be an interior point of  $A$ . We say that  $f$  is **differentiable** at  $z_0$  if  $\lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0}$  exists and is a complex number. We call this limit **derivative** of  $f$  at  $z_0$  and denote it

$$f'(z_0) = \frac{df}{dz}(z_0) = \lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0}.$$

**Example 3.1.1.** The constant function  $c$  is differentiable at every point of  $\mathbb{C}$  and its derivative is the constant function 0. Indeed, for every  $z_0$  we have  $\frac{dc}{dz}(z_0) = \lim_{z \rightarrow z_0} \frac{c - c}{z - z_0} = \lim_{z \rightarrow z_0} 0 = 0$ .

**Example 3.1.2.** The function  $z$  is differentiable at every point of  $\mathbb{C}$  and its derivative is the constant function 1: for every  $z_0$  we have  $\frac{dz}{dz}(z_0) = \lim_{z \rightarrow z_0} \frac{z - z_0}{z - z_0} = \lim_{z \rightarrow z_0} 1 = 1$ .

**Example 3.1.3.** Let  $f(z) = \bar{z}$ . We shall prove that the  $\lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0} = \lim_{z \rightarrow z_0} \frac{\bar{z} - \bar{z}_0}{z - z_0}$  does not exist, i.e.  $f$  is not differentiable at any  $z_0$ .

Let  $z_0 = x_0 + iy_0$ . The limit of  $\frac{f(z) - f(z_0)}{z - z_0}$  when  $z \rightarrow z_0$  on the horizontal line containing  $z_0$  is

$$\lim_{x \rightarrow x_0} \frac{\overline{(x + iy_0)} - \overline{(x_0 + iy_0)}}{(x + iy_0) - (x_0 + iy_0)} = \lim_{x \rightarrow x_0} \frac{x - x_0}{x - x_0} = \lim_{x \rightarrow x_0} 1 = 1$$

and the limit of  $\frac{f(z) - f(z_0)}{z - z_0}$  when  $z \rightarrow z_0$  on the vertical line containing  $z_0$  is

$$\lim_{y \rightarrow y_0} \frac{\overline{(x_0 + iy)} - \overline{(x_0 + iy_0)}}{(x_0 + iy) - (x_0 + iy_0)} = \lim_{y \rightarrow y_0} \frac{-iy + iy_0}{iy - iy_0} = \lim_{y \rightarrow y_0} (-1) = -1.$$

Since these two limits are different, the  $\lim_{z \rightarrow z_0} \frac{\bar{z} - \bar{z}_0}{z - z_0}$  does not exist.

The proofs of the following four propositions are identical with the proofs of the well-known analogous propositions for real functions of a real variable and we omit them.

**Proposition 3.1.** *If  $f : A \rightarrow \mathbb{C}$  is differentiable at the interior point  $z_0$  of  $A \subseteq \mathbb{C}$ , then  $f$  is continuous at  $z_0$ .*

**Proposition 3.2.** *If  $f, g : A \rightarrow \mathbb{C}$  are differentiable at the interior point  $z_0$  of  $A \subseteq \mathbb{C}$ , then  $f + g, f - g, fg : A \rightarrow \mathbb{C}$  are also differentiable at  $z_0$ . Furthermore, if  $g(z) \neq 0$  for all  $z \in A$ , then  $\frac{f}{g} : A \rightarrow \mathbb{C}$  is differentiable at  $z_0$ . Finally,  $(f + g)'(z_0) = f'(z_0) + g'(z_0)$ ,  $(f - g)'(z_0) = f'(z_0) - g'(z_0)$ ,  $(fg)'(z_0) = f'(z_0)g(z_0) + f(z_0)g'(z_0)$  and  $(\frac{f}{g})'(z_0) = \frac{f'(z_0)g(z_0) - f(z_0)g'(z_0)}{(g(z_0))^2}$ .*

**Proposition 3.3.** *If  $f : A \rightarrow B$  is differentiable at the interior point  $z_0$  of  $A \subseteq \mathbb{C}$  and  $g : B \rightarrow \mathbb{C}$  is differentiable at the interior point  $w_0 = f(z_0)$  of  $B \subseteq \mathbb{C}$ , then  $g \circ f : A \rightarrow \mathbb{C}$  is differentiable at  $z_0$ . Also,  $(g \circ f)'(z_0) = g'(w_0)f'(z_0)$ .*

**Proposition 3.4.** Let  $f : A \rightarrow B$  be one-to-one from  $A \subseteq \mathbb{C}$  onto  $B \subseteq \mathbb{C}$  and let  $f^{-1} : B \rightarrow A$  be the inverse function. Let also  $z_0$  be an interior point of  $A$  and  $w_0 = f(z_0)$  be an interior point of  $B$ . If  $f$  is differentiable at  $z_0$  and  $f'(z_0) \neq 0$  and  $f^{-1}$  is continuous at  $w_0$ , then  $f^{-1}$  is differentiable at  $w_0$  and  $(f^{-1})'(w_0) = \frac{1}{f'(z_0)}$ .

**Example 3.1.4.** Starting with the derivatives of the constant function  $c$  and the function  $z$  and using the usual algebraic rules for derivatives, we get that every polynomial function is differentiable at every point of  $\mathbb{C}$  and that its derivative is another polynomial function: if  $p(z) = a_0 + a_1z + a_2z^2 + \cdots + a_nz^n$ , then  $p'(z) = a_1 + 2a_2z + \cdots + na_nz^{n-1}$ .

**Example 3.1.5.** Every rational function is differentiable at every point of its domain of definition and its derivative is another rational function.

**Example 3.1.6.** If  $h(z) = (z^2 - 3z + 2)^{15} - 3(z^2 - 3z + 2)^2$ , then by the chain rule we get  $h'(z) = 15(z^2 - 3z + 2)^{14}(2z - 3) - 6(z^2 - 3z + 2)(2z - 3)$ .

Let  $f$  be a complex function defined in  $A \subseteq \mathbb{C}$  and  $z_0$  be an interior point of  $A$ . We say that  $f$  is **holomorphic** (or **analytic**) at  $z_0$  if there is  $r > 0$  so that  $D_{z_0}(r) \subseteq A$  and  $f$  is differentiable at every point of  $D_{z_0}(r)$ .

The notion of holomorphy is stronger than the notion of differentiability: for a function to be holomorphic at a point it is necessary for it to be differentiable at this point *and* at all nearby points.

**Example 3.1.7.** Every polynomial function is holomorphic at every point of  $\mathbb{C}$ .

**Example 3.1.8.** Every rational function is holomorphic at every point of its domain of definition.

**Example 3.1.9.** Let  $f(z) = |z|^2$ . We have  $\lim_{z \rightarrow 0} \frac{f(z) - f(0)}{z - 0} = \lim_{z \rightarrow 0} \bar{z} = 0$  and  $f$  is differentiable at 0 with  $f'(0) = 0$ .

We take an arbitrary  $z_0 \neq 0$  and we shall prove that the  $\lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0} = \lim_{z \rightarrow z_0} \frac{|z|^2 - |z_0|^2}{z - z_0}$  does not exist and therefore  $f$  is not differentiable at  $z_0$ .

Let  $z_0 = x_0 + iy_0$ . The limit of  $\frac{f(z) - f(z_0)}{z - z_0}$  when  $z \rightarrow z_0$  on the horizontal line containing  $z_0$  is

$$\lim_{x \rightarrow x_0} \frac{|x + iy_0|^2 - |x_0 + iy_0|^2}{(x + iy_0) - (x_0 + iy_0)} = \lim_{x \rightarrow x_0} \frac{x^2 - x_0^2}{x - x_0} = \lim_{x \rightarrow x_0} (x + x_0) = 2x_0$$

and the limit of  $\frac{f(z) - f(z_0)}{z - z_0}$  when  $z \rightarrow z_0$  on the vertical line containing  $z_0$  is

$$\lim_{y \rightarrow y_0} \frac{|x_0 + iy|^2 - |x_0 + iy_0|^2}{(x_0 + iy) - (x_0 + iy_0)} = \lim_{y \rightarrow y_0} \frac{y^2 - y_0^2}{iy - iy_0} = -i \lim_{y \rightarrow y_0} (y + y_0) = -2iy_0.$$

Since  $z_0 \neq 0$ , these two limits are different and the  $\lim_{z \rightarrow z_0} \frac{|z|^2 - |z_0|^2}{z - z_0}$  does not exist.

We conclude that  $f$  is differentiable only at 0 and that it is nowhere holomorphic.

The set of points at which  $f$  is holomorphic is called **domain of holomorphy** of  $f$ .

**Proposition 3.5.** If  $B \subseteq \mathbb{C}$  is the set of the points at which the complex function  $f$  is differentiable, then the domain of holomorphy of  $f$  is the interior of  $B$ . In particular, the domain of holomorphy of  $f$  is an open set.

*Proof.* Let  $U$  be the domain of holomorphy of  $f$ . If  $z \in U$ , there is  $r > 0$  so that  $f$  is differentiable at every point of  $D_z(r)$  and hence  $D_z(r) \subseteq B$ . Thus  $z$  is an interior point of  $B$ , i.e.  $z \in B^\circ$ .

Conversely, let  $z \in B^\circ$ . Then there is  $r > 0$  so that  $D_z(r) \subseteq B$ , and so  $f$  is differentiable at every point of  $D_z(r)$ . Therefore  $f$  is holomorphic at  $z$ , i.e.  $z \in U$ .  $\square$

**Example 3.1.10.** The domain of holomorphy of any polynomial function is  $\mathbb{C}$ .

**Example 3.1.11.** The domain of holomorphy of any rational function is its domain of definition.

**Example 3.1.12.** The domain of holomorphy of both functions  $f(z) = \bar{z}$  and  $f(z) = |z|^2$  is the empty set.

Let  $\Omega \subseteq \mathbb{C}$  be an open set. We say that the complex function  $f$  is holomorphic (or analytic) in  $\Omega$  if it is holomorphic at every point of  $\Omega$  or, equivalently, if  $\Omega$  is a subset of the domain of holomorphy of  $f$ .

Clearly, the largest open set  $\Omega$  in which  $f$  is holomorphic is its domain of holomorphy. It is also clear that *if  $f$  is differentiable at every point of an open set  $\Omega$ , then  $f$  is holomorphic in  $\Omega$ .*

Let the complex function  $f$  be defined in the neighborhood  $D_\infty(r)$  of  $\infty$ . We consider the complex function  $g$  defined in  $D_0(r)$  by

$$g(w) = f(1/w), \quad \text{for every } w \text{ with } |w| < r.$$

We say that  $f$  is differentiable or holomorphic at  $\infty$  if  $g$  is differentiable or holomorphic, respectively, at 0.

We observe that  $g(0) = f(\infty)$  and that the inverse functions  $w = \frac{1}{z}$  and  $z = \frac{1}{w}$  map each of the neighborhoods  $D_\infty(r) = \{z \mid |z| > \frac{1}{r}\} \cup \{\infty\}$  and  $D_0(r) = \{w \mid |w| < r\}$  onto the other. Now we shall see that the condition of differentiability of  $f$  at  $\infty$ , i.e. the differentiability of  $g$  at 0, can be translated into an equivalent condition in terms of  $f$  itself.

**Proposition 3.6.** *Let  $f$  be a complex function defined in  $D_\infty(r)$ . Then  $f$  is differentiable at  $\infty$  if and only if  $\lim_{z \rightarrow \infty} z(f(z) - f(\infty)) \in \mathbb{C}$ .*

*Proof.* Let  $g(w) = f(\frac{1}{z})$  be the function considered in the above definition. Through the change of variable  $w = \frac{1}{z}$ , we have  $\frac{g(w) - g(0)}{w - 0} = z(f(z) - f(\infty))$ . Thus, the existence of  $\lim_{w \rightarrow 0} \frac{g(w) - g(0)}{w - 0}$  is equivalent to the existence of  $\lim_{z \rightarrow \infty} z(f(z) - f(\infty))$ . In fact the two limits are equal.  $\square$

It is easy to see that differentiability of  $f$  at  $\infty$  implies continuity of  $f$  at  $\infty$ .

**Example 3.1.13.** We shall check the differentiability (and hence holomorphy) of polynomial and rational functions. We recall the notation and the results of examples 1.3.1 and 1.3.2.

A polynomial function  $p$  is continuous and complex-valued at  $\infty$  only if it is a constant  $p(z) = a_0$  and provided we define  $p(\infty) = a_0$ . In this case it is also differentiable at  $\infty$ , since

$$\lim_{z \rightarrow \infty} z(p(z) - p(\infty)) = \lim_{z \rightarrow \infty} 0 = 0.$$

A rational function  $r(z) = \frac{a_n z^n + \dots + a_1 z + a_0}{b_m z^m + \dots + b_1 z + b_0}$  is continuous and complex-valued at  $\infty$  only if  $n \leq m$ , where  $n$  and  $m$  are the degrees of its numerator and denominator. If  $n = m$ , then we define  $r(\infty) = \frac{a_n}{b_n}$  and then, after some algebraic manipulations, we get

$$\lim_{z \rightarrow \infty} z(r(z) - r(\infty)) = \lim_{z \rightarrow \infty} z \left( \frac{a_n z^n + \dots + a_1 z + a_0}{b_n z^n + \dots + b_1 z + b_0} - \frac{a_n}{b_n} \right) = \frac{a_{n-1} b_n - a_n b_{n-1}}{b_n^2}.$$

If  $n < m$ , then we define  $r(\infty) = 0$  and then we get

$$\lim_{z \rightarrow \infty} z(r(z) - r(\infty)) = \lim_{z \rightarrow \infty} z \frac{a_n z^n + \dots + a_1 z + a_0}{b_m z^m + \dots + b_1 z + b_0} = \begin{cases} \frac{a_n}{b_{n+1}}, & \text{if } n + 1 = m \\ 0, & \text{if } n + 1 < m \end{cases}$$

### Exercises.

**3.1.1.** Check the differentiability of the functions  $\operatorname{Re} z$ ,  $\operatorname{Im} z$  and  $|z|$ .

**3.1.2.** Let  $\Omega$  be open and  $f : \Omega \rightarrow \mathbb{C}$ . We take  $\Omega^* = \{z \mid \bar{z} \in \Omega\}$  and  $f^* : \Omega^* \rightarrow \mathbb{C}$  given by  $f^*(z) = \overline{f(\bar{z})}$  for every  $z \in \Omega^*$ . Prove that  $\Omega^*$  is open and that, if  $f$  is differentiable at  $z_0 \in \Omega$ , then  $f^*$  is differentiable at  $\bar{z}_0 \in \Omega^*$ .

**3.1.3.** Consider open sets  $U, V$  and  $f : V \rightarrow U, g : U \rightarrow \mathbb{C}, h : V \rightarrow \mathbb{C}$  so that  $h$  is one-to-one and  $h = g \circ f$ . If  $h$  is differentiable at  $w_0 \in V$ ,  $g$  is differentiable at  $z_0 = f(w_0)$ ,  $g'(z_0) \neq 0$  and  $f$  is continuous at  $w_0$ , prove that  $f$  is differentiable at  $w_0$  and  $f'(w_0) = \frac{h'(w_0)}{g'(z_0)}$ .

**3.1.4.** (i) If  $p$  is a polynomial of degree  $n$  with roots  $z_1, \dots, z_n$ , prove  $\frac{p'(z)}{p(z)} = \frac{1}{z-z_1} + \dots + \frac{1}{z-z_n}$  for every  $z \neq z_1, \dots, z_n$ . Then prove that, if the roots of  $p$  are contained in a closed halfplane, then the roots of  $p'$  are contained in the same halfplane. Conclude that the roots of  $p'$  are contained in the smallest convex polygon which contains the roots of  $p$ .

(ii) For every  $a$  and every  $n \in \mathbb{N}, n \geq 2$  prove that the equation  $1 + z + az^n = 0$  has at least one root  $z \in \overline{D}_0(2)$ .

**3.1.5.** (i) Let  $a_1, \dots, a_n$  be distinct and  $q(z) = (z - a_1) \cdots (z - a_n)$ . If the polynomial  $p$  has degree  $< n$ , prove  $\frac{p(z)}{q(z)} = \sum_{k=1}^n \frac{p(a_k)}{q'(a_k)(z-a_k)}$  for every  $z \neq a_1, \dots, a_n$ .

(ii) Let  $a_1, \dots, a_n$  be distinct. Prove that for every  $c_1, \dots, c_n$  there is a unique polynomial  $p$  of degree  $< n$  so that  $p(a_k) = c_k$  for every  $k = 1, \dots, n$ .

**3.1.6.** Let  $f$  have continuous derivative in a neighborhood of  $z_0$ . Prove that  $\frac{f(z'_n) - f(z''_n)}{z'_n - z''_n} \rightarrow f'(z_0)$  if  $z'_n \rightarrow z_0, z''_n \rightarrow z_0$  and  $z'_n \neq z''_n$  for every  $n$ .

## 3.2 The Cauchy-Riemann equations.

Now we shall relate the differentiability of  $f$ , as a complex function of  $z = x + iy$ , at some interior point  $z_0 = x_0 + iy_0$  of its domain  $A \subseteq \mathbb{C}$  with the partial derivatives of  $u = \operatorname{Re} f$  and  $v = \operatorname{Im} f$ , as functions of  $(x, y)$  at the same point  $(x_0, y_0)$ .

**Theorem 3.1.** *Let  $f$  be a complex function defined in  $A \subseteq \mathbb{C}, z_0 = (x_0, y_0)$  be an interior point of  $A$  and let  $u, v$  be the real and imaginary part of  $f$ . If  $f$  is differentiable at  $z_0$ , then  $u, v$  have partial derivatives with respect to  $x$  and  $y$  at  $(x_0, y_0)$  and*

$$\frac{\partial u}{\partial x}(x_0, y_0) = \frac{\partial v}{\partial y}(x_0, y_0), \quad \frac{\partial u}{\partial y}(x_0, y_0) = -\frac{\partial v}{\partial x}(x_0, y_0). \quad (3.1)$$

*Proof.* We assume

$$\lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0} = f'(z_0) = \mu + i\nu, \quad \mu, \nu \in \mathbb{R}. \quad (3.2)$$

Since the limit of  $\frac{f(z) - f(z_0)}{z - z_0}$  exists when  $z$  tends to  $z_0$ , the limits of the same expression when  $z$  tends to  $z_0$  on the horizontal line containing  $z_0$  as well as on the vertical line containing  $z_0$  also exist and have the same value:

$$\lim_{x \rightarrow x_0} \frac{f(x, y_0) - f(x_0, y_0)}{x - x_0} = \mu + i\nu, \quad \lim_{y \rightarrow y_0} \frac{f(x_0, y) - f(x_0, y_0)}{iy - iy_0} = \mu + i\nu. \quad (3.3)$$

From the first limit in (3.3) we get  $\lim_{x \rightarrow x_0} \frac{u(x, y_0) + iv(x, y_0) - u(x_0, y_0) - iv(x_0, y_0)}{x - x_0} = \mu + i\nu$ , and hence

$$\begin{aligned} \frac{\partial u}{\partial x}(x_0, y_0) &= \lim_{x \rightarrow x_0} \frac{u(x, y_0) - u(x_0, y_0)}{x - x_0} = \mu, \\ \frac{\partial v}{\partial x}(x_0, y_0) &= \lim_{x \rightarrow x_0} \frac{v(x, y_0) - v(x_0, y_0)}{x - x_0} = \nu. \end{aligned} \quad (3.4)$$

From the second limit in (3.3) we find  $\lim_{y \rightarrow y_0} \frac{u(x_0, y) + iv(x_0, y) - u(x_0, y_0) - iv(x_0, y_0)}{iy - iy_0} = \mu + i\nu$ , and hence

$$\begin{aligned} \frac{\partial v}{\partial y}(x_0, y_0) &= \lim_{y \rightarrow y_0} \frac{v(x_0, y) - v(x_0, y_0)}{y - y_0} = \mu, \\ \frac{\partial u}{\partial y}(x_0, y_0) &= \lim_{y \rightarrow y_0} \frac{u(x_0, y) - u(x_0, y_0)}{y - y_0} = -\nu. \end{aligned} \quad (3.5)$$

Comparing (3.4) and (3.5) we get (3.1). □



The equalities (3.1) are called **(system of) Cauchy-Riemann equations** at the point  $(x_0, y_0)$ . We observe that, if  $f$  is differentiable at  $z_0$ , then (3.2), (3.4) and (3.5) imply

$$f'(z_0) = \frac{\partial u}{\partial x}(x_0, y_0) + i \frac{\partial v}{\partial x}(x_0, y_0) = \frac{\partial v}{\partial y}(x_0, y_0) - i \frac{\partial u}{\partial y}(x_0, y_0).$$

The next result is the converse of theorem 3.1 but with extra assumptions.

**Theorem 3.2.** *Let  $f$  be a complex function defined in  $A \subseteq \mathbb{C}$ ,  $z_0 = (x_0, y_0)$  be an interior point of  $A$  and let  $u, v$  be the real and imaginary part of  $f$ . If  $u, v$  have partial derivatives with respect to  $x$  and  $y$  at every point of some neighborhood of  $(x_0, y_0)$  and if these partial derivatives are continuous at  $(x_0, y_0)$  and if they satisfy the system of C-R equations at  $(x_0, y_0)$ , then  $f$  is differentiable at  $z_0$ .*

*Proof.* Using the C-R equations, we define the real numbers  $\mu$  and  $\nu$  by:

$$\mu = \frac{\partial u}{\partial x}(x_0, y_0) = \frac{\partial v}{\partial y}(x_0, y_0), \quad \nu = -\frac{\partial u}{\partial y}(x_0, y_0) = \frac{\partial v}{\partial x}(x_0, y_0). \quad (3.6)$$

Now take an arbitrary  $\epsilon > 0$ . Since  $\frac{\partial u}{\partial x}, \frac{\partial u}{\partial y}$  are continuous at  $(x_0, y_0)$ , there is  $r > 0$  so that

$$\left| \frac{\partial u}{\partial x}(x, y) - \mu \right| < \frac{\epsilon}{4}, \quad \left| \frac{\partial u}{\partial y}(x, y) + \nu \right| < \frac{\epsilon}{4} \quad \text{for every } (x, y) \in D_{(x_0, y_0)}(r). \quad (3.7)$$

We take any  $(x, y) \in D_{(x_0, y_0)}(r)$  and we write

$$u(x, y) - u(x_0, y_0) = u(x, y) - u(x_0, y) + u(x_0, y) - u(x_0, y_0). \quad (3.8)$$

By the mean value theorem, there is  $x'$  between  $x$  and  $x_0$  so that

$$u(x, y) - u(x_0, y) = \frac{\partial u}{\partial x}(x', y)(x - x_0) \quad (3.9)$$

and  $y'$  between  $y$  and  $y_0$  so that

$$u(x_0, y) - u(x_0, y_0) = \frac{\partial u}{\partial y}(x_0, y')(y - y_0). \quad (3.10)$$

The  $x', y'$  depend on  $x, y$ , but the points  $(x', y), (x_0, y')$  belong to  $D_{(x_0, y_0)}(r)$ . Therefore, (3.7) implies

$$\left| \frac{\partial u}{\partial x}(x', y) - \mu \right| < \frac{\epsilon}{4}, \quad \left| \frac{\partial u}{\partial y}(x_0, y') + \nu \right| < \frac{\epsilon}{4}. \quad (3.11)$$

Combining (3.8), (3.9) and (3.10), we find

$$\begin{aligned} u(x, y) - u(x_0, y_0) - (\mu(x - x_0) - \nu(y - y_0)) \\ &= (u(x, y) - u(x_0, y) - \mu(x - x_0)) + (u(x_0, y) - u(x_0, y_0) + \nu(y - y_0)) \\ &= \left( \frac{\partial u}{\partial x}(x', y) - \mu \right)(x - x_0) + \left( \frac{\partial u}{\partial y}(x_0, y') + \nu \right)(y - y_0) \end{aligned}$$

and, because of (3.11),

$$\begin{aligned} &\left| u(x, y) - u(x_0, y_0) - (\mu(x - x_0) - \nu(y - y_0)) \right| \\ &\leq \left| \frac{\partial u}{\partial x}(x', y) - \mu \right| |x - x_0| + \left| \frac{\partial u}{\partial y}(x_0, y') + \nu \right| |y - y_0| \\ &< \frac{\epsilon}{4} |x - x_0| + \frac{\epsilon}{4} |y - y_0| < \frac{\epsilon}{2} \sqrt{(x - x_0)^2 + (y - y_0)^2}. \end{aligned} \quad (3.12)$$

In the same manner, for the function  $v$  we get

$$\left| v(x, y) - v(x_0, y_0) - (\nu(x - x_0) + \mu(y - y_0)) \right| < \frac{\epsilon}{2} \sqrt{(x - x_0)^2 + (y - y_0)^2}. \quad (3.13)$$

The inequalities (3.12) and (3.13) hold at every  $(x, y) \in D_{(x_0, y_0)}(r)$ .

We observe that the expressions inside the absolute values of the left sides of (3.12) and (3.13) are, respectively, the real and the imaginary part of the number

$$f(z) - f(z_0) - (\mu + i\nu)(z - z_0) = f(x, y) - f(x_0, y_0) - (\mu + i\nu)((x - x_0) + i(y - y_0)).$$

Therefore, (3.12) and (3.13) imply

$$|f(z) - f(z_0) - (\mu + i\nu)(z - z_0)| < \epsilon \sqrt{(x - x_0)^2 + (y - y_0)^2} = \epsilon |z - z_0|$$

for every  $z \in D_{z_0}(r)$  and hence  $|\frac{f(z)-f(z_0)}{z-z_0} - (\mu + i\nu)| < \epsilon$  for every  $z \in D_{z_0}(r), z \neq z_0$ . Thus,  $\lim_{z \rightarrow z_0} \frac{f(z)-f(z_0)}{z-z_0} = \mu + i\nu$ , and  $f$  is differentiable at  $z_0$  with  $f'(z_0) = \mu + i\nu$ .  $\square$

**Example 3.2.1.** The real and the imaginary parts of the function  $f(z) = z^2$  are  $u(x, y) = x^2 - y^2$  and  $v(x, y) = 2xy$ . We find  $\frac{\partial u}{\partial x}(x, y) = 2x, \frac{\partial u}{\partial y}(x, y) = -2y, \frac{\partial v}{\partial x}(x, y) = 2y$  and  $\frac{\partial v}{\partial y}(x, y) = 2x$  and we see that the partial derivatives are continuous in the whole plane and they satisfy the C-R equations at every point. Theorem 3.2 implies that  $f(z) = z^2$  is differentiable at every point and  $f'(z) = \frac{\partial u}{\partial x}(x, y) + i \frac{\partial v}{\partial x}(x, y) = 2x + i2y = 2z$ .

**Example 3.2.2.** We reconsider the function  $f(z) = \bar{z}$  of example 3.1.3. Its real and imaginary parts are  $u(x, y) = x$  and  $v(x, y) = -y$ . The partial derivatives  $\frac{\partial u}{\partial x}(x, y) = 1, \frac{\partial u}{\partial y}(x, y) = 0, \frac{\partial v}{\partial x}(x, y) = 0$  and  $\frac{\partial v}{\partial y}(x, y) = -1$  do not satisfy the C-R equations at any point  $(x, y)$ . Theorem 3.1 implies that  $f$  is not differentiable at any point.

**Example 3.2.3.** We reconsider the function  $f(z) = |z|^2$  of example 3.1.9. Its real and imaginary parts are  $u(x, y) = x^2 + y^2$  and  $v(x, y) = 0$ . The partial derivatives are  $\frac{\partial u}{\partial x}(x, y) = 2x, \frac{\partial u}{\partial y}(x, y) = 2y, \frac{\partial v}{\partial x}(x, y) = 0$  and  $\frac{\partial v}{\partial y}(x, y) = 0$  and they satisfy the C-R equations only at the point  $(0, 0)$ . Theorem 3.1 implies that  $f$  is not differentiable at any point besides, perhaps, the point  $(0, 0)$ . Now, since the partial derivatives are continuous and satisfy the C-R equations at  $(0, 0)$ , theorem 3.2 implies that  $f$  is differentiable at 0 and  $f'(0) = \frac{\partial u}{\partial x}(0, 0) + i \frac{\partial v}{\partial x}(0, 0) = 0 + i0 = 0$ .

**Example 3.2.4.** We shall see that the assumption of continuity of the partial derivatives of  $u, v$  at  $(x_0, y_0)$  in theorem 3.2 is crucial. We consider the function

$$f(z) = f(x, y) = \begin{cases} \frac{xy}{\sqrt{x^2+y^2}}, & \text{if } (x, y) \neq (0, 0) \\ 0, & \text{if } (x, y) = (0, 0) \end{cases}$$

Then its real and imaginary parts are

$$u(x, y) = \begin{cases} \frac{xy}{\sqrt{x^2+y^2}}, & \text{if } (x, y) \neq (0, 0) \\ 0, & \text{if } (x, y) = (0, 0) \end{cases} \quad v(x, y) = 0.$$

It is clear that  $\frac{\partial u}{\partial x}(x, y) = 0$  and  $\frac{\partial u}{\partial y}(x, y) = 0$  and the partial derivatives of  $v$  are continuous at every  $(x, y)$ . Moreover,

$$\frac{\partial u}{\partial x}(x, y) = \begin{cases} \frac{y^3}{\sqrt{(x^2+y^2)^3}}, & \text{if } (x, y) \neq (0, 0) \\ 0, & \text{if } (x, y) = (0, 0) \end{cases} \quad \frac{\partial u}{\partial y}(x, y) = \begin{cases} \frac{x^3}{\sqrt{(x^2+y^2)^3}}, & \text{if } (x, y) \neq (0, 0) \\ 0, & \text{if } (x, y) = (0, 0) \end{cases}$$

The partial derivatives of  $u$  are continuous at every  $(x, y) \neq (0, 0)$  but they are *not* continuous at  $(0, 0)$ . For instance, the limit of  $\frac{y^3}{\sqrt{(x^2+y^2)^3}}$  when  $(x, y)$  tends to  $(0, 0)$  on the line with equation  $y = x$  does not exist:  $\lim_{x \rightarrow \pm 0} \frac{x^3}{\sqrt{(x^2+x^2)^3}} = \pm 1/\sqrt{8}$ .

We will see now that  $f$  is *not* differentiable at 0, *even though*  $u, v$  do satisfy the C-R equations at 0. In fact the limit of  $\frac{f(z)-f(0)}{z-0} = \frac{(xy)/\sqrt{x^2+y^2}}{x+iy}$  when  $z$  tends to 0 on the line with equation  $y = x$  is  $\lim_{x \rightarrow 0} \frac{x^2/\sqrt{x^2+x^2}}{x+ix} = \frac{1}{(1+i)\sqrt{2}} \lim_{x \rightarrow 0} \frac{x}{|x|}$  and it does not exist.

The next proposition is a corollary of theorem 3.2. It is the form of theorem 3.2 in which this is usually applied.

**Proposition 3.7.** Let  $f$  be a complex function defined in the open set  $\Omega \subseteq \mathbb{C}$  and let  $u, v$  be the real and the imaginary part of  $f$ . If  $u, v$  have partial derivatives which are continuous and which satisfy the C-R equations at every point of  $\Omega$ , then  $f$  is holomorphic in  $\Omega$ .

*Proof.* We take an arbitrary  $z \in \Omega$  and a neighborhood of  $z$  which is contained in  $\Omega$ . Theorem 3.2 implies that  $f$  is differentiable at  $z$ . Thus  $f$  is differentiable at every point of  $\Omega$  and, since  $\Omega$  is open,  $f$  is holomorphic in  $\Omega$ .  $\square$

An open and connected set  $\Omega$  is called **region**.

**Theorem 3.3.** Let  $f$  be holomorphic in the region  $\Omega \subseteq \mathbb{C}$ . If  $f'(z) = 0$  for every  $z \in \Omega$ , then  $f$  is constant in  $\Omega$ .

*First proof.* Using  $f' = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = \frac{\partial v}{\partial y} - i \frac{\partial u}{\partial y}$ , we find  $\frac{\partial u}{\partial x} = \frac{\partial v}{\partial x} = \frac{\partial v}{\partial y} = \frac{\partial u}{\partial y} = 0$  in  $\Omega$ . We take any linear segment  $[z_1, z_2]$  in  $\Omega$  and its parametric equation  $\gamma(t) = (1-t)z_1 + tz_2, t \in [0, 1]$ . By the mean value theorem, there is  $t_0 \in (0, 1)$  so that

$$\begin{aligned} u(z_2) - u(z_1) &= (u \circ \gamma)(1) - (u \circ \gamma)(0) = \frac{d(u \circ \gamma)}{dt}(t_0) \\ &= \frac{\partial u}{\partial x}(\gamma(t_0))(x_2 - x_1) + \frac{\partial u}{\partial y}(\gamma(t_0))(y_2 - y_1) = 0, \end{aligned}$$

where  $z_1 = x_1 + iy_1$  and  $z_2 = x_2 + iy_2$ . Thus, the values of  $u$  at the endpoints of any line segment in  $\Omega$  are equal. Now we take arbitrary  $z', z'' \in \Omega$ . Then there is a polygonal line inside  $\Omega$  which connects the two points  $z'$  and  $z''$ . The values of  $u$  at the endpoints of every line segment of the polygonal line are equal and hence  $u(z') = u(z'')$ . Therefore  $u$  is constant in  $\Omega$ . Clearly, the same is true for the function  $v$  and hence for  $f = u + iv$ .

*Second proof.* We take arbitrary  $z, w \in \Omega$ . Since  $\Omega$  is a region, there is a piecewise smooth curve  $\gamma : [a, b] \rightarrow \Omega$  such that  $\gamma(a) = z, \gamma(b) = w$ . In fact we may choose  $\gamma$  to have a polygonal line in  $\Omega$  as its trajectory. Then we have

$$f(w) - f(z) = (f \circ \gamma)(b) - (f \circ \gamma)(a) = \int_a^b (f \circ \gamma)'(t) dt = \int_a^b f'(\gamma(t))\gamma'(t) dt = 0$$

because  $f'(\gamma(t)) = 0$  for every  $t \in [a, b]$ . We conclude that  $f(w) = f(z)$  for every  $w, z \in \Omega$  and hence  $f$  is constant in  $\Omega$ .  $\square$

Let  $f$  be a complex function and let  $u, v$  be the real and imaginary part of  $f$ . If  $u, v$  have partial derivatives with respect to  $x, y$  at the point  $z_0 = (x_0, y_0)$ , it is trivial to prove that at the point  $z_0$  we have

$$\frac{\partial f}{\partial x} = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}, \quad \frac{\partial f}{\partial y} = \frac{\partial u}{\partial y} + i \frac{\partial v}{\partial y}. \quad (3.14)$$

We define the following differential operators:

$$\frac{\partial}{\partial z} = \frac{1}{2} \left( \frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right), \quad \frac{\partial}{\partial \bar{z}} = \frac{1}{2} \left( \frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right). \quad (3.15)$$

Applying the differential operators  $\frac{\partial}{\partial z}$  and  $\frac{\partial}{\partial \bar{z}}$  to  $f$  and using (3.14), we have at the point  $z_0$ :

$$\begin{aligned} \frac{\partial f}{\partial z} &= \frac{1}{2} \left( \frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right), \\ \frac{\partial f}{\partial \bar{z}} &= \frac{1}{2} \left( \frac{\partial u}{\partial x} - \frac{\partial v}{\partial y} \right) + \frac{i}{2} \left( \frac{\partial v}{\partial x} + \frac{\partial u}{\partial y} \right). \end{aligned} \quad (3.16)$$

From the second of equations (3.16) we see that the system of C-R equations at the point  $z_0$  is equivalent to the single equation

$$\frac{\partial f}{\partial \bar{z}} = 0$$

at  $z_0$ . Moreover, if the system of C-R equations is satisfied, then the first equation (3.16) implies

$$\frac{\partial f}{\partial z} = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = f'$$

at  $z_0$ . We summarize.

**Proposition 3.8.** *If the complex function  $f$  is differentiable at  $z_0$ , then  $\frac{\partial f}{\partial z}(z_0) = f'(z_0)$  and  $\frac{\partial f}{\partial \bar{z}}(z_0) = 0$ . Conversely, if  $\frac{\partial f}{\partial z}$  and  $\frac{\partial f}{\partial \bar{z}}$  exist in a neighborhood of the point  $z_0$  and they are continuous at  $z_0$  and if  $\frac{\partial f}{\partial \bar{z}}(z_0) = 0$ , then  $f$  is differentiable at  $z_0$ .*

*Proof.* Trivial. The converse is a restatement of theorem 3.2. Indeed, (3.16) implies that the existence or the continuity of  $\frac{\partial f}{\partial z}$ ,  $\frac{\partial f}{\partial \bar{z}}$  at a point is equivalent to the existence or the continuity, respectively, of  $\frac{\partial u}{\partial x}$ ,  $\frac{\partial u}{\partial y}$ ,  $\frac{\partial v}{\partial x}$ ,  $\frac{\partial v}{\partial y}$  at the point.  $\square$

Sometimes a complex function  $f$  is given to us through an expression  $f(x, y)$  as a function of two real variables and we are interested in finding an expression  $f(z)$  of the function in terms of the single complex variable  $z$ . We then write  $x = \frac{z+\bar{z}}{2}$ ,  $y = \frac{z-\bar{z}}{2i}$  and hence

$$f(x, y) = f\left(\frac{z+\bar{z}}{2}, \frac{z-\bar{z}}{2i}\right). \quad (3.17)$$

In general, even after performing various algebraic simplifications we end up with an expression in terms of *both* variables  $z$  and  $\bar{z}$ . In order to end up with the occurrence of  $z$  only, it is reasonable to impose the condition that the derivative of  $f(x, y)$  with respect to  $\bar{z}$  vanishes. From (3.17) and a formal chain rule we get

$$\frac{\partial f}{\partial \bar{z}} = \frac{1}{2}\left(\frac{\partial f}{\partial x} + i\frac{\partial f}{\partial y}\right).$$

This is exactly the second differential operator (3.15) applied to  $f$  and we saw that the condition  $\frac{\partial f}{\partial \bar{z}} = 0$  is equivalent to the system of C-R equations. We conclude that *the function  $f(x, y)$  is a function of the single variable  $z$  if and only if its real and imaginary parts satisfy the C-R equations.*

### Exercises.

**3.2.1.** Solve exercise 3.1.1 under the light of C-R equations.

**3.2.2.** (i) Prove that  $F(x, y) = \sqrt{|xy|}$  satisfies the C-R equations at 0 but that it is not differentiable at 0.

(ii) Prove that the function with  $G(x, y) = \frac{x^2y}{x^4+y^2}$  if  $(x, y) \neq (0, 0)$  and with  $G(0, 0) = 0$  satisfies the C-R equations at 0, that  $\frac{G(z)-G(0)}{z-0}$  has a limit when  $z \rightarrow 0$  on every line which contains 0, but that  $G$  is not differentiable at 0.

**3.2.3.** Let  $f = u + iv$  be a complex function and  $\frac{\partial u}{\partial x}$ ,  $\frac{\partial u}{\partial y}$ ,  $\frac{\partial v}{\partial x}$ ,  $\frac{\partial v}{\partial y}$  exist in a neighborhood of  $z_0$  and be continuous at  $z_0$ .

(i) If  $\lim_{z \rightarrow z_0} \operatorname{Re} \frac{f(z)-f(z_0)}{z-z_0}$  exists and is a real number, prove that  $f$  is differentiable at  $z_0$ .

(ii) If  $\lim_{z \rightarrow z_0} \left| \frac{f(z)-f(z_0)}{z-z_0} \right|$  exists and is a real number, prove that either  $f$  is differentiable at  $z_0$  or  $\bar{f}$  is differentiable at  $z_0$ .

**3.2.4.** Let  $f = u + iv$  be holomorphic in the region  $\Omega \subseteq \mathbb{C}$ .

(i) If either  $u$  or  $v$  is constant in  $\Omega$ , prove that  $f$  is constant in  $\Omega$ .

(ii) More generally, if for some line  $l$  it is true that  $f(z) \in l$  for every  $z \in \Omega$ , prove that  $f$  is constant in  $\Omega$ .

(iii) Consider (ii) using a circle  $C$  instead of a line  $l$ .

**3.2.5.** This exercise juxtaposes the notion of differentiability of a function of two real variables, which we learn in multivariable calculus, and the notion of differentiability of a function of one complex variable, which we learn in complex analysis: to distinguish between them we call the first  $\mathbb{R}$ -differentiability and the second  $\mathbb{C}$ -differentiability.

We recall from multivariable calculus that a *real* function  $u$  defined in  $A \subseteq \mathbb{R}^2$  is  $\mathbb{R}$ -differentiable at the interior point  $(x_0, y_0)$  of  $A$  if there are  $a, b \in \mathbb{R}$  so that

$$\lim_{(x,y) \rightarrow (x_0,y_0)} \frac{u(x,y)-u(x_0,y_0)-(a(x-x_0)+b(y-y_0))}{\sqrt{(x-x_0)^2+(y-y_0)^2}} = 0.$$

In this case we have that  $\frac{\partial u}{\partial x}(x_0, y_0) = a$  and  $\frac{\partial u}{\partial y}(x_0, y_0) = b$ .

We also recall that a *vector* function  $f = (u, v)$  defined in  $A \subseteq \mathbb{R}^2$  is  $\mathbb{R}$ -differentiable at the interior point  $(x_0, y_0)$  of  $A$  if its real components  $u$  and  $v$  are both  $\mathbb{R}$ -differentiable at  $(x_0, y_0)$ , i.e. if there are  $a, b, c, d \in \mathbb{R}$  so that

$$\lim_{(x,y) \rightarrow (x_0,y_0)} \frac{u(x,y) - u(x_0,y_0) - (a(x-x_0) + b(y-y_0))}{\sqrt{(x-x_0)^2 + (y-y_0)^2}} = 0,$$

$$\lim_{(x,y) \rightarrow (x_0,y_0)} \frac{v(x,y) - v(x_0,y_0) - (c(x-x_0) + d(y-y_0))}{\sqrt{(x-x_0)^2 + (y-y_0)^2}} = 0.$$

In this case we have that  $\frac{\partial u}{\partial x}(x_0, y_0) = a$ ,  $\frac{\partial u}{\partial y}(x_0, y_0) = b$ ,  $\frac{\partial v}{\partial x}(x_0, y_0) = c$ ,  $\frac{\partial v}{\partial y}(x_0, y_0) = d$  and that the  $\mathbb{R}$ -derivative of  $f$  is the  $2 \times 2$  matrix  $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$ .

Prove that  $f = (u, v) = u + iv$  is  $\mathbb{C}$ -differentiable at  $z_0 = (x_0, y_0)$ , i.e. that the  $\lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0}$  exists and is a complex number, if and only if  $f$  is  $\mathbb{R}$ -differentiable at  $z_0 = (x_0, y_0)$  and its  $\mathbb{R}$ -derivative is an antisymmetric matrix:  $\begin{bmatrix} a & -b \\ b & a \end{bmatrix}$ . In this case the  $\mathbb{C}$ -derivative and the  $\mathbb{R}$ -derivative of  $f$  are related by  $f'(z_0) = a + ib$ .

**3.2.6.** Consider the functions  $z^n, \bar{z}^n, |z|^2$  and, using the differential operator  $\frac{\partial}{\partial \bar{z}}$ , examine whether they are functions of  $z$  only or, equivalently, whether they are holomorphic.

**3.2.7.** Let  $f$  be a complex function. If  $\frac{\partial f}{\partial x}$  and  $\frac{\partial f}{\partial y}$  exist in a neighborhood of the point  $z_0$  and are continuous at  $z_0$ , prove that  $\lim_{r \rightarrow 0} \frac{1}{2\pi i r^2} \int_{C_{z_0}(r)} f(z) dz = \frac{\partial f}{\partial \bar{z}}(z_0)$ .

### 3.3 Conformality.

Let the complex function  $f$  be continuous in  $A \subseteq \mathbb{C}$  and  $\gamma : [a, b] \rightarrow A$  be a curve. Thus the trajectory of  $\gamma$  is contained in the domain of definition of  $f$ . We define the function

$$f(\gamma) = f \circ \gamma : [a, b] \rightarrow \mathbb{C},$$

which is continuous in  $[a, b]$ . Then  $f(\gamma)$  is a curve and we call it **image of  $\gamma$  through  $f$** .

Now we also consider an interior point  $z$  of  $A$  and we assume that  $f$  is differentiable at  $z$  and  $f(z) = w$ ,  $f'(z) \neq 0$ . We also take any curve  $\gamma : [a, b] \rightarrow A$  with  $\gamma(a) = z$ . Then  $\gamma$  has  $z$  as its initial point and its trajectory is contained in  $A$ . We also assume that  $\gamma$  is differentiable at  $a$  and that  $\gamma'(a) \neq 0$ , i.e. that  $\gamma$  has a non-zero tangent vector at the point  $z$ . The image curve  $f(\gamma) : [a, b] \rightarrow \mathbb{C}$  has  $f(\gamma)(a) = (f \circ \gamma)(a) = f(\gamma(a)) = f(z) = w$  as its initial point and its tangent vector at  $w$  is  $f(\gamma)'(a) = (f \circ \gamma)'(a) = f'(\gamma(a))\gamma'(a) = f'(z)\gamma'(a) \neq 0$ . From this equality we have two conclusions. The first is that

$$|f(\gamma)'(a)| = |f'(z)||\gamma'(a)|.$$

Thus, the length of the tangent vector of  $f(\gamma)$  at its initial point  $w$  equals the length of the tangent vector of  $\gamma$  at its initial point  $z$  multiplied with the factor  $|f'(z)| > 0$ . We express this as:

*$f$  multiplies the lengths of tangent vectors at  $z$  with the factor  $|f'(z)| > 0$  or, in other words,  $f$  expands the tangent vectors at  $z$  by the factor  $|f'(z)| > 0$ .*

The second conclusion is that

$$\arg f(\gamma)'(a) = \arg f'(z) + \arg \gamma'(a). \quad (3.18)$$

Thus, we find the angle of the tangent vector of  $f(\gamma)$  at its initial point  $w$  by adding the angle of  $f'(z)$  to the angle of the tangent vector of  $\gamma$  at its initial point  $z$ . We express this as:

$f$  increases the angles of the tangent vectors at  $z$  by the angle of  $f'(z)$  or, in other words,  $f$  rotates the tangent vectors at  $z$  by the angle of  $f'(z)$ .

We observe that the expansion and the rotation of the tangent vectors at  $z$  is uniform over all these vectors: *independently of their directions and their lengths, all these tangent vectors are expanded by the same factor  $|f'(z)|$  and they are rotated by the same angle  $\arg f'(z)$* . Since, any two of these tangent vectors are rotated by the same angle, we conclude that their relative angles remain unchanged! Indeed, let us consider two of the above curves,  $\gamma_1$  and  $\gamma_2$ . Then the angle between their tangent vectors at  $z$  is  $\arg \gamma_2'(a) - \arg \gamma_1'(a)$  and the angle between the tangent vectors of  $f(\gamma_1)$  and  $f(\gamma_2)$  at  $w$  is  $\arg f(\gamma_2)'(a) - \arg f(\gamma_1)'(a)$ . From (3.18) for  $\gamma_1$  and  $\gamma_2$  we get

$$\arg f(\gamma_2)'(a) - \arg f(\gamma_1)'(a) = \arg \gamma_2'(a) - \arg \gamma_1'(a).$$

Therefore, the angle between the tangent vectors of  $f(\gamma_1)$  and  $f(\gamma_2)$  at  $w$  is the same as the angle between the tangent vectors of  $\gamma_1$  and  $\gamma_2$  at  $z$ . We say:

$f$  preserves the angle between tangent vectors at  $z$ .

This last property of  $f$  is called **conformality** of  $f$  at  $z$  and holds, as we just saw, under the assumption that  $f$  is differentiable at  $z$  and  $f'(z) \neq 0$ .

### Exercises.

**3.3.1.** Consider the holomorphic function  $w = f(z) = az + b$  with  $a \neq 0$ .

(i) Prove that  $f$  is one-to-one from  $\mathbb{C}$  onto  $\mathbb{C}$ .

(ii) Prove that  $f$  maps lines and circles onto lines and circles, respectively.

(iii) Consider two lines with equations  $kx + ly = m$  and  $k'x + l'y = m'$ . Which is the condition for the two lines to intersect? Under this condition, find their intersection point and the angle of the two lines at this point. Then find the equations of the images of the two lines through  $f$  and find their intersection point and their angle at this point. Confirm the conformality of  $f$ .

**3.3.2.** Consider the holomorphic function  $w = z^2$ .

(i) With any fixed  $u_0, v_0$ , consider the hyperbolas with equations  $x^2 - y^2 = u_0$  and  $2xy = v_0$  on the  $z$ -plane ( $z = x + iy$ ). Do they intersect and at which points? Find the angle of the two hyperbolas at each of their common points.

(ii) With any fixed  $x_0, y_0 \neq 0$ , consider the parabolas with equations  $u = \frac{1}{4y_0^2}v^2 - y_0^2$  and  $u = -\frac{1}{4x_0^2}v^2 + x_0^2$  on the  $w$ -plane ( $w = u + iv$ ). Do they intersect and at which points? Find the angle of the two parabolas at each of their common points.

**3.3.3.** Let  $f$  be holomorphic in the open set  $U \subseteq \mathbb{C}$  so that  $f'$  is continuous in  $U$ , let  $\gamma$  be a piecewise smooth curve in  $U$  and  $\Gamma = f(\gamma)$  be the image of  $\gamma$  through  $f$ . If the complex function  $\phi$  is continuous in  $\Gamma^*$ , prove that  $\int_{\Gamma} \phi(w) dw = \int_{\gamma} \phi(f(z))f'(z) dz$ .

## Chapter 4

# Examples of holomorphic functions.

### 4.1 Linear fractional transformations.

Every rational function of the form

$$T(z) = \frac{az+b}{cz+d}$$

is called **linear fractional transformation**. We assume that  $ad - bc \neq 0$ . It is easy to show that  $ad - bc \neq 0$  if and only if the function  $T$  is not constant.

In order to have the full picture of the definition of a linear fractional transformation  $T$ , we have to say something about the values of  $T$  at the roots of the denominator and at  $\infty$ . There are two cases. If  $c = 0$ , then because of  $ad - bc \neq 0$  we have  $ad \neq 0$  and then  $T(z) = \frac{a}{d}z + \frac{b}{d}$  for all  $z \in \mathbb{C}$ . Since  $\frac{a}{d} \neq 0$ , we have that  $T(\infty) = \infty$ . Thus

$$T(z) = \begin{cases} \frac{a}{d}z + \frac{b}{d}, & \text{if } z \in \mathbb{C} \\ \infty, & \text{if } z = \infty \end{cases} \quad \text{if } c = 0. \quad (4.1)$$

If  $c \neq 0$ , then the denominator has  $z = -\frac{d}{c}$  as its root, which, because of  $ad - bc \neq 0$ , is not a root of the numerator. Hence  $T(-\frac{d}{c}) = \infty$ . Also  $T(\infty) = \frac{a}{c}$ . Thus

$$T(z) = \begin{cases} \frac{az+b}{cz+d}, & \text{if } z \in \mathbb{C}, z \neq -\frac{d}{c} \\ \infty, & \text{if } z = -\frac{d}{c} \\ \frac{a}{c}, & \text{if } z = \infty \end{cases} \quad \text{if } c \neq 0. \quad (4.2)$$

We conclude that every linear fractional transformation (l.f.t.) is a function  $T : \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$  and, even though we write  $T(z) = \frac{az+b}{cz+d}$ , we must have in mind the full formulas (4.1) and (4.2).

It is very easy to see that every l.f.t. is one-to-one from  $\widehat{\mathbb{C}}$  onto  $\widehat{\mathbb{C}}$ . The formula of the inverse l.f.t. of  $T$  is

$$T^{-1}(z) = \frac{dz-b}{-cz+a}.$$

The identity function  $id(z) = z$  is clearly a l.f.t. with  $a = d = 1, b = c = 0$ , and we easily see that the composition of two l.f.t. is another l.f.t. Indeed, if  $T(z) = \frac{az+b}{cz+d}$  and  $S(z) = \frac{a'z+b'}{c'z+d'}$ , then

$$(S \circ T)(z) = \frac{a'T(z)+b'}{c'T(z)+d'} = \frac{a'\frac{az+b}{cz+d}+b'}{c'\frac{az+b}{cz+d}+d'} = \frac{(a'a+b'c)z+(a'b+b'd)}{(c'a+d'c)z+(c'b+d'd)}.$$

We observe that  $(a'a + b'c)(c'b + d'd) - (a'b + b'd)(c'a + d'c) = (a'd' - b'c')(ad - bc) \neq 0$ .

Thus, the set of all l.f.t. is a group with the binary operation of composition. The neutral element of this group is the identity function.

Since a l.f.t. is a rational function, it is continuous in  $\widehat{\mathbb{C}}$ , and, as a particular instance of example 3.1.13, it is holomorphic in  $\widehat{\mathbb{C}}$  except at the point at which it takes the value  $\infty$ .

We shall make a comment on an interesting relation between circles and lines. We observe that the equations of circles and lines can be unified in the following manner: if  $\alpha, \beta, \gamma \in \mathbb{R}$ ,  $w \in \mathbb{C}$ ,  $w \neq 0$ ,  $\alpha^2 + \beta^2 \neq 0$  and  $\beta^2|w|^2 \geq 4\alpha\gamma$ , then the equation

$$\alpha|z|^2 + \beta \operatorname{Re}(\bar{w}z) + \gamma = 0$$

is the equation of a line, if  $\alpha = 0$ , and the equation of a circle, if  $\alpha \neq 0$ . In fact, if  $\alpha = 0$ , then  $\beta \neq 0$  and the equation becomes  $\operatorname{Re}(\bar{w}z) = -\frac{\gamma}{\beta}$  and this is the equation of a line. If  $\alpha \neq 0$ , the equation becomes  $|z + \frac{\beta}{2\alpha}w|^2 = \frac{\beta^2|w|^2 - 4\alpha\gamma}{4\alpha^2}$ . This is the equation of the circle with center  $-\frac{\beta}{2\alpha}w$  and radius  $\frac{\sqrt{\beta^2|w|^2 - 4\alpha\gamma}}{2|\alpha|}$ . Conversely, every circle and every line have equations of this form. If, for instance, we take the equation  $\operatorname{Re}(\bar{w}z) = c$  of a line, with  $w \in \mathbb{C}$ ,  $w \neq 0$ , and  $c \in \mathbb{R}$ , we may write it in the form  $\alpha|z|^2 + \beta \operatorname{Re}(\bar{w}z) + \gamma = 0$  by taking  $\alpha = 0$ ,  $\beta = 1$  and  $\gamma = -c$ . If we take the equation  $|z - z_0| = r$  of a circle with  $z_0 \in \mathbb{C}$  and  $r \geq 0$ , we may write it as  $|z|^2 - 2 \operatorname{Re}(\bar{z}_0z) + |z_0|^2 = r^2$ . This becomes  $\alpha|z|^2 + \beta \operatorname{Re}(\bar{w}z) + \gamma = 0$  by taking  $\alpha = 1$ ,  $\gamma = |z_0|^2 - r^2$  and:  $\beta = -2$  and  $w = z_0$ , in case  $z_0 \neq 0$ , or  $\beta = 0$  and  $w = 1$ , in case  $z_0 = 0$ . In all cases the choices of the parameters satisfy the restrictions:  $\alpha, \beta, \gamma \in \mathbb{R}$ ,  $w \in \mathbb{C}$ ,  $w \neq 0$ ,  $\alpha^2 + \beta^2 \neq 0$  and  $\beta^2|w|^2 \geq 4\alpha\gamma$ .

This consideration of the equations of a line and a circle as special cases of one equation permits us to unify the notions of circle and line into the single notion of **generalized circle** in  $\mathbb{C}$ . If we attach the point  $\infty$  to any line (and leave circles unchanged), then we are talking about generalized circles in  $\widehat{\mathbb{C}}$ . Look at exercise 1.3.2 for another interesting unification of the notions of circle and line: *generalized circles in  $\widehat{\mathbb{C}}$  are the images of circles in  $\mathbb{S}^2$  through stereographic projection.*

Now, an important property of every l.f.t. is that it maps generalized circles in  $\widehat{\mathbb{C}}$  onto generalized circles in  $\widehat{\mathbb{C}}$ . To prove it we consider three special cases first.

**Example 4.1.1.** Every function  $T(z) = z + b$  is a l.f.t. with  $a = 1$ ,  $c = 0$ ,  $d = 1$  and, for an obvious reason, it is called **translation** by  $b$ .

Every such  $T$  is holomorphic in  $\mathbb{C}$ , one-to-one from  $\mathbb{C}$  onto  $\mathbb{C}$  and satisfies  $T(\infty) = \infty$ . It is trivial to prove that  $T$  maps lines in  $\widehat{\mathbb{C}}$  onto lines in  $\widehat{\mathbb{C}}$  and circles in  $\mathbb{C}$  onto circles in  $\mathbb{C}$ .

**Example 4.1.2.** Every function  $T(z) = az$  with  $a \neq 0$  is a l.f.t. with  $b = c = 0$ ,  $d = 1$  and it is called **homothety** with center 0.

Every such  $T$  rotates points around 0 by the fixed angle  $\arg a$ . Indeed, if  $w = T(z) = az$ , then  $\arg w = \arg z + \arg a$ . Moreover,  $T$  multiplies distances between points by the fixed factor  $|a|$ . Indeed, if  $w_1 = T(z_1) = az_1$  and  $w_2 = T(z_2) = az_2$ , then  $|w_1 - w_2| = |a||z_1 - z_2|$ . Also  $T$  is holomorphic in  $\mathbb{C}$ , one-to-one from  $\mathbb{C}$  onto  $\mathbb{C}$ , satisfies  $T(\infty) = \infty$  and it is easy to prove that  $T$  maps lines in  $\widehat{\mathbb{C}}$  onto lines in  $\widehat{\mathbb{C}}$  and circles in  $\mathbb{C}$  onto circles in  $\mathbb{C}$ .

**Example 4.1.3.** The function  $T(z) = \frac{1}{z}$  is a l.f.t. with  $a = d = 0$ ,  $c = b = 1$  and it is called **inversion** with respect to the circle  $\mathbb{T} = C_0(1)$ .

The inversion  $T$  is holomorphic in  $\widehat{\mathbb{C}} \setminus \{0\}$ , one-to-one from  $\widehat{\mathbb{C}} \setminus \{0, \infty\}$  onto  $\widehat{\mathbb{C}} \setminus \{0, \infty\}$  and satisfies  $T(0) = \infty$  and  $T(\infty) = 0$ . Moreover, it is easy to show that  $T$  maps (i) lines in  $\widehat{\mathbb{C}}$  which do not contain 0 onto circles in  $\mathbb{C}$  which contain 0, (ii) lines in  $\widehat{\mathbb{C}}$  which contain 0 onto lines in  $\widehat{\mathbb{C}}$  which contain 0, (iii) circles in  $\mathbb{C}$  which contain 0 onto lines in  $\widehat{\mathbb{C}}$  which do not contain 0 and (iv) circles in  $\mathbb{C}$  which do not contain 0 onto circles in  $\mathbb{C}$  which do not contain 0.

**Lemma 4.1.** *Every l.f.t. is a composition of finitely many translations, homotheties and inversions.*

*Proof.* Let  $T(z) = \frac{az+b}{cz+d}$ .

If  $c = 0$ , then  $T(z) = \frac{a}{d}z + \frac{b}{d}$ . If we consider the homothety  $T_1(z) = \frac{a}{d}z$  and the translation  $T_2(z) = z + \frac{b}{d}$ , then  $T = T_2 \circ T_1$ .

If  $c \neq 0$ , then

$$T(z) = \frac{\frac{a}{c}(cz+d) + (b - \frac{ad}{c})}{cz+d} = \frac{a}{c} + \frac{bc-ad}{c^2} \frac{1}{z + \frac{d}{c}}.$$



If we consider the translation  $T_1(z) = z + \frac{d}{c}$ , the inversion  $T_2(z) = \frac{1}{z}$ , the homothety  $T_3(z) = \frac{bc-ad}{c^2}z$  and the translation  $T_4(z) = z + \frac{a}{c}$ , then  $T = T_4 \circ T_3 \circ T_2 \circ T_1$ .  $\square$

**Proposition 4.1.** Every l.f.t. maps generalized circles in  $\widehat{\mathbb{C}}$  onto generalized circles in  $\widehat{\mathbb{C}}$ .

*Proof.* A corollary of lemma 4.1 and of the examples 4.1.1, 4.1.2 and 4.1.3.  $\square$

**Proposition 4.2.** Take the distinct  $z_1, z_2, z_3 \in \widehat{\mathbb{C}}$  and the distinct  $w_1, w_2, w_3 \in \widehat{\mathbb{C}}$ . Then there is a unique l.f.t.  $T$  so that  $T(z_j) = w_j$  for  $j = 1, 2, 3$ .

*Proof.* We consider the l.f.t.  $S$  which, depending on whether one of  $z_1, z_2, z_3$  is  $\infty$  or not, has the formula

$$S(z) = \begin{cases} \frac{z_2 - z_3}{z_2 - z_1} \frac{z - z_1}{z - z_3}, & \text{if } z_1, z_2, z_3 \neq \infty \\ \frac{z - z_1}{z_2 - z_1}, & \text{if } z_3 = \infty \\ \frac{z - z_1}{z - z_3}, & \text{if } z_2 = \infty \\ \frac{z_2 - z_3}{z - z_3}, & \text{if } z_1 = \infty \end{cases}$$

The l.f.t.  $S$  has values:  $S(z_1) = 0, S(z_2) = 1, S(z_3) = \infty$ .

There is a similar l.f.t.  $R$  with values:  $R(w_1) = 0, R(w_2) = 1, R(w_3) = \infty$ .

Then the l.f.t.  $T = R^{-1} \circ S$  has values:  $T(z_1) = w_1, T(z_2) = w_2, T(z_3) = w_3$ .

To prove the uniqueness of  $T$  with  $T(z_1) = w_1, T(z_2) = w_2, T(z_3) = w_3$  we consider the previous l.f.t.  $S, R$  and then the l.f.t.  $Q = R \circ T \circ S^{-1}$  has values:  $Q(0) = 0, Q(1) = 1, Q(\infty) = \infty$ . Since  $Q(\infty) = \infty$ , we get that  $Q$  has the form  $Q(z) = az + b$  with  $a \neq 0$ . Now from  $Q(0) = 0, Q(1) = 1$  we find  $a = 1, b = 0$  and hence  $Q$  is the identity l.f.t.  $id$  with  $id(z) = z$ . Thus  $R \circ T \circ S^{-1} = id$  and hence  $T = R^{-1} \circ S$ .  $\square$

When we apply the previous results we should bear in mind that every three distinct points in  $\widehat{\mathbb{C}}$  belong to a unique generalized circle in  $\widehat{\mathbb{C}}$ .

**Example 4.1.4.** The l.f.t. which maps the triple  $i, 2, 1$  onto the triple  $0, 1, \infty$  is

$$w = T(z) = \frac{2-i}{2-i} \frac{z-i}{z-1} = \frac{2+i}{5} \frac{z-i}{z-1} = \frac{(2+i)z + (1-2i)}{5z-5}.$$

The points  $i, 2, 1$  in the  $z$ -plane are not co-linear and hence belong to a circle  $A$ . The points  $0, 1$  in the  $w$ -plane belong to the real axis  $m$ . Thus the points  $0, 1, \infty$  belong to the line  $B = m \cup \{\infty\}$  in  $\widehat{\mathbb{C}}$ . Now,  $T$  maps the circle  $A$  in the  $z$ -plane onto some generalized circle  $T(A)$  in the  $w$ -plane. Since  $A$  contains  $i, 2, 1$ ,  $T(A)$  must contain the images of  $i, 2, 1$ , i.e.  $0, 1, \infty$ . Thus  $T(A) = B$ . If we want to determine the circle  $A = C_{z_0}(r)$  which contains  $i, 2, 1$ , we have to find  $z_0, r$  so that  $i, 2, 1$  satisfy the equation  $|z - z_0| = r$ . We just solve a system of three equations in three real unknowns:  $x_0, y_0, r$ . But there is a second and probably easier way to find the equation of  $A$ . Indeed,  $w$  belongs to  $m$  if and only if  $\text{Im } w = 0$  if and only if  $\text{Im} \frac{(2+i)z + (1-2i)}{5z-5} = 0$  (and  $z \neq 1$ ) if and only if  $|z|^2 - 3 \text{Re}((1-i)z) = -2$  (and  $z \neq 1$ ) if and only if  $|z - \frac{3}{2}(1+i)|^2 = -2 + \frac{9}{4}|1+i|^2 = \frac{5}{2}$  (and  $z \neq 1$ ) if and only if  $z$  belongs to  $C_{3(1+i)/2}(\sqrt{5/2})$  except 1. Since  $z = 1$  is mapped onto  $w = \infty$ , we have that  $w$  belongs to  $B$  if and only if  $z$  belongs to the circle  $C_{3(1+i)/2}(\sqrt{5/2})$ . We conclude that  $A = C_{3(1+i)/2}(\sqrt{5/2})$ .

### Exercises.

**4.1.1.** Find l.f.t.  $T$  so that  $T(1) = i, T(i) = 0, T(-1) = -i$ . Find  $T(\mathbb{T})$  and  $T(\mathbb{D})$ .

**4.1.2.** Find l.f.t.  $T$  so that  $T(\mathbb{D}) = \{z \mid \text{Im } z > 0\}, T(i) = 1, T(1) = 0, T(a) = -1$ , where  $a \in \mathbb{T}$ . Can  $a$  be an arbitrary point of  $\mathbb{T}$ ?

**4.1.3.** (i) Let  $T_1(z) = \frac{a_1z+b_1}{c_1z+d_1}$  and  $T_2(z) = \frac{a_2z+b_2}{c_2z+d_2}$ . Prove that  $T_1, T_2$  are the same function if and only if there is  $\lambda \neq 0$  so that  $a_2 = \lambda a_1, b_2 = \lambda b_1, c_2 = \lambda c_1, d_2 = \lambda d_1$ .

(ii) Prove that every l.f.t.  $T$  can take the form  $T(z) = \frac{az+b}{cz+d}$  with  $ad - bc = 1$ .

**4.1.4.** Let  $A$  be a generalized circle of the  $z$ -plane  $\widehat{\mathbb{C}}$  and  $B$  be a generalized circle of the  $w$ -plane  $\widehat{\mathbb{C}}$ . Then, in an obvious way,  $A$  splits  $\widehat{\mathbb{C}}$  into two disjoint sets  $A_+$  and  $A_-$  and, similarly,  $B$  splits  $\widehat{\mathbb{C}}$  into two disjoint sets  $B_+$  and  $B_-$ . Now, let  $T$  be a l.f.t. and let  $T(A) = B$ . Assume that  $z_0 \in A_+$  and  $w_0 = T(z_0) \in B_+$ . Prove that  $T(A_+) = B_+$  and  $T(A_-) = B_-$ .

**4.1.5.** A point  $z \in \widehat{\mathbb{C}}$  is called **fixed point** of the l.f.t.  $T$  if  $T(z) = z$ . If the l.f.t.  $T$  is not the identity (in which case  $T$  has infinitely many fixed points), prove that  $T$  has either one or two fixed points in  $\widehat{\mathbb{C}}$ . In each case, which are the images through  $T$  of the generalized circles which contain its fixed points?

Apply the above to each of:  $T(z) = z + 2, T(z) = 2z - 1, T(z) = \frac{z-1}{z+1}$  and  $T(z) = \frac{3z-4}{z-1}$ .

**4.1.6.** (i) The points  $a, b \in \widehat{\mathbb{C}}$  are called **symmetric** with respect to  $C_{z_0}(r)$  if either  $a = z_0, b = \infty$  or  $a = \infty, b = z_0$  or  $a, b \in \mathbb{C}$  are on the same halfline with vertex  $z_0$  and  $|a - z_0||b - z_0| = r^2$ . Observe that either  $a, b$  coincide with one and the same point of  $C_{z_0}(r)$  or  $a, b$  are on different sides of  $C_{z_0}(r)$ . Given  $a \in \widehat{\mathbb{C}} \setminus \{z_0, \infty\}$ , describe a geometric construction “with ruler and compass” of its symmetric point,  $b \in \widehat{\mathbb{C}} \setminus \{z_0, \infty\}$ , with respect to  $C_{z_0}(r)$ . Prove that  $a, b$  are symmetric with respect to  $C_{z_0}(r)$  if and only if  $b = z_0 + \frac{r^2}{a - z_0}$ .

(ii) The points  $a, b \in \widehat{\mathbb{C}}$  are called **symmetric** with respect to the line  $\widehat{l} = l \cup \{\infty\}$  in  $\widehat{\mathbb{C}}$  if either  $a = b = \infty$  or  $a, b \in \mathbb{C}$  are symmetric with respect to  $l$ . Prove that  $a, b$  are symmetric with respect to  $\widehat{l}$  if and only if  $b = z_1 + \frac{z_2 - z_1}{z_2 - \bar{z}_1}(\bar{a} - \bar{z}_1)$ , where  $z_1, z_2$  are two distinct fixed points of the line  $l$ .

(iii) We take a l.f.t.  $w = T(z)$  and generalized circles  $A$  in the  $z$ -plane  $\widehat{\mathbb{C}}$  and  $B$  in the  $w$ -plane  $\widehat{\mathbb{C}}$ . Prove that, if  $T$  maps  $A$  onto  $B$ , then  $T$  maps symmetric points with respect to  $A$  onto symmetric points with respect to  $B$ .

(iv) Find l.f.t.  $T$  so that  $T(C_0(1)) = C_i(3), T(i) = 3 + i, T(\frac{1}{2}) = 0$ .

**4.1.7.** The l.f.t.  $w = T(z)$  is called **real** if it maps the real line (with  $\infty$ ) in the  $z$ -plane  $\widehat{\mathbb{C}}$  onto the real line (with  $\infty$ ) in the  $w$ -plane  $\widehat{\mathbb{C}}$ .

(i) Prove that the l.f.t.  $T$  is real if and only if there are  $a, b, c, d \in \mathbb{R}$  with  $ad - bc \neq 0$  so that  $T(z) = \frac{az+b}{cz+d}$ .

(ii) If the l.f.t.  $T$  is real and  $T(z) = \frac{az+b}{cz+d}$ , with  $a, b, c, d \in \mathbb{R}, ad - bc \neq 0$ , we define  $\text{sign } T$  to be the sign of  $ad - bc$ . Using exercise 4.1.3(i), prove that  $\text{sign } T$  is well defined.

(iii) Prove that, if the l.f.t.  $T$  is real, then  $T^{-1}$  is real, and that, if the l.f.t.  $S, T$  are real, then  $S \circ T$  is real. Also prove that  $\text{sign } T^{-1} = \text{sign } T$  and  $\text{sign}(S \circ T) = \text{sign } S \text{ sign } T$ .

(iv) Take a real l.f.t.  $T$ . Prove that  $T$  maps the upper halfplane onto the upper halfplane (and the lower onto the lower) if and only if  $\text{sign } T = +1$  and that  $T$  maps the upper halfplane onto the lower halfplane (and the lower onto the upper) if and only if  $\text{sign } T = -1$ .

**4.1.8.** (i) Let  $z_0 \in \mathbb{D}$  and  $|\lambda| = 1$  and consider the l.f.t.  $T(z) = \lambda \frac{z - z_0}{1 - \bar{z}_0 z}$ . Prove that  $T(\mathbb{T}) = \mathbb{T}$  and  $T(z_0) = 0$ . Find  $T(\mathbb{D})$ .

(ii) Let  $z_0 \in \mathbb{D}$  and let  $T$  be a l.f.t. such that  $T(\mathbb{T}) = \mathbb{T}$  and  $T(z_0) = 0$ . Prove that there is  $\lambda$  with  $|\lambda| = 1$  so that  $T(z) = \lambda \frac{z - z_0}{1 - \bar{z}_0 z}$ .

(iii) Let  $a, b \in \mathbb{D}$  and let  $T$  be a l.f.t. such that  $T(\mathbb{T}) = \mathbb{T}$  and  $T(a) = b$ . Prove that there is  $\lambda$  with  $|\lambda| = 1$  so that  $\frac{T(z) - b}{1 - \bar{b}T(z)} = \lambda \frac{z - a}{1 - \bar{a}z}$ .

**4.1.9.** Consider  $\mathbb{H}_+ = \{z \mid \text{Im } z > 0\}$  and  $\mathbb{H}_- = \{z \mid \text{Im } z < 0\}$ .

(i) Let  $z_0 \in \mathbb{H}_+$  and  $|\lambda| = 1$  and consider the l.f.t.  $T(z) = \lambda \frac{z - z_0}{z - \bar{z}_0}$ . Prove that  $T(\mathbb{R} \cup \{\infty\}) = \mathbb{T}$  and  $T(z_0) = 0$ . Find  $T(\mathbb{H}_+)$ .

(ii) Let  $z_0 \in \mathbb{H}_+$  and let  $T$  be a l.f.t. such that  $T(\mathbb{R} \cup \{\infty\}) = \mathbb{T}$  and  $T(z_0) = 0$ . Prove that there is  $\lambda$  with  $|\lambda| = 1$  so that  $T(z) = \lambda \frac{z - z_0}{z - \bar{z}_0}$ .

**4.1.10.** Consider distinct  $z_1, z_2, z_3, z_4 \in \widehat{\mathbb{C}}$ . We define the **double ratio** of  $z_1, z_2, z_3, z_4$  (in this order) to be  $(z_1, z_2, z_3, z_4) = \frac{z_1 - z_3}{z_1 - z_4} \frac{z_2 - z_4}{z_2 - z_3}$ , if  $z_1, z_2, z_3, z_4 \in \mathbb{C}$ . If one of the  $z_1, z_2, z_3, z_4$  is  $\infty$ , say  $z_j = \infty$ , then we define  $(z_1, z_2, z_3, z_4)$  as the limit of the above formula when  $z_j \rightarrow \infty$ .

(i) Prove that  $(T(z_1), T(z_2), T(z_3), T(z_4)) = (z_1, z_2, z_3, z_4)$  for every l.f.t.  $T$  and every distinct  $z_1, z_2, z_3, z_4 \in \widehat{\mathbb{C}}$ .

(ii) Prove that the distinct  $z_1, z_2, z_3, z_4 \in \widehat{\mathbb{C}}$  belong to the same generalized circle if and only if  $(z_1, z_2, z_3, z_4) \in \mathbb{R} \setminus \{0\}$ .

(iii) If  $(z_1, z_2, z_3, z_4) = \lambda$ , find all values (depending on  $\lambda$ ) which result from this double ratio after all rearrangements of  $z_1, z_2, z_3, z_4$ .

**4.1.11.** Prove that the group of all l.f.t. is simple, i.e. that its only normal subgroups are itself and  $\{I\}$ , where  $I$  is the identity l.f.t.

## 4.2 The exponential function.

We define the **exponential function**  $\exp : \mathbb{C} \rightarrow \mathbb{C}$  by

$$\exp z = e^x (\cos y + i \sin y) \quad \text{for every } z = x + iy.$$

If  $z \in \mathbb{R}$ , i.e.  $z = x + i0$ , then  $\exp z = e^x (\cos 0 + i \sin 0) = e^x = e^z$ . This implies that we may use the symbol  $e^z$  instead of  $\exp z$  without the danger of contradiction, in the case that  $z$  is real, between the symbol  $e^z$  as we just defined it and the symbol  $e^z$  as we know it from infinitesimal calculus. Therefore, we define

$$e^z = \exp z = e^x (\cos y + i \sin y) \quad \text{for every } z = x + iy.$$

Since  $z = x + iy$  implies  $|e^z| = |e^x| |\cos y + i \sin y| = e^x$ , we have that

$$|e^z| = e^{\operatorname{Re} z}.$$

From  $e^z = e^x (\cos y + i \sin y)$  and  $|e^z| = e^x$  we get  $e^z = |e^z| (\cos y + i \sin y)$ . So  $y$  is one of the elements of  $\arg e^z$  and hence

$$\arg e^z = \{\operatorname{Im} z + k2\pi \mid k \in \mathbb{Z}\}.$$

We have the basic equality

$$e^{z_1} e^{z_2} = e^{z_1 + z_2}.$$

Indeed:  $e^{x_1} (\cos y_1 + i \sin y_1) e^{x_2} (\cos y_2 + i \sin y_2) = e^{x_1 + x_2} (\cos(y_1 + y_2) + i \sin(y_1 + y_2))$  from the addition formulas of  $\cos$  and  $\sin$ .

If  $z_2 - z_1 = k2\pi i$  for some  $k \in \mathbb{Z}$ , then  $e^{z_2} = e^{z_1} e^{k2\pi i} = e^{z_1} (\cos(k2\pi) + i \sin(k2\pi)) = e^{z_1}$ . Conversely, assume  $e^{z_2} = e^{z_1}$  and let  $z_2 - z_1 = x + iy$ . Then  $e^x (\cos y + i \sin y) = e^{z_2 - z_1} = \frac{e^{z_2}}{e^{z_1}} = 1$  and hence  $e^x = 1$ ,  $\cos y = 1$  and  $\sin y = 0$ . Therefore,  $x = 0$  and  $y = k2\pi$  for some  $k \in \mathbb{Z}$ . Thus,  $z_2 - z_1 = k2\pi i$  with  $k \in \mathbb{Z}$ . We proved that

$$e^{z_2} = e^{z_1} \quad \Leftrightarrow \quad z_2 - z_1 = k2\pi i \text{ for some } k \in \mathbb{Z}.$$

For all  $z = x + iy$  we have  $|e^z| = e^x > 0$  and hence

$$e^z \neq 0.$$

On the other hand, if we take any  $w \neq 0$  and if we use the notation  $\ln : (0, +\infty) \rightarrow \mathbb{R}$  for the well known logarithmic function from infinitesimal calculus, then the solutions of the equation  $e^z = w$  are described as follows:

$$e^z = w \quad \Leftrightarrow \quad z = \ln |w| + iy \text{ for some } y \in \arg w.$$

Indeed, if we write  $z = x + iy$ , then the equality  $w = e^z$  becomes  $w = e^x(\cos y + i \sin y)$  and it just means that its right side is one of the polar representations of  $w$ . Hence,  $w = e^z$  if and only if  $e^x = |w|$  and  $y$  is a value of  $\arg w$ . Now,  $e^x = |w|$  is equivalent to  $x = \ln |w|$ . Therefore, the equation  $e^z = w$  has these infinitely many solutions:  $z = \ln |w| + iy$  where  $y$  is any value of  $\arg w$ . All these solutions have the same real part,  $x = \ln |w|$ , and their imaginary parts are the elements of  $\arg w$ .

From what we said already, it is clear that the exponential function is onto  $\mathbb{C} \setminus \{0\}$  but not one-to-one in  $\mathbb{C}$ . In fact the exponential function is *infinity-to-one* since there are infinitely many values of  $z$  corresponding to the same value of  $w \neq 0$ .

Based on the equality  $e^{iy} = \cos y + i \sin y$ , we may write the polar representations of any  $z \neq 0$  in an equivalent form:

$$z = r(\cos \theta + i \sin \theta) \quad \Leftrightarrow \quad z = re^{i\theta}.$$

The second form is simpler and we shall use it extensively in the rest of the course. For instance, we may rewrite the examples 2.3.8 and 2.3.9 as follows.

**Example 4.2.1.** Using the parametric equation  $z = \gamma(t) = z_0 + re^{it}$ ,  $t \in [0, 2\pi]$ , for the circle  $C_{z_0}(r)$ , we have

$$\oint_{C_{z_0}(r)} f(z) dz = \oint_{\gamma} f(z) dz = \int_0^{2\pi} f(z_0 + re^{it}) ire^{it} dt.$$

**Example 4.2.2.** If  $n \in \mathbb{Z}$ , we have  $\int_0^{2\pi} e^{int} dt = 2\pi$ , if  $n = 0$ , and  $\int_0^{2\pi} e^{int} dt = 0$ , if  $n \neq 0$ . Therefore, if  $n \in \mathbb{Z}$ , we get

$$\oint_{C_{z_0}(r)} (z - z_0)^n dz = \int_0^{2\pi} r^n e^{int} ire^{it} dt = ir^{n+1} \int_0^{2\pi} e^{i(n+1)t} dt = \begin{cases} 2\pi i, & \text{if } n = -1 \\ 0, & \text{if } n \neq -1 \end{cases}$$

The real and imaginary parts of  $e^z$  are  $u(x, y) = e^x \cos y$  and  $v(x, y) = e^x \sin y$ . Therefore,  $u, v$  have partial derivatives  $\frac{\partial u}{\partial x}(x, y) = e^x \cos y$ ,  $\frac{\partial u}{\partial y} = -e^x \sin y$ ,  $\frac{\partial v}{\partial x} = e^x \sin y$  and  $\frac{\partial v}{\partial y} = e^x \cos y$ , which are continuous and satisfy the system of C-R equations in  $\mathbb{C}$  and hence  $e^z$  is holomorphic in  $\mathbb{C}$ . To calculate the derivative of  $e^z$  we write

$$\frac{\partial u}{\partial x}(x, y) + i \frac{\partial v}{\partial x}(x, y) = e^x \cos y + ie^x \sin y = e^x(\cos y + i \sin y)$$

and hence

$$\frac{de^z}{dz} = e^z.$$

We shall now examine the mapping properties of the function  $w = e^z$ . We write  $z = x + iy$  and  $w = u + iv$ .

If  $z = x + iy$  varies on the horizontal line  $h_y$  in the  $z$ -plane which intersects the  $y$ -axis at the fixed point  $iy$ , then  $w = e^z = e^x(\cos y + i \sin y)$  varies on the halfline  $r_y$  in the  $w$ -plane with vertex 0 (without 0) which forms angle  $y$  with the positive  $u$ -semiaxis. Also, if  $z$  varies on the horizontal line  $h_y$  from left to right, i.e. when  $x$  increases from  $-\infty$  to  $+\infty$ , then  $w = e^z$  varies on the halfline  $r_y$  from 0 to  $\infty$ . If  $y$  increases by  $\Delta y > 0$ , i.e. if the horizontal line  $h_y$  moves upward, then the corresponding halfline  $r_y$  rotates in the positive direction around 0 by the angle  $\Delta y$ . The two horizontal lines  $h_y$  and  $h_{y+2\pi}$  are mapped onto the same halfline  $r_y = r_{y+2\pi}$ .

If the point  $z = x + iy$  varies on the vertical line  $v_x$  in the  $z$ -plane which intersects the  $x$ -axis at the fixed point  $x$ , then  $w = e^z = e^x(\cos y + i \sin y)$  varies on the circle  $C_0(e^x)$ , call it  $c_x$ , in the  $w$ -plane. Also, if  $z$  moves upward on the vertical line  $v_x$ , i.e. if  $y$  increases from  $-\infty$  to  $+\infty$ , then  $w = e^z$  rotates on the circle  $c_x$  infinitely many times in the positive direction. If  $y$  increases over an interval of length  $2\pi$ , then  $w = e^z$  describes the whole circle  $c_x$  once in the positive direction. If  $x$  increases by  $\Delta x > 0$ , i.e. if the vertical line  $v_x$  moves to the right, then the circle  $c_x$  with radius  $e^x$  becomes the circle  $c_{x+\Delta x}$  with radius  $e^{x+\Delta x} = e^x e^{\Delta x}$ .

We may combine the above results. For instance, if we consider the open rectangle

$$\Pi = \{x + iy \mid x_1 < x < x_2, y_1 < y < y_2\}$$

in the  $z$ -plane with sides parallel to the two coordinate axes, then  $\Pi$  is the intersection of the open horizontal zone between the lines  $h_{y_1}$  and  $h_{y_2}$  and the open vertical zone between the lines  $v_{x_1}$  and  $v_{x_2}$ . If  $y_2 - y_1 < 2\pi$ , then  $\Pi$  is mapped onto the open “circular rectangle”

$$R = \{re^{i\theta} \mid e^{x_1} < r < e^{x_2}, y_1 < \theta < y_2\},$$

in the  $w$ -plane, which is the intersection of the angular region between the halflines  $r_{y_1}$  and  $r_{y_2}$  and the open ring between the circles  $c_{x_1}$  and  $c_{x_2}$ . If  $y_2 - y_1 = 2\pi$ , then the “circular rectangle”  $R$  is the open ring between the circles  $c_{x_1}$  and  $c_{x_2}$  without its linear segment which belongs to the halfline  $r_{y_1} = r_{y_2}$ . Of course, in this case, if  $\Pi$  includes at least one of its horizontal sides, then its image  $R$  is the whole open ring between the circles  $c_{x_1}$  and  $c_{x_2}$ .

Starting from  $e^{iy} = \cos y + i \sin y$  and  $e^{-iy} = \cos(-y) + i \sin(-y) = \cos y - i \sin y$ , we easily find that  $\cos y = \frac{1}{2}(e^{iy} + e^{-iy})$  and  $\sin y = \frac{1}{2i}(e^{iy} - e^{-iy})$  for every  $y \in \mathbb{R}$ . Now we extend the trigonometric functions **cosine** and **sine** from  $\mathbb{R}$  to  $\mathbb{C}$  by defining

$$\cos z = \frac{1}{2}(e^{iz} + e^{-iz}), \quad \sin z = \frac{1}{2i}(e^{iz} - e^{-iz}) \quad \text{for every } z \in \mathbb{C}.$$

It is clear from the holomorphy of the exponential function that  $\cos$  and  $\sin$  are holomorphic in  $\mathbb{C}$  and that

$$\frac{d \cos z}{dz} = -\sin z, \quad \frac{d \sin z}{dz} = \cos z.$$

It is also easy to show that  $\cos$  and  $\sin$  are  $2\pi$ -periodic.

Now we extend the **tangent** and the **cotangent** from  $\mathbb{R}$  to  $\mathbb{C}$  by defining

$$\tan z = \frac{\sin z}{\cos z}, \quad \cot z = \frac{\cos z}{\sin z} \quad \text{for every } z \in \mathbb{C}.$$

The solutions of  $\cos z = 0$  are  $z = \frac{\pi}{2} + k\pi$ ,  $k \in \mathbb{Z}$ , and the solutions of  $\sin z = 0$  are  $z = k\pi$ ,  $k \in \mathbb{Z}$ . Therefore,  $\tan$  is defined and holomorphic in the open set  $\mathbb{C} \setminus \{\frac{\pi}{2} + k\pi \mid k \in \mathbb{Z}\}$  and  $\cot$  is defined and holomorphic in the open set  $\mathbb{C} \setminus \{k\pi \mid k \in \mathbb{Z}\}$ . Both functions are  $\pi$ -periodic.

### Exercises.

**4.2.1.** Prove that  $\overline{e^z} = e^{\bar{z}}$  for all  $z$ .

**4.2.2.** Prove that  $|e^z - 1| \leq e^{|z|} - 1 \leq |z|e^{|z|}$ .

**4.2.3.** Let  $z \rightarrow \infty$  on any halfline. Depending on the halfline, study the existence of the  $\lim e^z$  in  $\widehat{\mathbb{C}}$ . Which characteristic of the halfline determines the existence and the value of the limit?

**4.2.4.** Find the images through the exponential function of:  $\{x + iy \mid a < x < b, \theta < y < \theta + \pi\}$ ,  $\{x + iy \mid a < x < b, \theta < y < \theta + 2\pi\}$ ,  $\{x + iy \mid x < b, \theta < y < \theta + \pi\}$ ,  $\{x + iy \mid x < b, \theta < y < \theta + 2\pi\}$ ,  $\{x + iy \mid a < x, \theta < y < \theta + \pi\}$ ,  $\{x + iy \mid a < x, \theta < y < \theta + 2\pi\}$ .

**4.2.5.** Every horizontal and every vertical line in the  $z$ -plane are perpendicular. Also, every halfline with vertex 0 and every circle with center 0 in the  $w$ -plane are perpendicular. How do these facts relate to the conformality of the function  $w = e^z$ ?

**4.2.6.** (i) Prove that

$$\cdot \sin^2 z + \cos^2 z = 1,$$

$$\cdot \sin(z + w) = \sin z \cos w + \cos z \sin w, \quad \cos(z + w) = \cos z \cos w - \sin z \sin w,$$

$$\cdot |\cos(x + iy)|^2 = \cos^2 x + \sinh^2 y, \quad |\sin(x + iy)|^2 = \sin^2 x + \sinh^2 y.$$

(ii) Study the function  $w = \sin z$  in the vertical zone  $\{x + iy \mid -\frac{\pi}{2} < x < \frac{\pi}{2}\}$  and the function  $w = \cos z$  in the vertical zone  $\{x + iy \mid 0 < x < \pi\}$ . Examine the images through these functions of the various horizontal linear segments (of length  $\pi$ ) and the various vertical lines inside these two vertical zones.

### 4.3 Branches of the logarithmic function.

In the last section we proved, for every  $w \neq 0$ , the equivalence

$$e^z = w \Leftrightarrow z = \ln |w| + iy \text{ for some } y \in \arg w. \quad (4.3)$$

For every  $w \neq 0$  we consider the set

$$\log w = \{\ln |w| + iy \mid y \in \arg w\}$$

and we call it **logarithm** of  $w$ . So the elements of  $\log w$  are the solutions of  $e^z = w$ , i.e.

$$e^z = w \Leftrightarrow z \in \log w.$$

If we take  $y = \text{Arg } w$ , then we get the particular element

$$\text{Log } w = \ln |w| + i \text{Arg } w$$

of  $\log w$  and this is called **principal logarithm** of  $w$ .

If  $r = |w|$  and if  $\theta$  is any of the values of the argument of  $w$ , i.e. if  $w = r(\cos \theta + i \sin \theta) = re^{i\theta}$  is any of the polar representations of  $w$ , then the values of  $\arg w$  are the numbers  $\theta + k2\pi$ ,  $k \in \mathbb{Z}$ . Hence the values of  $\log w$  are the numbers  $\ln r + i(\theta + k2\pi)$ ,  $k \in \mathbb{Z}$ .

**Example 4.3.1.** (i)  $\text{Log } 1 = 0$  and  $\log 1 = \{i2k\pi \mid k \in \mathbb{Z}\}$ .

(ii)  $\text{Log}(-1) = i\pi$  and  $\log(-1) = \{i(2k+1)\pi \mid k \in \mathbb{Z}\}$ .

(iii)  $\text{Log } i = i\frac{\pi}{2}$  and  $\log i = \{i(2k + \frac{1}{2})\pi \mid k \in \mathbb{Z}\}$ .

(iv)  $\text{Log}(-3i) = \ln 3 - i\frac{\pi}{2}$  and  $\log(-i) = \{\ln 3 + i(2k - \frac{1}{2})\pi \mid k \in \mathbb{Z}\}$ .

(v)  $\text{Log}(1+i) = \ln \sqrt{2} + i\frac{\pi}{4}$  and  $\log(1+i) = \{\ln \sqrt{2} + i(2k + \frac{1}{4})\pi \mid k \in \mathbb{Z}\}$ .

(vi)  $\text{Log}(1-i\sqrt{3}) = \ln 2 - i\frac{\pi}{3}$  and  $\log(1-i\sqrt{3}) = \{\ln 2 + i(2k - \frac{1}{3})\pi \mid k \in \mathbb{Z}\}$ .

For any fixed  $w \neq 0$  the set  $\log w$  has infinitely many elements, and any two of them differ by an integral multiple of  $i2\pi$ . All elements of  $\log w$  have the same real part  $x = \ln |w|$ , and hence they are on the same vertical line  $v_x$  with equation  $x = \ln |w|$ , and the vertical differences between them are the integral multiples of  $2\pi$ . Therefore, every vertical segment of the line  $v_x$ , which has length  $2\pi$  and includes only one of its endpoints, contains exactly one element of  $\log w$ . Moreover, every horizontal zone, which has vertical width  $2\pi$  and includes only one of its boundary lines (either the upper or the lower one), contains exactly one element of  $\log w$  for every  $w \neq 0$ . More precisely, if we consider any  $\theta_0$  and the horizontal zone

$$Z_{\theta_0} = \{x + iy \mid \theta_0 < y \leq \theta_0 + 2\pi\} \quad \text{or} \quad Z_{\theta_0} = \{x + iy \mid \theta_0 \leq y < \theta_0 + 2\pi\},$$

then  $Z_{\theta_0}$  contains exactly one element of  $\log w$ : the one with imaginary part  $y$  equal to the (unique)  $\theta \in \arg w$  satisfying  $\theta_0 < \theta \leq \theta_0 + 2\pi$  or  $\theta_0 \leq \theta < \theta_0 + 2\pi$ , respectively. For instance, if we consider the special zone determined by  $\theta_0 = -\pi$  which contains its upper boundary line, i.e.

$$Z_{-\pi} = \{x + iy \mid -\pi < y \leq \pi\},$$

then, for every  $w \neq 0$ , the unique element of  $\log w$  which is contained in this zone is the principal logarithm  $\text{Log } w$ .

**Proposition 4.3.** For all  $w_1, w_2 \neq 0$  we have

$$\log(w_1 w_2) = \log w_1 + \log w_2.$$

By this we mean that the sum of any element of  $\log w_1$  and any element of  $\log w_2$  is an element of  $\log(w_1 w_2)$  and, conversely, any element of  $\log(w_1 w_2)$  is the sum of an element of  $\log w_1$  and an element of  $\log w_2$ .

*Proof.* A corollary of proposition 1.1 and of the equality  $\ln |w_1 w_2| = \ln |w_1| + \ln |w_2|$ .  $\square$

It is already clear that the exponential function  $w = \exp z = e^z$  from  $\mathbb{C}$  onto  $\mathbb{C} \setminus \{0\}$  is not one-to-one. Therefore, there is no inverse of the exponential function. If we want to produce some kind of inverse of the exponential function, we may take any  $w$  in the range  $\mathbb{C} \setminus \{0\}$  of the function and select one value of  $z$  out of the infinitely many in  $\mathbb{C}$  which satisfy the  $e^z = w$ . There are many instances of this method at a more elementary level. Let us consider for instance the function  $y = x^2$  from  $(-\infty, +\infty)$  onto  $[0, +\infty)$ , which is not one-to-one in  $(-\infty, +\infty)$ . We take any  $y \in [0, +\infty)$  (the range of  $y = x^2$ ) and find one  $x$  such that  $x^2 = y$ . There are exactly two such  $x$ :  $x = \sqrt{y}$  and  $x = -\sqrt{y}$ . Therefore, one might say that we have only two choices for the inverse function: the choice  $x = \sqrt{y}$  for every  $y \in [0, +\infty)$  and the choice  $x = -\sqrt{y}$  for every  $y \in [0, +\infty)$ . *But this is not correct.* We may choose  $x = \sqrt{y}$  for some  $y \in [0, +\infty)$  and  $x = -\sqrt{y}$  for the remaining  $y \in [0, +\infty)$ . It is obvious that there are infinitely many such inverse functions, depending on the particular choice we make between  $x = \sqrt{y}$  and  $x = -\sqrt{y}$  for each value of  $y$ . Nevertheless, there is a criterion which reduces the number of our inverse functions to *exactly two*: the criterion of *continuity*! We observe that the last function, with the double formula, is not continuous. On the contrary, the function  $x = \sqrt{y}$  for every  $y \in [0, +\infty)$  and the function  $x = -\sqrt{y}$  for every  $y \in [0, +\infty)$  are both continuous. To prove that these are the only continuous inverse functions is a simple exercise in real analysis. Indeed, assume that there is some continuous inverse function  $x = f(y)$  of  $y = x^2$  defined in  $[0, +\infty)$  (the range of  $y = x^2$ ). I.e.  $f : [0, +\infty) \rightarrow \mathbb{R}$  is continuous in  $[0, +\infty)$  and  $f(y)^2 = y$  for every  $y \in [0, +\infty)$ . Let there be  $y_1, y_2 > 0$  with  $y_1 \neq y_2$  such that  $f(y_1) = \sqrt{y_1}$  and  $f(y_2) = -\sqrt{y_2}$ . Since  $f$  is continuous in the interval between  $y_1, y_2$  and its values at the endpoints are opposite, there is some  $y$  in this interval so that  $f(y) = 0$ . This is impossible, because  $y > 0$  and either  $f(y) = \sqrt{y} > 0$  or  $f(y) = -\sqrt{y} < 0$ . Therefore, there are no such  $y_1, y_2 > 0$  and hence we have exactly two cases: either  $f(y) = \sqrt{y}$  for every  $y \geq 0$  or  $f(y) = -\sqrt{y}$  for every  $y \geq 0$ . We may say that there are exactly two *continuous branches of the square root* in  $[0, +\infty)$ : the branch  $x = \sqrt{y}$  and the branch  $x = -\sqrt{y}$ .

Now let us go back to the determination of possible inverses of the exponential function.

Let  $A \subseteq \mathbb{C} \setminus \{0\}$ . We say that the function  $f$  is a **continuous branch of log** in  $A$  if  $f$  is continuous in  $A$  and for every  $w \in A$  we have that  $f(w)$  is an element of  $\log w$  or, equivalently,

$$e^{f(w)} = w \quad \text{for every } w \in A.$$

Proposition 4.4 gives many useful examples of continuous branches of the logarithm.

**Proposition 4.4.** *Let  $\theta_0 \in \mathbb{R}$ . We consider the set*

$$A_{\theta_0} = \{re^{i\theta} \mid 0 < r < +\infty, \theta_0 < \theta < \theta_0 + 2\pi\}$$

*in the  $w$ -plane (i.e.  $\mathbb{C}$  without the halfline with vertex 0 which forms angle  $\theta_0$  with the positive  $u$ -semiaxis, where  $w = u + iv$ ) and the open horizontal zone*

$$Z_{\theta_0} = \{x + iy \mid -\infty < x < +\infty, \theta_0 < y < \theta_0 + 2\pi\}$$

*in the  $z$ -plane. We define the function  $f : A_{\theta_0} \rightarrow Z_{\theta_0}$  as follows: for every  $w \in A_{\theta_0}$  we take  $f(w)$  to be the unique element of  $\log w$  in the zone  $Z_{\theta_0}$ . Then  $f$  is continuous in  $A_{\theta_0}$  and so it is a continuous branch of  $\log$  in  $A_{\theta_0}$ .*

*Proof.* Assume that  $f$  is not continuous at some  $w$  in  $A_{\theta_0}$ . Then there is a sequence  $(w_n)$  in  $A_{\theta_0}$  so that  $w_n \rightarrow w$  and  $f(w_n) \not\rightarrow f(w)$ . This implies that there is  $\delta > 0$  so that  $|f(w_n) - f(w)| \geq \delta > 0$  for infinitely many  $n$ . These infinitely many  $n$  define a subsequence of  $(w_n)$ . Now we ignore the rest of the sequence  $(w_n)$  and concentrate on the specific subsequence. For simplicity we rename the subsequence and call it  $(w_n)$  again. Therefore, we have a sequence  $(w_n)$  in  $A_{\theta_0}$  such that

$$w_n \rightarrow w \quad \text{and} \quad |f(w_n) - f(w)| \geq \delta > 0 \quad \text{for every } n. \quad (4.4)$$

We set  $z = f(w) \in Z_{\theta_0}$  and  $z_n = f(w_n) \in Z_{\theta_0}$  for every  $n$ . Then  $e^z = w$  and  $e^{z_n} = w_n$  for every  $n$  and (4.4) becomes

$$e^{z_n} \rightarrow e^z \quad \text{and} \quad |z_n - z| \geq \delta > 0 \quad \text{for every } n. \quad (4.5)$$

The real parts of the  $z_n$  are equal to  $\ln |w_n|$  and, since  $\ln |w_n| \rightarrow \ln |w|$ , the real parts of the  $z_n$  are bounded. Moreover, since  $z_n \in Z_{\theta_0}$ , the imaginary parts of the  $z_n$  are also bounded. Therefore, the sequence  $(z_n)$  is bounded and the Bolzano-Weierstrass theorem implies that there is a subsequence  $(z_{n_k})$  so that  $z_{n_k} \rightarrow z'$  for some  $z'$ . Since all  $z_{n_k}$  belong to  $Z_{\theta_0}$ , we see that  $z'$  belongs to the closed zone  $\bar{Z}_{\theta_0} = \{x + iy \mid -\infty < x < +\infty, \theta_0 \leq y \leq \theta_0 + 2\pi\}$ . Taking the limit in (4.5), we get that  $e^{z'} = e^z$  and  $|z' - z| \geq \delta$ . Therefore,  $z'$  and  $z$  differ by a non-zero integral multiple of  $i2\pi$ . But this is impossible, because  $z$  belongs to the open zone  $Z_{\theta_0}$  and  $z'$  belongs to the closed zone  $\bar{Z}_{\theta_0}$ . Thus  $f$  is continuous at every  $w$  in  $A_{\theta_0}$ .  $\square$

Our study of the mapping properties of the exponential function in the previous section gives the following information about the mapping properties of the continuous branch  $f : A_{\theta_0} \rightarrow Z_{\theta_0}$  of  $\log$ , which is defined in proposition 4.4:  $f$  maps the halflines in  $A_{\theta_0}$  with vertex 0 (without 0) onto the horizontal lines in  $Z_{\theta_0}$  and the circles with center 0 (without their point on the halfline which is excluded from  $A_{\theta_0}$ ) onto the vertical segments of  $Z_{\theta_0}$ .

Choosing any real  $\theta_0$ , we have defined a continuous branch of  $\log$  in the subset  $A_{\theta_0}$  of the  $w$ -plane, whose range is the zone  $Z_{\theta_0}$  of the  $z$ -plane. If, instead of  $\theta_0$ , we consider  $\theta_0 + k2\pi$  with any  $k \in \mathbb{Z}$ , then the domain  $A = A_{\theta_0+k2\pi}$  remains the same but the range, i.e. the zone  $Z_{\theta_0+k2\pi}$ , moves vertically by  $k2\pi$ . The various zones  $Z_{\theta_0+k2\pi}$  are successive and cover the whole  $z$ -plane (except for their boundary lines with equations  $y = \theta_0 + k2\pi$ ). We summarize:

*If we exclude from the  $w$ -plane a halfline with vertex 0, then in the remaining open set  $A$  there are infinitely many continuous branches of  $\log$  defined. Each of them maps  $A$  onto some open horizontal zone of the  $z$ -plane of width  $2\pi$ . These various open zones, which correspond to the various continuous branches of  $\log$  (in the same set  $A$ ), are mutually disjoint, successive and cover the  $z$ -plane (except for their boundary lines). Of course, if we change the original halfline which determines the set  $A$ , then the corresponding zones and the corresponding continuous branches of  $\log$  also change.*

**Example 4.3.2.** One particular example of a continuous branch of  $\log$  is defined when we choose  $\theta_0 = -\pi$ . Then the set  $A_{-\pi} = \{re^{i\theta} \mid 0 < r < +\infty, -\pi < \theta < \pi\}$  is the  $w$ -plane without the negative  $u$ -semiaxis (where  $w = u + iw$ ) and the range of the branch is the zone  $Z_{-\pi} = \{x + iy \mid -\infty < x < +\infty, -\pi < y < \pi\}$ . It is obvious that this branch is the function which maps every  $w \in A_{-\pi}$  onto the principal value  $z = \text{Log } w$  of  $\log w$ . I.e. we get the so-called **principal branch of  $\log$**

$$\text{Log} : A_{-\pi} \rightarrow Z_{-\pi}.$$

We must keep in mind that in the same set  $A_{-\pi}$  of the  $w$ -plane, besides the principal branch, there are infinitely many other continuous branches of  $\log$  defined. Each of them maps  $A_{-\pi}$  in a corresponding zone  $Z_{-\pi+k2\pi}$ , with  $k \in \mathbb{Z}$ , which is  $Z_{-\pi}$  moved vertically by  $k2\pi$ . This branch results from the principal branch  $\text{Log}$  by adding the constant  $ik2\pi$  and its formula is  $\text{Log} + ik2\pi$ .

Now, we introduce a slight generalization of the notion of the branch of  $\log$ , i.e. we define the notion of the branch of  $\log g$ , where  $g$  is a more general function than the identity  $g(w) = w$ .

Let  $A \subseteq \mathbb{C}$  and  $g : A \rightarrow \mathbb{C} \setminus \{0\}$  be continuous in  $A$ . We say that the function  $f$  is a **continuous branch of  $\log g$**  in  $A$  if  $f$  is continuous in  $A$  and for every  $w \in A$  we have that  $f(w)$  is an element of  $\log g(w)$  or, equivalently,

$$e^{f(w)} = g(w) \quad \text{for every } w \in A.$$



**Example 4.3.3.** Let  $g : A \rightarrow \mathbb{C} \setminus \{0\}$  be continuous in  $A \subseteq \mathbb{C}$ . If there is a continuous branch  $h$  of  $\log$  in  $g(A)$ , then  $f = h \circ g$  is a continuous branch of  $\log g$  in  $A$ .

Indeed,  $f = h \circ g$  is continuous in  $A$  and, since  $e^{h(z)} = z$  for every  $z \in g(A)$ , we also have  $e^{f(w)} = e^{h(g(w))} = g(w)$  for every  $w \in A$ .

This is a standard way to produce continuous branches of  $\log g$  when we know continuous branches of  $\log$  in the range of  $g$ .

For instance, if  $g(w) = w - w_0$  and  $A = \mathbb{C} \setminus l$ , where  $l$  is a halfline with vertex  $w_0$ , then  $g(A) = \mathbb{C} \setminus l'$ , where  $l'$  is the halfline with vertex 0 which is parallel to  $l$ . We know that there are infinitely many branches of  $\log$  defined in  $g(A)$  and hence there are infinitely many branches of  $\log(w - w_0)$  defined in  $A$ .

**Proposition 4.5.** Let  $g : A \rightarrow \mathbb{C} \setminus \{0\}$  be continuous in  $A \subseteq \mathbb{C}$  and let  $f$  be any continuous branch of  $\log g$  in  $A$ . If  $w_0$  is an interior point of  $A$  and  $g$  is differentiable at  $w_0$ , then  $f$  is differentiable at  $w_0$  and  $f'(w_0) = \frac{g'(w_0)}{g(w_0)}$ . Hence, if  $g$  is holomorphic in the interior of  $A$ , then  $f$  is also holomorphic in the interior of  $A$ .

*Proof.* We set  $z_0 = f(w_0)$  and  $z = f(w)$  for every  $w \in A$ . Then  $e^{z_0} = g(w_0)$  and  $e^z = g(w)$ . Since  $f$  is continuous,  $w \rightarrow w_0$  implies  $z \rightarrow z_0$ . Therefore, using the derivative of the exponential function at  $z_0$ , we see that

$$\frac{f(w) - f(w_0)}{w - w_0} = \frac{z - z_0}{e^z - e^{z_0}} \frac{g(w) - g(w_0)}{w - w_0} \rightarrow \frac{g'(w_0)}{e^{z_0}} = \frac{g'(w_0)}{g(w_0)} \quad \text{when } w \rightarrow w_0.$$

Thus  $f$  is differentiable at  $w_0$  and  $f'(w_0) = \frac{g'(w_0)}{g(w_0)}$ .  $\square$

Therefore, if  $g : A \rightarrow \mathbb{C} \setminus \{0\}$  is holomorphic in the open set  $A$ , every continuous branch of  $\log g$  can be called **holomorphic branch** of  $\log g$  in  $A$ .

**Example 4.3.4.** We have defined infinitely many continuous branches of  $\log$  in the open set which results when we exclude any halfline with vertex 0 from the  $w$ -plane. All these branches are *holomorphic branches* of  $\log$ . In particular the principal branch  $\text{Log} : A_{-\pi} \rightarrow Z_{-\pi}$  is holomorphic in  $A_{-\pi}$ .

**Proposition 4.6.** Let  $g : A \rightarrow \mathbb{C} \setminus \{0\}$  be continuous in  $A \subseteq \mathbb{C}$ .

(i) If  $f_1$  is a continuous branch of  $\log g$  in  $A$  and  $f_2 - f_1 = ik2\pi$  in  $A$ , where  $k$  is a fixed integer, then  $f_2$  is also a continuous branch of  $\log g$  in  $A$ .

(ii) If, moreover,  $A$  is connected and  $f_1, f_2$  are continuous branches of  $\log g$  in  $A$ , then  $f_2 - f_1 = ik2\pi$  in  $A$ , where  $k$  is a fixed integer. In particular, if  $f_1(w_0) = f_2(w_0)$  for some  $w_0 \in A$ , then  $f_1 = f_2$  in  $A$ .

*Proof.* (i) The continuity of  $f_1$  in  $A$  implies the continuity of  $f_2$  in  $A$ . We also have  $e^{f_1(w)} = g(w)$  for every  $w \in A$  and hence  $e^{f_2(w)} = e^{f_1(w) + ik2\pi} = e^{f_1(w)} e^{ik2\pi} = g(w)1 = g(w)$  for every  $w \in A$ . Therefore,  $f_2$  is a continuous branch of  $\log g$  in  $A$ .

(ii) We consider the function  $k = \frac{1}{i2\pi} (f_2 - f_1)$ . Since for every  $w \in A$  both  $f_2(w)$  and  $f_1(w)$  are elements of  $\log g(w)$ , we have that  $k(w)$  is an integer. Also, since both  $f_1, f_2$  are continuous in  $A$ ,  $k$  is continuous in  $A$ . Now,  $k$  is a continuous real function in the connected set  $A$ , and hence it has the intermediate value property. But since its only values are integers, it is constant in  $A$ . So there is a fixed integer  $k$  so that  $\frac{1}{i2\pi} (f_2 - f_1) = k$  or, equivalently,  $f_2 - f_1 = ik2\pi$  in  $A$ .

If  $f_2(w_0) = f_1(w_0)$  for some  $w_0 \in A$ , then the integer  $k$  is 0 and we get that  $f_2 = f_1$  in  $A$ .  $\square$

Thus, if we know one continuous branch of  $\log g$  in the connected set  $A$ , then we find every other possible continuous branch of  $\log g$  in  $A$  by adding to the known branch an arbitrary constant of the form  $ik2\pi$  with  $k \in \mathbb{Z}$ .

**Example 4.3.5.** Let  $A = A_{-\pi}$  be the  $w$ -plane without the negative  $u$ -semiaxis (where  $w = u + iv$ ). We want to find a continuous branch of  $\log$  in  $A$  having value  $z = 0$  when  $w = 1$ .

We already know that the principal branch  $\text{Log}$  of the logarithm has value  $z = \text{Log } 1 = 0$  at  $w = 1$ . Since  $A$  is connected, there is no other such continuous branch of  $\log$  in  $A$ .

Now, in the same set  $A = A_{-\pi}$  we want to find a continuous branch of  $\log$  taking the value  $z = i4\pi$  at  $w = 1$ .

Since  $A$  is connected the branch we are looking for has the form  $\text{Log} + ik2\pi$  for some fixed integer  $k$ . We try  $w = 1$  in this equality and get  $k = 2$ .

**Example 4.3.6.** Let  $A = A_0 = \{re^{i\theta} \mid 0 < r < +\infty, 0 < \theta < 2\pi\}$  be the  $w$ -plane without the positive  $u$ -semiaxis (where  $w = u + iv$ ). We want to find a continuous branch of  $\log$  in  $A$  taking the value  $z = i(\frac{\pi}{2} + 4\pi)$  at  $w = i$ .

We consider the horizontal zones in the  $z$ -plane which correspond to the set  $A$ : to each  $k \in \mathbb{Z}$  corresponds the zone  $Z_{0+k2\pi} = \{x + iy \mid -\infty < x < +\infty, k2\pi < y < 2\pi + k2\pi\}$ . Now we choose the particular zone which contains the value  $z = i(\frac{\pi}{2} + 4\pi)$ . This zone corresponds to  $k = 2$  and it is  $Z_{4\pi} = \{x + iy \mid -\infty < x < +\infty, 4\pi < y < 6\pi\}$ . Then we consider the continuous branch  $f$  of  $\log$  which maps  $A$  onto  $Z_{4\pi}$ :

$$f(w) = \ln r + i\theta \quad \text{for } w = re^{i\theta} \text{ and } r = |w| > 0, 4\pi < \theta < 6\pi,$$

where  $\theta$  is the unique value of  $\arg w$  which is contained in the interval  $(4\pi, 6\pi)$ .

Since  $A$  is connected, there is no other such continuous branch of  $\log$  in  $A$ .

### Exercises.

**4.3.1.** Let  $z \neq 0$ . Prove that the only element of  $\exp(\log z)$  is  $z$  and that the elements of  $\log(\exp z)$  are  $z + k2\pi i$ ,  $k \in \mathbb{Z}$ .

**4.3.2.** If  $A$  is any of the sets  $\{w \mid r_1 \leq |w| \leq r_2\} \setminus [-r_2, -r_1]$ ,  $\{w \mid 0 < |w| \leq r_2\} \setminus [-r_2, 0]$ ,  $\{w \mid r_1 \leq |w| < +\infty\} \setminus (-\infty, -r_1]$ , find  $\text{Log}(A)$ .

**4.3.3.** Work on the following in both cases:  $\theta_0 = -\pi$  and  $\theta_0 = 0$ .

Consider  $A_{\theta_0}$ , i.e. the  $w$ -plane without the halfline with vertex 0 which forms angle  $\theta_0$  with the positive  $u$ -semiaxis. Consider also  $\theta_1, \theta_2$  with  $\theta_0 < \theta_1 < \theta_2 < \theta_0 + 2\pi$  as well as  $r_1, r_2$  with  $0 < r_1 < r_2 < +\infty$ . Draw the set  $P = \{w = re^{i\theta} \mid r_1 < r < r_2, \theta_1 < \theta < \theta_2\}$  and its images through the various continuous branches of  $\log$  in  $A_{\theta_0}$ .

**4.3.4.** Let  $P = \{re^{i\theta} \mid 1 < r < 2, -\frac{3\pi}{4} < \theta < \frac{3\pi}{4}\}$ ,  $Q = \{w = re^{i\theta} \mid 1 < r < 2, \frac{\pi}{4} < \theta < \frac{7\pi}{4}\}$ . We know that there is a continuous branch  $f$  of  $\log$  in  $P$  and a continuous branch  $g$  of  $\log$  in  $Q$ . Is it possible for  $f$  and  $g$  to coincide in  $P \cap Q$ ?

**4.3.5.** Look back at exercise 1.2.1 and find all the possible values of  $\text{Log}(z_1 z_2) - \text{Log } z_1 - \text{Log } z_2$ .

**4.3.6.** Prove that there is no continuous branch of  $\log$  defined in any circle  $C_0(r)$  and hence in any set  $A$  which contains such a circle.

**4.3.7.** Define  $w^a = e^{a \text{Log } w}$  for every  $w \in D_1(1)$ , and prove that  $\lim_{x \rightarrow +\infty} (1 + \frac{z}{x})^x = e^z$  for every  $z$ .

**4.3.8.** Let  $A \subseteq \mathbb{C} \setminus \{0\}$ . If  $A$  is connected and if  $f_1, f_2$  are two different continuous branches of  $\log$  in  $A$ , prove that  $f_1(A) \cap f_2(A) = \emptyset$ . (Observe how this result is confirmed by the special case of  $A$  being  $\mathbb{C}$  without a halfline with vertex 0 in which case the various continuous branches of  $\log$  in  $A$  map  $A$  onto disjoint horizontal zones.)

**4.3.9.** Let  $a < b$ . Discuss the geometric meaning of the number  $\text{Log} \frac{z-b}{z-a}$  for every  $z$  with  $\text{Im } z > 0$ .

## 4.4 Powers and branches of roots.

If  $n \in \mathbb{N}$ ,  $n \geq 2$ , the function

$$w = z^n$$

is holomorphic in the  $z$ -plane  $\mathbb{C}$  and we shall examine some mapping properties of this function.

We work with polar representations:

$$z = re^{i\theta}, \quad w = r^n e^{in\theta}.$$

If  $\theta \in \mathbb{R}$  is constant and  $r$  varies in  $(0, +\infty)$ , i.e. if  $z$  moves on the halfline  $r_\theta$  in the  $z$ -plane with vertex 0 (without 0) which forms angle  $\theta$  with the positive  $x$ -semiaxis, then  $w = z^n$  moves on the halfline  $r_\phi$  in the  $w$ -plane with vertex 0 (without 0) which forms angle  $\phi = n\theta$  with the positive  $u$ -semiaxis. Also, if  $z$  moves on the halfline  $r_\theta$  from 0 to  $\infty$ , then  $w = z^n$  moves on the halfline  $r_\phi$  from 0 to  $\infty$ . If  $\theta$  increases by  $\Delta\theta > 0$ , i.e. if the halfline  $r_\theta$  turns in the positive direction by an angle  $\Delta\theta$ , then the corresponding halfline  $r_\phi$  turns in the positive direction by an angle  $\Delta\phi = n\Delta\theta$ . The two halflines  $r_\theta$  and  $r_{\theta+\frac{2\pi}{n}}$  are mapped onto the same halfline  $r_\phi = r_{\phi+2\pi}$ .

If  $r \in (0, +\infty)$  is constant and  $\theta$  varies in  $\mathbb{R}$ , i.e. if the point  $z$  moves on the circle  $C_0(r)$  in the  $z$ -plane, then  $w = z^n$  moves on the circle  $C_0(r^n)$  in the  $w$ -plane. Also, if  $z$  rotates once on  $C_0(r)$  in the positive direction, i.e. if  $\theta$  increases in an interval of length  $2\pi$ , then  $w = z^n$  rotates  $n$  times on  $C_0(r^n)$  in the positive direction. If  $\theta$  increases in an interval of length  $\frac{2\pi}{n}$ , then  $w = z^n$  rotates once on  $C_0(r^n)$  in the positive direction. If  $r$  increases, i.e. if the circle  $C_0(r)$  expands, then the corresponding circle  $C_0(r^n)$  also expands.

In the proof of the following proposition as well as in the whole course, we shall use the symbol  $\sqrt[n]{x}$  only to denote the unique nonnegative  $n$ -th root of a nonnegative real number  $x$ .

If  $n \in \mathbb{N}$ ,  $n \geq 2$  and if we take any  $w = Re^{i\Theta} \neq 0$ , then the equation  $z^n = w$  has  $n$  solutions which are described as follows:

$$z^n = w = Re^{i\Theta} \Leftrightarrow z = \sqrt[n]{R} e^{i(\frac{\Theta}{n} + k\frac{2\pi}{n})} \text{ for some } k = 0, 1, \dots, n-1. \quad (4.6)$$

Indeed, if we write  $z = re^{i\theta}$ , then the equality  $z^n = w$  becomes  $r^n e^{in\theta} = Re^{i\Theta}$  and this is equivalent to  $r^n = R$  and  $n\theta = \Theta + k2\pi$  for some  $k \in \mathbb{Z}$ . Solving for  $r$  and  $\theta$ , we find the solutions  $z = \sqrt[n]{R} e^{i(\frac{\Theta}{n} + k\frac{2\pi}{n})}$ ,  $k \in \mathbb{Z}$ . It is trivial to see that two of these solutions are the same if and only if the corresponding values of  $k$  differ by a multiple of  $n$  and hence there are  $n$  distinct solutions corresponding to the values  $0, 1, \dots, n-1$  of  $k$ . We easily see that the solutions of  $z^n = w$  are the vertices of a regular  $n$ -gon inscribed in the circle  $C_0(\sqrt[n]{R})$ .

The set of the solutions of  $z^n = w$ , which appear in the right side of (4.6), is called  **$n$ -th root** of  $w$  and it is denoted  $w^{\frac{1}{n}}$  or  $w^{1/n}$ , i.e.

$$w^{\frac{1}{n}} = \left\{ \sqrt[n]{R} e^{i(\frac{\Theta}{n} + k\frac{2\pi}{n})} \mid k = 0, 1, \dots, n-1 \right\} \quad \text{when } w = Re^{i\Theta}.$$

Thus, we have the equivalence

$$z^n = w \Leftrightarrow z \in w^{\frac{1}{n}}.$$

Of course, if  $w = 0$ , then the equation  $z^n = w$  has the unique solution  $z = 0$  and then we define  $0^{\frac{1}{n}} = \{0\}$ .

To get (4.6) we considered a polar representation  $w = Re^{i\Theta}$ . This means that the values of  $\arg w$  are the numbers  $\theta = \Theta + k2\pi$ ,  $k \in \mathbb{Z}$ , and we have that

$$w^{\frac{1}{n}} = \left\{ \sqrt[n]{|w|} e^{i\frac{\theta}{n}} \mid \theta \in \arg w \right\} = \left\{ e^{\frac{\zeta}{n}} \mid \zeta \in \log w \right\}.$$

**Example 4.4.1.** The  $n$ -th root of 1 is called  **$n$ -th root of unity**.

Since  $1 = 1e^{i0}$ , the elements of the  $n$ -th root of unity are the numbers  $e^{ik\frac{2\pi}{n}}$ ,  $k = 0, 1, \dots, n-1$ .

Obviously, one of them is 1 and, if we denote  $e^{i\frac{2\pi}{n}}$  by the symbol  $\omega_n$ , we find that the elements of the  $n$ -th root of unity are the numbers

$$1, \omega_n, \omega_n^2, \dots, \omega_n^{n-1}.$$

This  $\omega_n$  is called **principal  $n$ -th root of unity**.

We saw that, if  $w \neq 0$ , then  $w^{\frac{1}{n}}$  has exactly  $n$  elements which are on the vertices of a regular  $n$ -gon inscribed in the circle  $C_0(\sqrt[n]{|w|})$  of the  $z$ -plane. Therefore, every arc of this circle with central angle  $\frac{2\pi}{n}$ , which includes only one of its endpoints, contains exactly one of the elements of  $w^{\frac{1}{n}}$ . Thus, every angular set in the  $z$ -plane with vertex 0 and angle  $\frac{2\pi}{n}$ , which includes only one of its boundary halflines, contains, for every  $w \neq 0$ , exactly one element of  $w^{\frac{1}{n}}$ . In particular, if we consider any  $\theta_0$  and the angular set

$$A_{\theta_0} = \{re^{i\theta} \mid r > 0, \theta_0 < \theta \leq \theta_0 + \frac{2\pi}{n}\} \quad \text{or} \quad A_{\theta_0} = \{re^{i\theta} \mid r > 0, \theta_0 \leq \theta < \theta_0 + \frac{2\pi}{n}\},$$

then  $A_{\theta_0}$  contains exactly one element of  $w^{\frac{1}{n}}$ .

Clearly, the function  $w = z^n$  from  $\mathbb{C} \setminus \{0\}$  onto  $\mathbb{C} \setminus \{0\}$  is  $n$ -to-one and has no inverse. So we shall define *branches* of an inverse of  $w = z^n$ .

Let  $A \subseteq \mathbb{C} \setminus \{0\}$ . We say that the function  $f$  is a **continuous branch of  $w^{\frac{1}{n}}$**  in  $A$  if  $f$  is continuous in  $A$  and for every  $w \in A$  we have that  $f(w)$  is an element of  $w^{\frac{1}{n}}$  or, equivalently,

$$f(w)^n = w \quad \text{for every } w \in A.$$

Proposition 4.7 gives many examples of continuous branches of  $w^{\frac{1}{n}}$ .

**Proposition 4.7.** *Let  $\phi_0 \in \mathbb{R}$ . We consider the set*

$$A_{\phi_0} = \{se^{i\phi} \mid s > 0, \phi_0 < \phi < \phi_0 + 2\pi\}$$

*in the  $w$ -plane (i.e.  $\mathbb{C}$  without the halfline with vertex 0 which forms angle  $\phi_0$  with the positive  $u$ -semiaxis, where  $w = u + iv$ ) and the angular region*

$$B_{\phi_0/n} = \{re^{i\theta} \mid r > 0, \frac{\phi_0}{n} < \theta < \frac{\phi_0}{n} + \frac{2\pi}{n}\}$$

*in the  $z$ -plane. We define the function  $f : A_{\phi_0} \rightarrow B_{\phi_0/n}$  as follows: for every  $w \in A_{\phi_0}$  we take  $f(w)$  to be the unique element of  $w^{\frac{1}{n}}$  in the angular region  $B_{\phi_0/n}$ . Then  $f$  is continuous in  $A_{\phi_0}$  and so it is a continuous branch of  $w^{\frac{1}{n}}$  in  $A_{\phi_0}$ .*

*Proof.* Assume that  $f$  is not continuous at some  $w$  in  $A_{\phi_0}$ . Then there is a sequence  $(w_k)$  in  $A_{\phi_0}$  so that  $w_k \rightarrow w$  and  $f(w_k) \not\rightarrow f(w)$ . Then there is  $\delta > 0$  so that  $|f(w_k) - f(w)| \geq \delta > 0$  for infinitely many  $k$ . These infinitely many  $k$  define a subsequence of  $(w_k)$ . Now we ignore the rest of the sequence  $(w_k)$  and concentrate on the specific subsequence. For simplicity we rename the subsequence and call it  $(w_k)$  again. Therefore, we have a sequence  $(w_k)$  in  $A_{\phi_0}$  such that

$$w_k \rightarrow w \quad \text{and} \quad |f(w_k) - f(w)| \geq \delta > 0 \quad \text{for every } k. \quad (4.7)$$

We set  $z = f(w) \in B_{\phi_0/n}$  and  $z_k = f(w_k) \in B_{\phi_0/n}$  for every  $k$ . Then  $z^n = w$  and  $z_k^n = w_k$  for every  $k$  and (4.7) becomes

$$z_k^n \rightarrow z^n \quad \text{and} \quad |z_k - z| \geq \delta > 0 \quad \text{for every } k. \quad (4.8)$$

Since  $|z_k|^n \rightarrow |z|^n$  and hence  $|z_k| \rightarrow |z|$ , we get that the sequence  $(z_k)$  is bounded and the Bolzano-Weierstrass theorem implies that there is a subsequence  $(z_{k_m})$  so that  $z_{k_m} \rightarrow z'$  for some  $z'$ . Since all  $z_{k_m}$  belong to  $B_{\phi_0/n}$ , we have that  $z'$  belongs to the closed angular region  $\overline{B_{\phi_0/n}} = \{z = re^{i\theta} \mid r \geq 0, \frac{\phi_0}{n} \leq \theta \leq \frac{\phi_0}{n} + \frac{2\pi}{n}\}$ . Taking the limit in (4.8), we get  $z'^n = z^n$  and  $|z' - z| \geq \delta$ . This is impossible, because  $z$  belongs to  $B_{\phi_0/n}$  and  $z'$  belongs to  $\overline{B_{\phi_0/n}}$ .

Thus  $f$  is continuous at every  $w$  in  $A_{\phi_0}$ . □

From the mapping properties of the function  $w = z^n$  we get the following for the mapping properties of the continuous branch  $f : A_{\phi_0} \rightarrow B_{\phi_0/n}$  of  $w^{\frac{1}{n}}$ , which is defined in proposition 4.7. The function  $f$  maps the halflines in  $A_{\phi_0}$  with vertex 0 (without 0) onto the halflines in  $B_{\phi_0/n}$  with vertex 0 (without 0) and the circular arcs in  $A_{\phi_0}$  with center 0 onto the circular arcs in  $B_{\phi_0/n}$  with center 0.

Choosing any real  $\phi_0$ , we have defined a continuous branch of  $w^{\frac{1}{n}}$  in the subset  $A_{\phi_0}$  of the  $w$ -plane, whose range is the angular region  $B_{\phi_0/n}$  of the  $z$ -plane. If, instead of  $\phi_0$ , we consider  $\phi_0 + k2\pi$  with any  $k = 0, 1, \dots, n-1$ , then the set  $A = A_{\phi_0+k2\pi}$  remains the same but the range, i.e. the angular region  $B_{(\phi_0+k2\pi)/n}$ , rotates by an angle  $k\frac{2\pi}{n}$ . The  $n$  angular regions  $B_{(\phi_0+k2\pi)/n}$  with  $k = 0, 1, \dots, n-1$  are successive and cover the  $z$ -plane (except for their  $n$  boundary halflines with vertex 0). We summarize:

*If we exclude from the  $w$ -plane any halfline with vertex 0, then in the remaining open set  $A$  there are  $n$  continuous branches of  $w^{\frac{1}{n}}$  defined. Each of them maps  $A$  onto some open angular region of the  $z$ -plane with vertex 0 and angle  $\frac{2\pi}{n}$ . These various angular regions, which correspond to the various continuous branches of  $w^{\frac{1}{n}}$  (in the same set  $A$ ), are mutually disjoint, successive and cover the  $z$ -plane (except for their boundary halflines). Of course, if we change the original halfline which determines the set  $A$ , then the corresponding angular regions and the corresponding branches of  $w^{\frac{1}{n}}$  also change.*

**Example 4.4.2.** We get a concrete example of a continuous branch of  $w^{\frac{1}{n}}$  when we take  $\phi_0 = -\pi$ . Then the set  $A_{-\pi} = \{se^{i\phi} \mid s > 0, -\pi < \phi < \pi\}$  is the  $w$ -plane without the negative  $u$ -semiaxis (where  $w = u + iv$ ) and the range of the continuous branch of  $w^{\frac{1}{n}}$  is the angular region  $B_{-\pi/n} = \{re^{i\theta} \mid r > 0, -\frac{\pi}{n} < \theta < \frac{\pi}{n}\}$ . This branch is given by

$$z = \sqrt[n]{s} e^{i\frac{\phi}{n}} \quad \text{for } w = se^{i\phi} \text{ with } -\pi < \phi < \pi.$$

Clearly,

$$z = \sqrt[n]{|w|} e^{i\frac{\text{Arg } w}{n}} = e^{\frac{\text{Log } w}{n}}.$$

On the same set  $A_{-\pi}$  of the  $w$ -plane, besides the above continuous branch of  $w^{\frac{1}{n}}$ , we may define  $n$  continuous branches of  $w^{\frac{1}{n}}$ . Each of them maps  $A_{-\pi}$  onto a corresponding angular region  $B_{(-\pi+k2\pi)/n}$  with  $k = 0, 1, \dots, n-1$ , which results by rotating  $B_{-\pi/n}$  in the positive direction by the angle  $k\frac{2\pi}{n}$ . This branch results from the original branch by multiplication by the constant  $e^{ik\frac{2\pi}{n}}$  and it is given by

$$z = \sqrt[n]{s} e^{i(\frac{\phi}{n} + k\frac{2\pi}{n})} \quad \text{for } w = se^{i\phi} \text{ with } -\pi < \phi < \pi.$$

Now we introduce a generalization of the notion of continuous branch of  $w^{\frac{1}{n}}$ . We define the notion of continuous branch of  $g^{\frac{1}{n}}$ , where  $g$  is a more general function than  $g(w) = w$ .

Let  $A \subseteq \mathbb{C}$  and  $g : A \rightarrow \mathbb{C} \setminus \{0\}$  be continuous in  $A$ . We say that the function  $f$  is a **continuous branch of  $g^{\frac{1}{n}}$**  in  $A$  if  $f$  is continuous in  $A$  and for every  $w \in A$  we have that  $f(w)$  is an element of  $g(w)^{\frac{1}{n}}$  or, equivalently,

$$f(w)^n = g(w) \quad \text{for every } w \in A.$$

**Example 4.4.3.** Let  $g : A \rightarrow \mathbb{C} \setminus \{0\}$  be continuous in  $A \subseteq \mathbb{C}$ . If there is a continuous branch  $h$  of  $w^{\frac{1}{n}}$  in  $g(A)$ , then  $f = h \circ g$  is a continuous branch of  $g^{\frac{1}{n}}$  in  $A$ .

Indeed,  $f = h \circ g$  is continuous in  $A$  and, since  $h(z)^n = z$  for every  $z \in g(A)$ , we also have  $f(w)^n = h(g(w))^n = g(w)$  for every  $w \in A$ .

**Example 4.4.4.** Let  $g : A \rightarrow \mathbb{C} \setminus \{0\}$  be continuous in  $A \subseteq \mathbb{C}$ . If there is a continuous branch  $h$  of  $\log$  in  $A$ , then  $f = e^{\frac{1}{n}h}$  is a continuous branch of  $g^{\frac{1}{n}}$  in  $A$ .

Indeed,  $f = e^{\frac{1}{n}h}$  is continuous in  $A$  and, since  $e^{h(w)} = g(w)$  for every  $w \in A$ , we also have  $f(w)^n = e^{h(w)} = g(w)$  for every  $w \in A$ .

This is a standard way to produce continuous branches of  $g^{\frac{1}{n}}$  when we know continuous branches of  $\log g$ .

**Proposition 4.8.** *Let  $g : A \rightarrow \mathbb{C} \setminus \{0\}$  be continuous in  $A \subseteq \mathbb{C}$  and  $f$  be any continuous branch of  $g^{\frac{1}{n}}$  in  $A$ . If  $w_0$  is an interior point of  $A$  and  $g$  is differentiable at  $w_0$ , then  $f$  is differentiable at  $w_0$  and  $f'(w_0) = \frac{g'(w_0)f(w_0)}{ng(w_0)}$ . Hence, if  $g$  is holomorphic in the interior of  $A$ , then  $f$  is also holomorphic in the interior of  $A$ .*

*Proof.* We set  $z_0 = f(w_0)$  and  $z = f(w)$  for every  $w \in A$ . Then  $z_0^n = g(w_0)$  and  $z^n = g(w)$ . Since  $f$  is continuous,  $w \rightarrow w_0$  implies  $z \rightarrow z_0$ . Therefore, using the derivative of the exponential function at  $z_0$ , we see that

$$\frac{f(w) - f(w_0)}{w - w_0} = \frac{z - z_0}{z^n - z_0^n} \frac{g(w) - g(w_0)}{w - w_0} \rightarrow \frac{g'(w_0)}{nz_0^{n-1}} = \frac{g'(w_0)f(w_0)}{ng(w_0)} \quad \text{when } w \rightarrow w_0.$$

Thus  $f$  is differentiable at  $w_0$  and  $f'(w_0) = \frac{g'(w_0)f(w_0)}{ng(w_0)}$ . □

Therefore, if  $g : A \rightarrow \mathbb{C} \setminus \{0\}$  is holomorphic in the open set  $A$ , every continuous branch of  $g^{\frac{1}{n}}$  can be called **holomorphic branch** of  $g^{\frac{1}{n}}$  in  $A$ .

**Example 4.4.5.** We have defined  $n$  distinct continuous branches of  $w^{\frac{1}{n}}$  in the open set  $A$  which results when we exclude any halfline with vertex 0 from the  $w$ -plane. All these branches are holomorphic branches of  $w^{\frac{1}{n}}$  in  $A$ .

**Proposition 4.9.** *Let  $g : A \rightarrow \mathbb{C} \setminus \{0\}$  be continuous in  $A \subseteq \mathbb{C}$ . Let also  $\omega_n = e^{i\frac{2\pi}{n}}$  be the principal  $n$ -th root of unity.*

(i) *If  $f_1$  is a continuous branch of  $g^{\frac{1}{n}}$  in  $A$  and  $\frac{f_2}{f_1} = \omega_n^k$  in  $A$ , where  $k = 0, 1, \dots, n-1$  is fixed, then  $f_2$  is also a continuous branch of  $g^{\frac{1}{n}}$  in  $A$ .*

(ii) *If, moreover,  $A$  is connected and  $f_1, f_2$  are continuous branches of  $g^{\frac{1}{n}}$  in  $A$ , then  $\frac{f_2}{f_1} = \omega_n^k$  in  $A$ , where  $k = 0, 1, \dots, n-1$  is fixed. In particular, if  $f_1(w_0) = f_2(w_0)$  for some  $w_0 \in A$ , then  $f_1 = f_2$  in  $A$ .*

*Proof.* (i) The continuity of  $f_1$  in  $A$  implies the continuity of  $f_2$  in  $A$ . We also have  $f_1(w)^n = g(w)$  for every  $w \in A$  and hence  $f_2(w)^n = f_1(w)^n (\omega_n^k)^n = g(w)(\omega_n^n)^k = g(w)$  for every  $w \in A$ . Thus,  $f_2$  is a continuous branch of  $g^{\frac{1}{n}}$  in  $A$ .

(ii) For each  $w \in A$  the numbers  $f_2(w), f_1(w)$  are elements of  $g(w)^{\frac{1}{n}}$ . Hence  $(\frac{f_2(w)}{f_1(w)})^n = \frac{g(w)}{g(w)} = 1$  and so  $\frac{f_2}{f_1} : A \rightarrow \{1, \omega_n, \dots, \omega_n^{n-1}\}$ . Now, the function  $\frac{f_2}{f_1}$  is continuous in  $A$  and  $A$  is connected, hence the set  $\frac{f_2}{f_1}(A)$  is also connected. Since  $\frac{f_2}{f_1}(A) \subseteq \{1, \omega_n, \dots, \omega_n^{n-1}\}$ , the set  $\frac{f_2}{f_1}(A)$  contains only one point. I.e.  $\frac{f_2}{f_1}$  is constant in  $A$  and hence  $\frac{f_2}{f_1} = \omega_n^k$  in  $A$ , where  $k = 0, 1, \dots, n-1$  is fixed.

In case  $f_2(w_0) = f_1(w_0)$ , then the integer  $k$  is 1 and we get  $f_2 = f_1$  in  $A$ . □

Thus, if we know one continuous branch of  $g^{\frac{1}{n}}$  in the connected set  $A$ , then we can find every other of the  $n$  possible continuous branches of  $g^{\frac{1}{n}}$  in  $A$  by multiplying the known branch with any constant  $n$ -th root of unity.

**Example 4.4.6.** Let  $A = A_{-\pi} = \{se^{i\phi} \mid s > 0, -\pi < \phi < \pi\}$  be the  $w$ -plane without the negative  $u$ -semiaxis (where  $w = u + iv$ ). We want to find a continuous branch of the square root  $w^{\frac{1}{2}}$  in  $A$  taking the value  $z = 1$  at  $w = 1$ .

From the example 4.4.2 we already know the continuous branch of the square root which maps  $A$

onto the angular region  $B_{-\pi/2} = \{re^{i\theta} \mid r > 0, -\frac{\pi}{2} < \theta < \frac{\pi}{2}\}$ , i.e. onto the right halfplane of the  $z$ -plane, which is given by

$$z = \sqrt{s} e^{i\frac{\phi}{2}} \quad \text{for } w = se^{i\phi} \text{ with } -\pi < \phi < \pi.$$

Since  $A$  is connected, there is no other continuous branch of the square root in  $A$  taking the value  $z = 1$  at  $w = 1$ .

**Example 4.4.7.** Let  $A = A_{-\pi} = \{se^{i\phi} \mid s > 0, -\pi < \phi < \pi\}$  again. Now we want to find a continuous branch of the square root  $w^{\frac{1}{2}}$  in  $A$  taking the value  $z = -1$  at  $w = 1$ .

In the previous example we found one continuous branch of the square root in  $A$ . Since  $A$  is connected, there are exactly two continuous branches of the square root in  $A$ . We consider the principal square root of 1, i.e.  $\omega_2 = e^{i\frac{2\pi}{2}} = e^{i\pi} = -1$ . (Trivial: the square roots of 1 are the solutions of  $z^2 = 1$ , i.e. the numbers 1,  $-1$ .) Then the second continuous branch of the square root in  $A$  is given by

$$z = \sqrt{s} e^{i\frac{\phi}{2}} \omega_2 = -\sqrt{s} e^{i\frac{\phi}{2}} \quad \text{for } w = se^{i\phi} \text{ with } -\pi < \phi < \pi,$$

i.e. the opposite of the previous branch. This branch maps  $A$  onto the angular region  $B_{(-\pi+2\pi)/2} = B_{\pi/2} = \{re^{i\theta} \mid r > 0, \frac{\pi}{2} < \theta < \frac{3\pi}{2}\}$ , i.e. onto the left halfplane of the  $z$ -plane.

### Exercises.

**4.4.1.** Describe the sets  $(-1)^{\frac{1}{2}}, (-1)^{\frac{1}{3}}, (-1)^{\frac{1}{4}}, i^{\frac{1}{2}}, i^{\frac{1}{3}}, i^{\frac{1}{4}}, (\frac{1-i\sqrt{3}}{2})^{\frac{1}{2}}, (\frac{1-i\sqrt{3}}{2})^{\frac{1}{3}}, (\frac{1-i\sqrt{3}}{2})^{\frac{1}{4}}$ .

**4.4.2.** (i) Find the elements of  $\log(i^2)$  and of  $2 \log i$  and observe that the two sets are different.

(ii) Prove that for every  $w \neq 0$  and every  $n \in \mathbb{N}$  the sets  $\log(w^{\frac{1}{n}})$  and  $\frac{1}{n} \log w$  are equal.

**4.4.3.** Let  $w \neq 0$  and  $z$  be any of the elements of  $w^{\frac{1}{n}}$ . Prove that the elements of  $w^{\frac{1}{n}}$  are the numbers  $z, z\omega_n, z\omega_n^2, \dots, z\omega_n^{n-1}$ .

**4.4.4.** The set  $\mathbb{C}^* = \mathbb{C} \setminus \{0\}$  is a group under multiplication. Let  $n \in \mathbb{N}, n \geq 2$ .

(i) Prove that the  $n$ -th root of unity, i.e. the set  $\{1, \omega_n, \omega_n^2, \dots, \omega_n^{n-1}\}$ , is a subgroup of  $\mathbb{C}^*$ .

(ii) Let  $z = \omega_n^k$  be any of the elements of the  $n$ -th root of unity and  $\langle z \rangle = \{z^m \mid m \in \mathbb{Z}\}$  be the group generated by  $z$ . Prove that  $z$  is a generator of  $\{1, \omega_n, \omega_n^2, \dots, \omega_n^{n-1}\}$  or, equivalently,  $\langle z \rangle = \{1, \omega_n, \omega_n^2, \dots, \omega_n^{n-1}\}$  if and only if  $\gcd\{k, n\} = 1$ .

(iii) Prove that  $\{1, \omega_n, \omega_n^2, \dots, \omega_n^{n-1}\}$  has no subgroups other than  $\{1\}$  and itself if and only if  $n$  is a prime number.

**4.4.5.** Look at exercise 3.3.2. Consider the curves on the  $z$ -plane with equations  $x^2 - y^2 = \alpha$  and  $2xy = \beta$ . If the two curves intersect at a point  $(x_0, y_0)$ , find in two ways their angle at this point.

**4.4.6.** Prove that there is no continuous branch of  $w^{\frac{1}{n}}$  in any circle  $C_0(r)$  and hence in any set  $A$  which contains such a circle.

**4.4.7.** Find the continuous branches of the square root in  $\mathbb{C} \setminus [0, +\infty)$ .

**4.4.8.** Find the continuous branches of the cube root in  $\mathbb{C} \setminus (-\infty, 0]$ .

**4.4.9.** (i) Considering a holomorphic branch of  $(w+1)^{\frac{1}{2}}$  in  $\mathbb{C} \setminus (-\infty, -1]$  and a holomorphic branch of  $(w-1)^{\frac{1}{2}}$  in  $\mathbb{C} \setminus [1, +\infty)$ , prove that there is a holomorphic branch of  $(w^2-1)^{\frac{1}{2}}$  in  $\Omega = \mathbb{C} \setminus ((-\infty, -1] \cup [1, +\infty))$ .

(ii) Considering a holomorphic branch of  $(w+1)^{\frac{1}{2}}$  in  $\mathbb{C} \setminus (-\infty, -1]$  and a holomorphic branch of  $(w-1)^{\frac{1}{2}}$  in  $\mathbb{C} \setminus (-\infty, 1]$ , prove that there is a holomorphic branch of  $(w^2-1)^{\frac{1}{2}}$  in  $\Omega' = \mathbb{C} \setminus [-1, 1]$ . (This is not as easy as (i).)

(iii) Prove that there is no continuous branch of  $(w^2-1)^{\frac{1}{2}}$  in any circle which surrounds one of the points  $\pm 1$  but not the other.

**4.4.10.** Prove that we can define a holomorphic branch  $f$  of  $(1-w)^{\frac{1}{2}} + (1+w)^{\frac{1}{2}}$  in the region  $A$  which results when we exclude from  $\mathbb{C}$  two non-intersecting halflines, one with vertex  $+1$  and another with vertex  $-1$ . Prove that every such  $f$  satisfies  $f(w)^4 - 4f(w)^2 + 4w^2 = 0$  for every  $w \in A$ . How many such branches  $f$  exist in  $A$ ?

**4.4.11.** (i) Let  $w \neq 0$  and  $a \in \mathbb{Z}$ . Prove that  $\{e^{az} \mid z \in \log w\}$  has only one element, namely  $w^a$ .  
(ii) Generalizing (i), let  $w \neq 0$  and  $a \notin \mathbb{Z}$ . We define  $w^a = \{e^{az} \mid z \in \log w\}$  and this set may have more than one elements. When does  $w^a$  have finitely many elements and when does it have infinitely many elements?

(iii) Describe the sets  $(\frac{1-i\sqrt{3}}{2})^{\frac{1}{2}}, i^{\frac{1}{4}}, 2^i, i^{\sqrt{2}}$  and draw their elements.

(iv) Prove that the elements of  $w^{a+b}$  are also elements of  $w^a w^b$ , and that the elements of  $w^{ab}$  are also elements of  $(w^a)^b$ .

(v) Let  $f$  be a continuous branch of  $\log$  in  $A \subseteq \mathbb{C} \setminus \{0\}$ . Prove that  $g = e^{af}$  is a continuous branch of  $w^a$  in  $A$  and that  $g$  is differentiable at every interior point  $w_0$  of  $A$  and  $g'(w_0) = \frac{ag(w_0)}{w_0}$ .

(vi) Prove that there is a unique holomorphic branch  $f$  of  $(1-w)^i = e^{i \log(1-w)}$  in  $\mathbb{D}$  so that  $f(0) = 1$ . Then prove that there are  $c_1, c_2 > 0$  so that  $c_1 < |f(w)| < c_2$  for every  $w \in \mathbb{D}$ . Find the best such  $c_1, c_2$ .

**4.4.12.** We define

$$\arccos w = \{z \mid \cos z = w\}, \quad \arcsin w = \{z \mid \sin z = w\}, \quad \arctan w = \{z \mid \tan z = w\}.$$

(i) Prove that the three sets are non-empty, except in the case of  $\arctan(\pm i)$ .

(ii) Express  $\arccos$ ,  $\arcsin$  and  $\arctan$  in terms of  $\log$ .

(iii) It should be clear from exercise 4.2.6 that  $\sin$  is one-to-one from  $\{x+iy \mid -\frac{\pi}{2} < x < \frac{\pi}{2}\}$  onto  $\Omega = \mathbb{C} \setminus ((-\infty, -1] \cup [1, +\infty))$ . Prove that the inverse function  $g_0$  is a continuous branch of  $\arcsin$  in  $\Omega$ , i.e.  $g_0$  is continuous in  $\Omega$  and  $\sin g_0(w) = w$  for every  $w \in \Omega$ . Describe all continuous branches  $g$  of  $\arcsin$  in  $\Omega$  and prove that they are holomorphic in  $\Omega$  with  $g'(w) = 1/(1-w^2)^{\frac{1}{2}}$  for every  $w \in \Omega$ , where at the denominator appears a specific holomorphic branch of  $(1-w^2)^{\frac{1}{2}}$  in  $\Omega$ .

(iv) From exercise 4.2.6 again, it is clear that  $\cos$  is one-to-one from  $\{x+iy \mid 0 < x < \pi\}$  onto  $\Omega = \mathbb{C} \setminus ((-\infty, -1] \cup [1, +\infty))$ . Prove that the inverse function  $h_0$  is a continuous branch of  $\arccos$  in  $\Omega$ , i.e.  $h_0$  is continuous in  $\Omega$  and  $\cos h_0(w) = w$  for every  $w \in \Omega$ . Describe all continuous branches  $h$  of  $\arccos$  in  $\Omega$  and prove that they are holomorphic in  $\Omega$  with  $h'(w) = -1/(1-w^2)^{\frac{1}{2}}$  for every  $w \in \Omega$ , where at the denominator appears a specific holomorphic branch of  $(1-w^2)^{\frac{1}{2}}$  in  $\Omega$ .

(v) Prove that  $\tan$  is one-to-one from  $\{x+iy \mid -\frac{\pi}{2} < x < \frac{\pi}{2}\}$  onto  $U = \mathbb{C} \setminus \{iv \mid v \leq -1 \text{ or } 1 \leq v\}$ . Prove that the inverse function  $k_0$  is a continuous branch of  $\arctan$  in  $U$ , i.e.  $k_0$  is continuous in  $U$  and  $\tan k_0(w) = w$  for every  $w \in U$ . Describe all continuous branches  $k$  of  $\arctan$  in  $U$  and prove that they are holomorphic in  $U$  with  $k'(w) = \frac{1}{1+w^2}$  for every  $w \in U$ .

**4.4.13.** Considering appropriate continuous branches of  $w^{\frac{1}{2}}$ , evaluate  $\int_{\gamma} \frac{1}{w^{1/2}} dw$  for both curves  $\gamma_1(t) = e^{it}, t \in [0, \pi]$ , and  $\gamma_2(t) = e^{-it}, t \in [0, \pi]$ .

## 4.5 Functions defined by curvilinear integrals.

### 4.5.1 Indefinite integrals.

Let the complex functions  $f, F$  be defined in the region  $\Omega \subseteq \mathbb{C}$ . We say that  $F$  is a **primitive** of  $f$  in  $\Omega$  if  $F'(z) = f(z)$  for every  $z \in \Omega$ .

**Proposition 4.10.** *Let the complex function  $f$  be continuous in the region  $\Omega \subseteq \mathbb{C}$ . Then the following are equivalent.*



(i)  $\oint_{\gamma} f(z) dz = 0$  for every piecewise smooth closed curve  $\gamma$  in  $\Omega$ .

(ii)  $\int_{\gamma_1} f(z) dz = \int_{\gamma_2} f(z) dz$  for every two piecewise smooth curves  $\gamma_1, \gamma_2$  in  $\Omega$  with the same endpoints.

(iii) There is a primitive of  $f$  in  $\Omega$ .

*Proof.* (iii)  $\Rightarrow$  (i) Let  $F$  be any primitive of  $f$  in  $\Omega$ . We take an arbitrary piecewise smooth curve  $\gamma : [a, b] \rightarrow \Omega$  with  $\gamma(a) = \gamma(b)$ . Then

$$\begin{aligned} \oint_{\gamma} f(z) dz &= \oint_{\gamma} F'(z) dz = \int_a^b F'(\gamma(t))\gamma'(t) dt = \int_a^b (F \circ \gamma)'(t) dt \\ &= (F \circ \gamma)(b) - (F \circ \gamma)(a) = F(\gamma(b)) - F(\gamma(a)) = 0. \end{aligned}$$

(i)  $\Rightarrow$  (ii) Assume that the piecewise smooth curves  $\gamma_1, \gamma_2$  in  $\Omega$  have the same endpoints. Then the piecewise smooth curve  $\gamma = \gamma_1 + (-\gamma_2)$  is a closed curve in  $\Omega$  and then

$$\int_{\gamma_1} f(z) dz - \int_{\gamma_2} f(z) dz = \int_{\gamma_1} f(z) dz + \int_{-\gamma_2} f(z) dz = \oint_{\gamma} f(z) dz = 0.$$

(ii)  $\Rightarrow$  (iii) We consider an arbitrary fixed  $z_0 \in \Omega$ . Then for every  $z \in \Omega$  there is at least one piecewise smooth curve  $\gamma$  in  $\Omega$  with initial point  $z_0$  and final point  $z$ . We define the function  $F : \Omega \rightarrow \mathbb{C}$  by

$$F(z) = \int_{\gamma} f(\zeta) d\zeta. \quad (4.9)$$

This formula defines  $F(z)$  uniquely, since the value of the curvilinear integral depends only on the point  $z$  and not on the particular piecewise smooth curve  $\gamma$  which we use to join  $z_0$  to  $z$ .

Now we shall prove that  $F$  is a primitive of  $f$  in  $\Omega$ . We take an arbitrary  $z \in \Omega$  and a disc  $D_z(r) \subseteq \Omega$ . We also take a piecewise smooth curve  $\gamma$  in  $\Omega$  with initial point  $z_0$  and final point  $z$ . Then the value of  $F(z)$  is given by (4.9). Now we consider any  $w \in D_z(r)$  and the curve  $\gamma + [z, w]$ . This curve is in  $\Omega$ , it is piecewise smooth and has initial point  $z_0$  and final point  $w$ . Therefore,

$$F(w) = \int_{\gamma + [z, w]} f(\zeta) d\zeta = \int_{\gamma} f(\zeta) d\zeta + \int_{[z, w]} f(\zeta) d\zeta. \quad (4.10)$$

From (4.9) and (4.10) we get

$$F(w) - F(z) - f(z)(w - z) = \int_{[z, w]} f(\zeta) d\zeta - f(z) \int_{[z, w]} d\zeta = \int_{[z, w]} (f(\zeta) - f(z)) d\zeta. \quad (4.11)$$

Now, since  $f$  is continuous, for every  $\epsilon > 0$  there is  $\delta > 0$  so that  $|f(\zeta) - f(z)| < \epsilon$  for every  $\zeta \in \Omega$  with  $|\zeta - z| < \delta$ . Taking  $w \in D_z(r)$  with  $|w - z| < \delta$  we automatically have  $|\zeta - z| < \delta$  for every  $\zeta \in [z, w]$  and (4.11) implies

$$|F(w) - F(z) - f(z)(w - z)| \leq \epsilon|w - z|.$$

Therefore,  $\left| \frac{F(w) - F(z)}{w - z} - f(z) \right| \leq \epsilon$  for every  $w$  with  $0 < |w - z| < \delta$  and hence  $F'(z) = f(z)$ .  $\square$

Let the complex function  $f$  be continuous in the region  $\Omega \subseteq \mathbb{C}$ . If either one of the equivalent conditions (i), (ii) of proposition 4.10 is satisfied, then as we saw in the proof of (ii)  $\Rightarrow$  (iii) of proposition 4.10, we may choose a fixed point  $z_0 \in \Omega$  and define  $F(z) = \int_{\gamma} f(\zeta) d\zeta$  for every  $z \in \Omega$ , where  $\gamma$  is an arbitrary piecewise smooth curve in  $\Omega$  with initial point  $z_0$  and final point  $z$ . Now, any function  $F$  of the form

$$F(z) = \int_{\gamma} f(\zeta) d\zeta + c \quad \text{for every } z \in \Omega,$$

where  $\gamma$  is any piecewise smooth curve in  $\Omega$  with fixed (but otherwise arbitrary) initial point  $z_0$  and final point  $z$  and where  $c$  is an arbitrary constant, is called **indefinite integral** of  $f$  in  $\Omega$ .

The crucial condition for the existence of an indefinite integral is (ii) (or its equivalent (i)) of proposition 4.10. As soon as this is satisfied, then by changing the *base point*  $z_0 \in \Omega$  or the constant  $c$  we get different indefinite integrals  $F$ .

In the proof of proposition 4.10 we saw that every indefinite integral of  $f$  is a primitive of  $f$ . The converse is also true. Indeed, let  $F$  be any primitive of  $f$  in the region  $\Omega$ , i.e. let  $F'(z) = f(z)$  for every  $z \in \Omega$ . Proposition 4.10 implies that condition (ii) is satisfied and, if we take any piecewise smooth curve  $\gamma : [a, b] \rightarrow \Omega$  with initial point a fixed  $z_0 \in \Omega$  and final point  $z \in \Omega$ , then

$$\begin{aligned} \int_{\gamma} f(\zeta) d\zeta &= \int_{\gamma} F'(\zeta) d\zeta = \int_a^b F'(\gamma(t))\gamma'(t) dt = \int_a^b (F \circ \gamma)'(t) dt \\ &= (F \circ \gamma)(b) - (F \circ \gamma)(a) = F(z) - F(z_0). \end{aligned} \quad (4.12)$$

Thus,  $F$  has the form  $F(z) = \int_{\gamma} f(\zeta) d\zeta + F(z_0)$  and hence it is an indefinite integral of  $f$  in  $\Omega$ .

We summarize. *Let the complex function  $f$  be continuous in the region  $\Omega \subseteq \mathbb{C}$ . Then the notion of primitive of  $f$  in  $\Omega$  coincides with the notion of indefinite integral of  $f$  in  $\Omega$ . Moreover, the existence of a primitive or, equivalently, of an indefinite integral of  $f$  in  $\Omega$  is equivalent to the validity of condition (ii) (or (i)) of proposition 4.10.*

Regarding the number of possible primitives of  $f$  in  $\Omega$  we may easily see that, if there is at least one primitive  $F$  of  $f$  in  $\Omega$ , then all others are of the form  $F + c$  for an arbitrary constant  $c$ . Indeed, it is obvious that  $F + c$  is a primitive of  $f$  in  $\Omega$ . Conversely, if  $G$  is a primitive of  $f$  in  $\Omega$ , then we have  $(G - F)'(z) = G'(z) - F'(z) = f(z) - f(z) = 0$  for every  $z \in \Omega$ . Now, theorem 3.3 implies that  $G - F$  is a constant in  $\Omega$ .

Since it is useful for calculations of curvilinear integrals, we state relation (4.12) as a separate proposition.

**Proposition 4.11.** *Let  $F$  be a primitive of the continuous function  $f$  in the region  $\Omega \subseteq \mathbb{C}$ . Then for every piecewise smooth curve  $\gamma$  in  $\Omega$  with initial endpoint  $z_1$  and final endpoint  $z_2$  we have  $\int_{\gamma} f(z) dz = F(z_2) - F(z_1)$ .*

**Example 4.5.1.** Every polynomial function  $p(z) = a_0 + a_1z + \dots + a_nz^n$  has the primitive  $a_0z + \frac{a_1}{2}z^2 + \dots + \frac{a_n}{n+1}z^{n+1}$  in  $\mathbb{C}$ . Therefore, we have  $\oint_{\gamma} p(z) dz = 0$  for every piecewise smooth closed curve  $\gamma$ .

In particular, if  $n \in \mathbb{Z}$ ,  $n \geq 0$ , we have  $\oint_{\gamma} (z - z_0)^n dz = 0$  for every piecewise smooth closed curve  $\gamma$ . A very special case of this, with the circle  $C_{z_0}(r)$ , we saw in examples 2.3.9 and 4.2.2.

**Example 4.5.2.** The exponential function  $e^z$  has the primitive  $e^z$  in  $\mathbb{C}$ . Hence  $\oint_{\gamma} e^z dz = 0$  for every piecewise smooth closed curve  $\gamma$ .

**Example 4.5.3.** Let  $z_0 \in \mathbb{C}$  and  $n \in \mathbb{N}$ ,  $n \geq 2$ . Then the function  $\frac{1}{(z-z_0)^n}$  has the primitive  $-\frac{1}{(n-1)(z-z_0)^{n-1}}$  in  $\mathbb{C} \setminus \{z_0\}$ . Therefore,  $\oint_{\gamma} \frac{1}{(z-z_0)^n} dz = 0$  for every piecewise smooth closed curve  $\gamma$  in  $\mathbb{C} \setminus \{z_0\}$ . A very special case of this, with the circle  $C_{z_0}(r)$ , we saw in examples 2.3.9 and 4.2.2.

**Example 4.5.4.** The function  $\frac{1}{z-z_0}$  (the case  $n = 1$  of the previous example) has no primitive in  $\mathbb{C} \setminus \{z_0\}$  or even in any open ring  $D_{z_0}(r_1, r_2) = \{z \mid r_1 < |z - z_0| < r_2\}$ .

Indeed, if  $\frac{1}{z-z_0}$  had a primitive in  $D_{z_0}(r_1, r_2)$ , then we would have  $\oint_{\gamma} \frac{1}{z-z_0} dz = 0$  for every piecewise smooth closed curve  $\gamma$  in  $D_{z_0}(r_1, r_2)$ . Now, if we take a radius  $r$  so that  $r_1 < r < r_2$  and the curve  $\gamma : [0, 2\pi] \rightarrow D_{z_0}(r_1, r_2)$  with parametric equation  $\gamma(t) = z_0 + re^{it}$ , then we have  $\oint_{\gamma} \frac{1}{z-z_0} dz = \oint_{C_{z_0}(r)} \frac{1}{z-z_0} dz = \int_0^{2\pi} \frac{1}{re^{it}} rie^{it} dt = 2\pi i \neq 0$ . In fact, we did exactly the same calculation in example 4.2.2.

The following result is important.

**Theorem 4.1.** *Let  $g : \Omega \rightarrow \mathbb{C} \setminus \{0\}$  be holomorphic in the region  $\Omega \subseteq \mathbb{C}$  and let  $g'$  be continuous in  $\Omega$ . Then a holomorphic branch of  $\log g$  exists in  $\Omega$  if and only if  $\oint_{\gamma} \frac{g'(z)}{g(z)} dz = 0$  for every piecewise smooth closed curve  $\gamma$  in  $\Omega$ .*

*Proof.* Assume that there is a holomorphic branch of  $\log g$  in  $\Omega$ , i.e. there is  $F$  holomorphic in  $\Omega$  so that  $e^{F(z)} = g(z)$  for every  $z \in \Omega$ . Then  $F'(z)e^{F(z)} = g'(z)$  for every  $z \in \Omega$  and hence  $F'(z) = \frac{g'(z)}{g(z)}$  for every  $z \in \Omega$ . Therefore,  $F$  is a primitive of  $\frac{g'}{g}$  in  $\Omega$  and thus,  $\oint_{\gamma} \frac{g'(z)}{g(z)} dz = 0$  for every piecewise smooth closed curve  $\gamma$  in  $\Omega$ .

Conversely, assume  $\oint_{\gamma} \frac{g'(z)}{g(z)} dz = 0$  for every piecewise smooth closed curve  $\gamma$  in  $\Omega$ . Then  $\frac{g'}{g}$  has a primitive, say  $F$ , in  $\Omega$ . Now, we have  $\frac{d}{dz}(g(z)e^{-F(z)}) = g'(z)e^{-F(z)} - g(z)F'(z)e^{-F(z)} = 0$  for every  $z \in \Omega$ . This implies that, for some constant  $c$ , we have  $g(z)e^{-F(z)} = c$  for every  $z \in \Omega$ . Since  $c \neq 0$ , there is a constant  $d$  so that  $e^d = c$  and we finally get that  $e^{F(z)+d} = g(z)$  for every  $z \in \Omega$ . Now the function  $F + d$  is a holomorphic branch of  $\log g$  in  $\Omega$ .  $\square$

In the next chapter we shall prove that for every holomorphic  $g$  the derivative  $g'$  is automatically continuous. Therefore, *a posteriori*, the assumption in theorem 4.1 that  $g'$  is continuous is unnecessary.

**Example 4.5.5.** If the region  $\Omega \subseteq \mathbb{C} \setminus \{z_0\}$  contains a circle  $C_{z_0}(r)$ , then there is no holomorphic branch of  $\log(z - z_0)$  in  $\Omega$ . In fact, example 4.5.4 shows that  $\oint_{C_{z_0}(r)} \frac{1}{z-z_0} dz \neq 0$ .

**Example 4.5.6.** Let  $g : \Omega \rightarrow \mathbb{C} \setminus \{0\}$  be holomorphic in the region  $\Omega \subseteq \mathbb{C}$ , let  $g'$  be continuous in  $\Omega$  and suppose that there is a halfline with vertex 0 so that  $g(\Omega) \subseteq \mathbb{C} \setminus l$ .

We know that a holomorphic branch of  $\log$  exists in  $\mathbb{C} \setminus l$  and now example 4.3.3 says that a holomorphic branch of  $\log g$  exists in  $\Omega$ . From theorem 4.1 we also get that  $\oint_{\gamma} \frac{g'(z)}{g(z)} dz = 0$  for every piecewise smooth closed curve  $\gamma$  in  $\Omega$ .

## 4.5.2 Integrals with parameter.

**Lemma 4.2.** Let  $n \in \mathbb{N}$  and  $\gamma$  be any piecewise smooth curve. If the complex function  $\phi$  is continuous in the trajectory  $\gamma^*$ , we define  $f(z) = \int_{\gamma} \frac{\phi(\zeta)}{(\zeta-z)^n} d\zeta$  for every  $z \notin \gamma^*$ . Then  $f$  is holomorphic in the open set  $\mathbb{C} \setminus \gamma^*$  and  $f'(z) = n \int_{\gamma} \frac{\phi(\zeta)}{(\zeta-z)^{n+1}} d\zeta$  for every  $z \notin \gamma^*$ .

*Proof.* We take any  $z \in \mathbb{C} \setminus \gamma^*$ . Since  $\mathbb{C} \setminus \gamma^*$  is open, there is  $\delta > 0$  so that  $D_z(\delta) \subseteq \mathbb{C} \setminus \gamma^*$ . We consider the smaller circle  $D_z(\frac{\delta}{2})$  and we have  $|\zeta - w| \geq \frac{\delta}{2}$  for every  $\zeta \in \gamma^*$  and every  $w \in D_z(\frac{\delta}{2})$ . Now for every  $w \in D_z(\frac{\delta}{2})$  we get

$$\frac{f(w)-f(z)}{w-z} - n \int_{\gamma} \frac{\phi(\zeta)}{(\zeta-z)^{n+1}} d\zeta = \int_{\gamma} \left( \frac{\frac{1}{(\zeta-w)^n} - \frac{1}{(\zeta-z)^n}}{w-z} - \frac{n}{(\zeta-z)^{n+1}} \right) \phi(\zeta) d\zeta. \quad (4.13)$$

To simplify the notation, we temporarily set  $a = \zeta - w$  and  $b = \zeta - z$ , and, to estimate the parenthesis in (4.13), we use the algebraic identity

$$\frac{\frac{1}{a^n} - \frac{1}{b^n}}{b-a} - \frac{n}{b^{n+1}} = (b-a) \left( \frac{1}{a^n b^2} + \frac{2}{a^{n-1} b^3} + \cdots + \frac{n-1}{a^2 b^n} + \frac{n}{a b^{n+1}} \right).$$

We have that  $|a| \geq \frac{\delta}{2}$  and  $|b| \geq \frac{\delta}{2}$  for every  $\zeta \in \gamma^*$  and  $w \in D_z(\frac{\delta}{2})$  and hence

$$\begin{aligned} \left| \frac{\frac{1}{a^n} - \frac{1}{b^n}}{b-a} - \frac{n}{b^{n+1}} \right| &\leq |b-a| \left( \frac{1}{|a|^n |b|^2} + \cdots + \frac{n}{|a| |b|^{n+1}} \right) \\ &\leq |w-z| \frac{1+2+\cdots+(n-1)+n}{(\delta/2)^{n+2}} \leq |w-z| \frac{n^2 2^{n+2}}{\delta^{n+2}}. \end{aligned} \quad (4.14)$$

Now, (4.13) and (4.14) imply

$$\left| \frac{f(w)-f(z)}{w-z} - n \int_{\gamma} \frac{\phi(\zeta)}{(\zeta-z)^{n+1}} d\zeta \right| \leq |w-z| \frac{n^2 2^{n+2}}{\delta^{n+2}} \sup_{\zeta \in \gamma^*} |\phi(\zeta)| l(\gamma)$$

for every  $w \in D_z(\frac{\delta}{2})$ . Therefore,  $\lim_{w \rightarrow z} \frac{f(w)-f(z)}{w-z} = n \int_{\gamma} \frac{\phi(\zeta)}{(\zeta-z)^{n+1}} d\zeta$  and  $f$  is differentiable at  $z$  with  $f'(z) = n \int_{\gamma} \frac{\phi(\zeta)}{(\zeta-z)^{n+1}} d\zeta$ .  $\square$

Observe that lemma 4.2 justifies the change of order of the operations of integration and differentiation with respect to the parameter  $z$ :

$$f'(z) = \frac{d}{dz} f(z) = \frac{d}{dz} \int_{\gamma} \frac{\phi(\zeta)}{(\zeta-z)^n} d\zeta = \int_{\gamma} \frac{d}{dz} \left( \frac{\phi(\zeta)}{(\zeta-z)^n} \right) d\zeta = n \int_{\gamma} \frac{\phi(\zeta)}{(\zeta-z)^{n+1}} d\zeta.$$

**Proposition 4.12.** *Let  $\gamma$  be any piecewise smooth curve and the complex function  $\phi$  be continuous in the trajectory  $\gamma^*$ . Then the function  $f(z) = \int_{\gamma} \frac{\phi(\zeta)}{\zeta-z} d\zeta$  is infinitely many times differentiable in the open set  $\mathbb{C} \setminus \gamma^*$  and  $f^{(n)}(z) = n! \int_{\gamma} \frac{\phi(\zeta)}{(\zeta-z)^{n+1}} d\zeta$  for every  $z \notin \gamma^*$ .*

*Proof.* Successive applications of lemma 4.2. □

### Exercises.

**4.5.1.** Let  $f, g$  be holomorphic in the region  $\Omega \subseteq \mathbb{C}$  and let  $f', g'$  be continuous in  $\Omega$ .

(i) If  $|f(z) - 1| < 1$  for every  $z \in \Omega$ , prove that  $\oint_{\gamma} \frac{f'(z)}{f(z)} dz = 0$  for every piecewise smooth closed curve  $\gamma$  in  $\Omega$ .

(ii) If  $|f(z) - g(z)| < |g(z)|$  for every  $z \in \Omega$ , prove that  $\oint_{\gamma} \frac{f'(z)}{f(z)} dz = \oint_{\gamma} \frac{g'(z)}{g(z)} dz$  for every piecewise smooth closed curve  $\gamma$  in  $\Omega$ .

**4.5.2.** Let  $\gamma$  be a piecewise smooth curve and the complex function  $\phi$  be continuous in  $\gamma^*$ . We know that the function  $f(z) = \int_{\gamma} \frac{\phi(\zeta)}{\zeta-z} d\zeta$  is holomorphic in  $\mathbb{C} \setminus \gamma^*$ . Prove that  $f$  is holomorphic at  $\infty$ .

**4.5.3.** Let the complex function  $f$  be continuous in  $\mathbb{R}$  and let  $\int_{-\infty}^{+\infty} \frac{|f(t)|}{1+|t|} dt < +\infty$ . Prove that the function  $F(z) = \int_{-\infty}^{+\infty} \frac{f(t)}{t-z} dt$  is holomorphic in  $\mathbb{C} \setminus \mathbb{R}$ .

**4.5.4.** Let the complex function  $f$  be continuous in  $\mathbb{R}$  and  $\int_{-\infty}^{+\infty} |f(t)|e^{M|t|} dt < +\infty$  for every  $M > 0$ . Prove that the function  $F(z) = \int_{-\infty}^{+\infty} f(t)e^{tz} dt$  is holomorphic in  $\mathbb{C}$ .

**4.5.5.** Find the domains of holomorphy of the functions  $f(z) = \int_0^1 \frac{1}{1+tz} dt$ ,  $g(z) = \int_{-1}^1 \frac{e^{tz}}{1+t^2} dt$ ,  $h(z) = \int_0^{+\infty} \frac{e^{tz}}{1+t^2} dt$  and  $k(z) = \int_0^{+\infty} e^{-tz^2} dt$ .

## 4.6 Functions defined by power series.

Every series of the form

$$\sum_{n=0}^{+\infty} a_n (z - z_0)^n = a_0 + a_1(z - z_0) + a_2(z - z_0)^2 + \dots$$

is called **power series** with **center**  $z_0$  and **coefficients**  $a_n$ . The  $R \in [0, +\infty]$  defined by

$$R = \frac{1}{\limsup \sqrt[n]{|a_n|}}$$

is called **radius of convergence** of the power series. (Of course we understand that  $R = 0$  if  $\limsup \sqrt[n]{|a_n|} = +\infty$  and  $R = +\infty$  if  $\limsup \sqrt[n]{|a_n|} = 0$ .)

**Proposition 4.13.** *Let  $\sum_{n=0}^{+\infty} a_n (z - z_0)^n$  be a power series with radius of convergence  $R$ .*

*If  $R = 0$ , then the series converges only at  $z_0$ . If  $R > 0$ , then:*

(i) *The power series converges absolutely at every  $z \in D_{z_0}(R)$ .*

(ii) *The power series diverges at every  $z \notin \overline{D}_{z_0}(R)$ .*

(iii) *The power series converges uniformly in every closed disc  $\overline{D}_{z_0}(r)$  with  $r < R$ .*

(iv) *The sum  $s(z) = \sum_{n=0}^{+\infty} a_n (z - z_0)^n$  is holomorphic in  $D_{z_0}(R)$ . The derivative of  $s$  in  $D_{z_0}(R)$  is the sum  $t(z) = \sum_{n=1}^{+\infty} n a_n (z - z_0)^{n-1}$  of the power series which results from  $\sum_{n=0}^{+\infty} a_n (z - z_0)^n$  by formal termwise differentiation.*

*Proof.* If  $z = z_0$ , then the power series consists only of its constant term  $a_0$  and hence converges. If  $z \neq z_0$ , then by the definition of  $R$  we get  $\overline{\lim} \sqrt[n]{|a_n(z - z_0)^n|} = \overline{\lim} \sqrt[n]{|a_n|} |z - z_0| = \frac{|z - z_0|}{R}$ . The root test of Cauchy for general series implies that the power series converges absolutely if  $|z - z_0| < R$  and diverges if  $|z - z_0| > R$  and this is the content of (i) and (ii).

(iii) Let  $0 < r < R$ . We take any  $R'$  with  $r < R' < R$ . Then  $\overline{\lim} \sqrt[n]{|a_n|} < \frac{1}{R'}$  and so there is  $n_0$  so that  $\sqrt[n]{|a_n|} \leq \frac{1}{R'}$  for every  $n \geq n_0$ . Then for every  $z \in \overline{D}_{z_0}(r)$  we have  $|a_n(z - z_0)^n| = |a_n| |z - z_0|^n \leq (\frac{r}{R'})^n$  for every  $n \geq n_0$ . Since  $\frac{r}{R'} < 1$ , we have  $\sum_{n=0}^{+\infty} (\frac{r}{R'})^n < +\infty$  and the test of Weierstrass implies that the power series  $\sum_{n=0}^{+\infty} a_n(z - z_0)^n$  converges uniformly in  $\overline{D}_{z_0}(r)$ .

(iv) Besides  $\sum_{n=0}^{+\infty} a_n(z - z_0)^n$ , we also consider the power series  $\sum_{n=1}^{+\infty} n a_n(z - z_0)^{n-1}$ . The second power series results from the first by formal termwise differentiation. We shall prove that the second series converges at every  $z \in D_{z_0}(R)$  and that its sum is the derivative of the sum  $s$  of the first series at every  $z \in D_{z_0}(R)$ .

Since  $\sqrt[n]{n} \rightarrow 1$ , we have  $\overline{\lim} \sqrt[n]{|n a_n|} = \overline{\lim} \sqrt[n]{n} \sqrt[n]{|a_n|} = \overline{\lim} \sqrt[n]{|a_n|}$  and the radius of convergence of the series  $\sum_{n=1}^{+\infty} n a_n(z - z_0)^{n-1}$  is also  $R$ . Thus,  $\sum_{n=1}^{+\infty} n a_n(z - z_0)^{n-1}$  converges at every  $z \in D_{z_0}(R)$ . We define  $t(z) = \sum_{n=1}^{+\infty} n a_n(z - z_0)^{n-1}$  at every  $z \in D_{z_0}(R)$ .

Now at every  $z, w \in D_{z_0}(R)$  we have  $s(w) - s(z) = \sum_{n=0}^{+\infty} a_n((w - z_0)^n - (z - z_0)^n)$ . For simplicity, we shall set temporarily  $a = z - z_0$  and  $b = w - z_0$  and then we have

$$\begin{aligned} \frac{s(w) - s(z)}{w - z} - t(z) &= \sum_{n=2}^{+\infty} a_n(b^{n-1} + b^{n-2}a + \dots + ba^{n-2} + a^{n-1} - na^{n-1}) \\ &= (w - z) \sum_{n=2}^{+\infty} a_n(b^{n-2} + 2b^{n-3}a + \dots + (n-2)ba^{n-3} + (n-1)a^{n-2}). \end{aligned} \quad (4.15)$$

We fix  $z \in D_{z_0}(R)$  and  $\delta = \frac{R - |z - z_0|}{2} > 0$ . We also set  $R_1 = |z - z_0| + \delta = R - \delta$ . If  $w \in D_z(\delta)$ , then  $|b| \leq R_1$  and  $|a| \leq R_1$  and (4.15) implies

$$\left| \frac{s(w) - s(z)}{w - z} - t(z) \right| \leq |w - z| \sum_{n=2}^{+\infty} n^2 |a_n| R_1^{n-2}.$$

Since  $\overline{\lim} \sqrt[n]{|n^2 a_n R_1^n|} = \frac{R_1}{R} < 1$ , the last sum is a finite number independent of  $w \in D_z(\delta)$ . Therefore,  $\lim_{w \rightarrow z} \frac{s(w) - s(z)}{w - z} = t(z)$  and  $s$  is differentiable at  $z$  with  $s'(z) = t(z)$ .  $\square$

If  $R$  is the radius of convergence of  $\sum_{n=0}^{+\infty} a_n(z - z_0)^n$ , then the open disc  $D_{z_0}(R)$  is called **disc of convergence** of the power series.

We saw that, if  $0 < R \leq +\infty$ , the sum  $s(z)$  of the power series is a holomorphic function in  $D_{z_0}(R)$ . In fact the derivative of  $s(z)$  is the function  $t(z)$  which is the sum of the power series we get by formal termwise differentiation of the original power series. We saw that the differentiated power series has the same disc of convergence as the original series and hence we may repeat our arguments: the function  $t(z)$  is holomorphic in  $D_{z_0}(R)$  and its derivative, i.e. the second derivative of  $s(z)$ , is the sum of the power series which we get by a second formal termwise differentiation of the original power series. We conclude that *the function  $s(z)$  is infinitely many times differentiable in the disc of convergence  $D_{z_0}(R)$  and*

$$s^{(k)}(z) = \sum_{n=k}^{+\infty} n(n-1) \cdots (n-k+1) a_n (z - z_0)^{n-k} \quad \text{for every } z \in D_{z_0}(R).$$

**Example 4.6.1.** For the power series  $\sum_{n=1}^{+\infty} \frac{z^n}{n}$  we get  $\overline{\lim} \sqrt[n]{|1/n|} = 1$ , and hence  $R = 1$ . The disc of convergence is  $\mathbb{D}$ . If  $s$  is the function defined by the power series in  $\mathbb{D}$ , then  $s'(z) = \sum_{n=1}^{+\infty} z^{n-1} = \frac{1}{1-z}$  for every  $z \in \mathbb{D}$ . We observe that  $-\text{Log}(1-z)$  is defined and is holomorphic in  $\mathbb{D}$ . Its derivative is  $\frac{1}{1-z}$  and its value at 0 is 0. Since the functions  $s(z)$  and  $-\text{Log}(1-z)$  have the same derivative in the region  $\mathbb{D}$  and the same value at 0, we conclude that

$$\sum_{n=1}^{+\infty} \frac{z^n}{n} = -\text{Log}(1-z) \quad \text{for every } z \in \mathbb{D}.$$

We shall come back to this identity when we study the Taylor series of the function  $-\text{Log}(1-z)$  in  $\mathbb{D}$ .

**Example 4.6.2.** For  $\sum_{n=1}^{+\infty} \frac{z^n}{n^2}$  we get  $\overline{\lim} \sqrt[n]{|1/n^2|} = 1$ , and hence  $R = 1$ . The disc of convergence is  $\mathbb{D}$ .

**Example 4.6.3.** For  $\sum_{n=0}^{+\infty} \frac{z^n}{n!}$  we have  $\overline{\lim} \sqrt[n]{|1/n!|} = 0$  and hence  $R = +\infty$ . The disc of convergence is  $\mathbb{C}$ . If  $s$  is the function defined by the power series in  $\mathbb{C}$ , then  $s'(z) = \sum_{n=1}^{+\infty} \frac{z^{n-1}}{(n-1)!} = \sum_{n=0}^{+\infty} \frac{z^n}{n!} = s(z)$  for every  $z$ . Now we have that  $\frac{d}{dz}(e^{-z}s(z)) = -e^{-z}s(z) + e^{-z}s'(z) = 0$  for every  $z$ . Since the value of  $e^{-z}s(z)$  at 0 is 1, we find that  $e^{-z}s(z) = 1$  for every  $z$  and thus

$$\sum_{n=0}^{+\infty} \frac{z^n}{n!} = e^z \quad \text{for every } z.$$

We shall reprove this identity later, when we study the Taylor series of the function  $e^z$ .

On the other hand, since the series  $\sum_{n=0}^{+\infty} \frac{z^n}{n!}$  and  $\sum_{n=0}^{+\infty} \frac{w^n}{n!}$  converge absolutely, proposition 2.3 implies that

$$\begin{aligned} \sum_{n=0}^{+\infty} \frac{z^n}{n!} \sum_{n=0}^{+\infty} \frac{w^n}{n!} &= \sum_{n=0}^{+\infty} \left( \sum_{k=0}^n \frac{z^k}{k!} \frac{w^{n-k}}{(n-k)!} \right) = \sum_{n=0}^{+\infty} \frac{1}{n!} \left( \sum_{k=0}^n \binom{n}{k} z^k w^{n-k} \right) \\ &= \sum_{n=0}^{+\infty} \frac{(z+w)^n}{n!}. \end{aligned}$$

This provides us with a second proof of the identity  $e^z e^w = e^{z+w}$ .

**Example 4.6.4.** For  $\sum_{n=1}^{+\infty} n!z^n$  we have  $\overline{\lim} \sqrt[n]{n!} = +\infty$ , and hence  $R = 0$ . The power series converges only at 0.

Every series of the form

$$\sum_{n=-\infty}^{n=-1} a_n (z - z_0)^n = \dots + \frac{a_{-3}}{(z-z_0)^3} + \frac{a_{-2}}{(z-z_0)^2} + \frac{a_{-1}}{z-z_0}$$

is called **power series of second type** with **center**  $z_0$  and **coefficients**  $a_n$ . The  $R \in [0, +\infty]$  defined by

$$R = \overline{\lim} \sqrt[m]{|a_{-m}|}$$

is called **radius of convergence** of the power series.

The usual power series of the form  $\sum_{n=0}^{+\infty} a_n (z - z_0)^n$  are also called *power series of first type*, to distinguish them from the power series of second type.

We observe that a power series of second type has no meaning at  $z_0$ , in the same way that any power series of first type (with  $a_n \neq 0$  for at least one  $n \geq 1$ ) has no meaning at  $\infty$ . On the other hand, if  $z = \infty$ , then a power series of second type becomes  $\sum_{-\infty}^{n=-1} 0 = 0$  and hence converges with sum 0.

From now on in these notes we shall use the notations

$$D_{z_0}(R, +\infty) = \{z \mid R < |z - z_0|\}, \quad \overline{D}_{z_0}(R, +\infty) = \{z \mid R \leq |z - z_0|\}$$

for the open and the closed unbounded ring with center  $z_0$  and internal radius  $R$ . We also use

$$D_{z_0}(R_1, R_2) = \{z \mid R_1 < |z - z_0| < R_2\}, \quad \overline{D}_{z_0}(R_1, R_2) = \{z \mid R_1 \leq |z - z_0| \leq R_2\}$$

to denote the open and the closed bounded ring with center  $z_0$ , internal radius  $R_1$  and external radius  $R_2$ .

**Proposition 4.14.** Let  $\sum_{-\infty}^{n=-1} a_n (z - z_0)^n$  be a power series of second type with radius of convergence  $R$ .

If  $R = +\infty$ , then the series converges only at  $\infty$ . If  $R < +\infty$ , then

(i) The power series converges absolutely at every  $z \in D_{z_0}(R, +\infty) \cup \{\infty\}$ .

(ii) The power series diverges at every  $z \notin \overline{D}_{z_0}(R, +\infty)$ .

(iii) The power series converges uniformly in every  $\overline{D}_{z_0}(r, +\infty) \cup \{\infty\}$  with  $r > R$ .

(iv) The sum  $s(z) = \sum_{-\infty}^{n=-1} a_n (z - z_0)^n$  is holomorphic in  $D_{z_0}(R, +\infty) \cup \{\infty\}$ . The derivative of  $s$  in  $D_{z_0}(R, +\infty) \cup \{\infty\}$  is the sum  $t(z) = \sum_{-\infty}^{n=-1} n a_n (z - z_0)^{n-1}$  of the power series which results from  $\sum_{-\infty}^{n=-1} a_n (z - z_0)^n$  by formal termwise differentiation.

*Proof.* The easiest way is to reduce a power series of second type to a power series of first type with the simple change of variable  $w = \frac{1}{z-z_0}$ . Then the power series  $\sum_{-\infty}^{n=-1} a_n(z-z_0)^n$  takes the form  $\sum_{-\infty}^{n=-1} a_n w^{-n} = \sum_{m=1}^{+\infty} a_{-m} w^m$  of a power series of first type with center 0. We also observe that  $z$  varies in the unbounded ring  $D_{z_0}(R, +\infty)$  if and only if  $w$  varies in the punctured disc  $D_0(\frac{1}{R}) \setminus \{0\}$ . Also,  $z$  varies in the unbounded ring  $\overline{D}_{z_0}(r, +\infty)$  if and only if  $w$  varies in the punctured disc  $\overline{D}_0(\frac{1}{r}) \setminus \{0\}$ . Now we can use everything we know about the series  $\sum_{m=1}^{+\infty} a_{-m} w^m$  from proposition 4.13 to get the corresponding results about the series  $\sum_{-\infty}^{n=-1} a_n(z-z_0)^n$ . For example, the differentiability of  $\sum_{-\infty}^{n=-1} a_n(z-z_0)^n$  results from the differentiability of  $\sum_{m=1}^{+\infty} a_{-m} w^m$  and the differentiability of the function  $w = \frac{1}{z-z_0}$ . We leave all the details to the reader. We shall only say a few things about the differentiability of  $s(z) = \sum_{-\infty}^{n=-1} a_n(z-z_0)^n$  at  $\infty$ , using again the transformed power series  $s_*(w) = \sum_{m=1}^{+\infty} a_{-m} w^m$ . Since  $s(\infty) = 0$  and  $s_*(0) = 0$ , we have

$$\lim_{z \rightarrow \infty} z(s(z) - s(\infty)) = \lim_{z \rightarrow \infty} z s(z) = \lim_{w \rightarrow 0} (1 + z_0 w) \frac{s_*(w)}{w} = s'_*(0) = a_{-1}.$$

Therefore,  $s$  is differentiable at  $\infty$  □

If  $R$  is the radius of convergence of  $\sum_{-\infty}^{n=-1} a_n(z-z_0)^n$ , then the open ring  $D_{z_0}(R, +\infty)$  is called **ring of convergence** of the power series. In fact the series converges in  $D_{z_0}(R, +\infty) \cup \{\infty\}$ , which is an open set in  $\widehat{\mathbb{C}}$  with respect to the chordal metric.

If  $0 \leq R < +\infty$ , we saw that the sum  $s(z)$  of the power series is a holomorphic function in  $D_{z_0}(R, +\infty) \cup \{\infty\}$ . In fact the derivative of the sum  $s(z)$  of the power series is the function  $t(z)$  which is the sum of the power series we get by formal termwise differentiation of the original power series. The differentiated power series converges in the same set  $D_{z_0}(R, +\infty) \cup \{\infty\}$ . Therefore, we may repeat our arguments: the function  $t(z)$  is holomorphic in  $D_{z_0}(R, +\infty) \cup \{\infty\}$  and its derivative, i.e. the second derivative of  $s(z)$ , is the sum of the power series which we get by a second formal termwise differentiation of the original power series. We conclude that *the function  $s(z)$  is infinitely many times differentiable in  $D_{z_0}(R, +\infty) \cup \{\infty\}$  and*

$$s^{(k)}(z) = \sum_{-\infty}^{n=-1} n(n-1) \cdots (n-k+1) a_n (z-z_0)^{n-k} \quad \text{for every } z \in D_{z_0}(R, +\infty) \cup \{\infty\}.$$

**Example 4.6.5.**  $\sum_{-\infty}^{n=-1} \frac{z^n}{-n} = \sum_{m=1}^{+\infty} \frac{1}{mz^m}$  converges in  $D_0(1, +\infty) \cup \{\infty\} = \widehat{\mathbb{C}} \setminus \overline{\mathbb{D}}$ .

**Example 4.6.6.**  $\sum_{-\infty}^{n=-1} \frac{z^n}{n^2} = \sum_{m=1}^{+\infty} \frac{1}{m^2 z^m}$  converges in  $D_0(1, +\infty) \cup \{\infty\} = \widehat{\mathbb{C}} \setminus \overline{\mathbb{D}}$ .

**Example 4.6.7.**  $\sum_{-\infty}^{n=-1} \frac{z^n}{(-n)!} = \sum_{m=1}^{+\infty} \frac{1}{m! z^m}$  converges in  $D_0(0, +\infty) \cup \{\infty\} = \widehat{\mathbb{C}} \setminus \{0\}$ .

**Example 4.6.8.**  $\sum_{-\infty}^{n=-1} (-n)! z^n = \sum_{m=1}^{+\infty} \frac{m!}{z^m}$  converges only at  $\infty$ .

Finally, we consider a series of the form

$$\sum_{-\infty}^{+\infty} a_n (z-z_0)^n = \cdots + \frac{a_{-2}}{(z-z_0)^2} + \frac{a_{-1}}{z-z_0} + a_0 + a_1(z-z_0) + a_2(z-z_0)^2 + \cdots$$

which consists of a power series of first type and a power series of second type. We assume that  $a_n \neq 0$  for at least one  $n < 0$  and for at least one  $n > 0$ . Then the original series is called **power series of third type** with **center**  $z_0$  and **coefficients**  $a_n$ . The radius of convergence  $R_1$  of  $\sum_{-\infty}^{n=-1} a_n(z-z_0)^n$  and the radius of convergence  $R_2$  of  $\sum_{n=0}^{+\infty} a_n(z-z_0)^n$  are called **radii of convergence** of our power series. We say that  $\sum_{-\infty}^{+\infty} a_n(z-z_0)^n$  converges at  $z$  if both  $\sum_{-\infty}^{n=-1} a_n(z-z_0)^n$  and  $\sum_{n=0}^{+\infty} a_n(z-z_0)^n$  converge at  $z$ , and we say that  $\sum_{-\infty}^{+\infty} a_n(z-z_0)^n$  diverges at  $z$  in all other cases.

A power series of third type with center  $z_0$  has no meaning at the points  $z_0$  and  $\infty$ .

A power series of third type is a combination of a power series of first type and a power series of second type. Therefore, we expect that the properties of a power series of this new type are a combination of properties of power series of the two previous types. Indeed, the next result is a direct combination of propositions 4.13 and 4.14 and we omit the proof.

**Proposition 4.15.** Let  $\sum_{-\infty}^{+\infty} a_n(z - z_0)^n$  be a power series of third type with radii of convergence  $R_1, R_2$ .

If  $R_2 \leq R_1$ , then the series diverges at every  $z$ , except in the case  $0 < R_1 = R_2 = R < +\infty$  and then it may converge only at some  $z \in C_{z_0}(R)$ . If  $R_1 < R_2$ , then

(i) The power series converges absolutely at every  $z \in D_{z_0}(R_1, R_2)$ .

(ii) The power series diverges at every  $z \notin \overline{D}_{z_0}(R_1, R_2)$ .

(iii) The power series converges uniformly in every  $\overline{D}_{z_0}(r_1, r_2)$  with  $R_1 < r_1 < r_2 < R_2$ .

(iv) The sum  $s(z) = \sum_{-\infty}^{+\infty} a_n(z - z_0)^n$  is holomorphic in  $D_{z_0}(R_1, R_2)$ . The derivative of  $s$  in  $D_{z_0}(R_1, R_2)$  is the sum  $t(z) = \sum_{-\infty}^{+\infty} na_n(z - z_0)^{n-1}$  of the power series which results from  $\sum_{-\infty}^{+\infty} a_n(z - z_0)^n$  by formal termwise differentiation.

If  $R_1 < R_2$ , then  $D_{z_0}(R_1, R_2)$  is called **ring of convergence** of  $\sum_{-\infty}^{+\infty} a_n(z - z_0)^n$  and the function  $s$  defined by the power series is infinitely many times differentiable in  $D_{z_0}(R_1, R_2)$ .

**Example 4.6.9.** We consider  $\sum_{-\infty}^{n=-1} \frac{2^n}{-n} z^n + 1 + \sum_{n=1}^{+\infty} \frac{1}{n^2} z^n$ .

Then  $\sum_{-\infty}^{n=-1} \frac{2^n}{-n} z^n$  has radius of convergence  $\frac{1}{2}$  and  $1 + \sum_{n=1}^{+\infty} \frac{1}{n^2} z^n$  has radius of convergence 1. Therefore,  $D_0(\frac{1}{2}, 1)$  is the ring of convergence of  $\sum_{-\infty}^{n=-1} \frac{2^n}{-n} z^n + 1 + \sum_{n=1}^{+\infty} \frac{1}{n^2} z^n$ .

### Exercises.

**4.6.1.** Find the discs of convergence of  $\sum_{n=1}^{+\infty} a_n z^n$  when  $a_n = n^{13}$ ,  $a_n = \frac{1}{n^5}$ ,  $a_n = \frac{1}{n^n}$ ,  $a_n = n^{\ln n}$ ,  $a_n = (\ln n)^n$ ,  $a_n = \frac{n!}{n^n}$ ,  $a_n = \frac{(n!)^2}{n^n}$ ,  $a_n = \frac{(n!)^2}{(2n)!}$ .

**4.6.2.** Find the rings of convergence of  $\sum_{-\infty}^{n=-1} a_n z^n$  when  $a_n = n^3$ ,  $a_n = \frac{1}{n^2}$ ,  $a_n = \frac{1}{2^n}$ ,  $a_n = 3^n$ ,  $a_n = \frac{1}{(-n)!n^n}$ .

**4.6.3.** Find the ring of convergence and the sum of  $\sum_{-\infty}^{n=-1} (-1)^n z^n + \sum_{n=1}^{+\infty} (\frac{1}{2i})^{n+1} z^n$ .

**4.6.4.** (i) Using the geometric series  $\sum_{n=0}^{+\infty} z^n$ , write  $\frac{1}{1-z}$  as a power series with disc of convergence  $D_0(1)$  and as power series with ring of convergence  $D_0(1, +\infty)$ .

(ii) Write  $\frac{1}{(z-3)(z-4)}$  as a power series with disc of convergence  $D_0(3)$ , as a power series with ring of convergence  $D_0(3, 4)$  and as a power series with ring of convergence  $D_0(4, +\infty)$ .

**4.6.5.** If  $m \in \mathbb{N}$ , using the geometric series  $\sum_{n=0}^{+\infty} z^n$ , write  $\frac{1}{(1-z)^m}$  as a power series  $\sum_{n=0}^{+\infty} a_n z^n$ , and determine its disc of convergence.

**4.6.6.** Find the radius of convergence of  $1 + \sum_{n=1}^{+\infty} \frac{a(a+1)\cdots(a+n-1)b(b+1)\cdots(b+n-1)}{1\cdot 2\cdots n\cdot c(c+1)\cdots(c+n-1)} z^n$ , where  $c \neq 0, -1, -2, \dots$ . This power series is called **hypergeometric series** with parameters  $a, b, c$ . Prove that the function  $w = F(z; a, b, c)$ , which is defined by the hypergeometric series in its disc of convergence, is a solution of the differential equation  $z(1-z)w'' + (c - (a+b+1)z)w' - abw = 0$ .

**4.6.7.** (i) Prove that, if two power series of the type  $\sum_{n=0}^{+\infty} a_n(z - z_0)^n$  with positive radii of convergence define the same function in the intersection of their discs of convergence, then the two series coincide, i.e. they have the same coefficients  $a_n$ .

(ii) Prove a result analogous to (i) for two power series of the type  $\sum_{-\infty}^{n=-1} a_n(z - z_0)^n$ .

**4.6.8.** Let  $0 < R < +\infty$ .

(i) If  $\sum_{n=0}^{+\infty} a_n(z - z_0)^n$  converges absolutely for some  $z \in C_{z_0}(R)$ , prove that it converges absolutely for every  $z \in \overline{D}_{z_0}(R)$ .

(ii) If  $\sum_{n=0}^{+\infty} a_n(z - z_0)^n$  converges for some  $z \in C_{z_0}(R)$ , prove that it converges absolutely for every  $z \in D_{z_0}(R)$ .

**4.6.9.** Let  $R', R''$  and  $R$  be the radii of convergence of  $\sum_{n=0}^{+\infty} a_n'(z - z_0)^n$ ,  $\sum_{n=0}^{+\infty} a_n''(z - z_0)^n$  and  $\sum_{n=0}^{+\infty} (a_n' + a_n'')(z - z_0)^n$ , respectively. If  $R' \neq R''$ , prove that  $R = \min\{R', R''\}$ . If  $R' = R''$ , prove that  $R \geq R' = R''$ .



**4.6.10.** Let  $\sum_{n=0}^{+\infty} a_n(z - z_0)^n$  and  $\sum_{n=0}^{+\infty} b_n(z - z_0)^n$  be two power series and assume that  $c_n = a_0b_n + a_1b_{n-1} + \cdots + a_{n-1}b_1 + a_nb_0$  for every  $n \geq 0$ . If the two power series converge in the disc  $D_{z_0}(R)$ , prove that the power series  $\sum_{n=0}^{+\infty} c_n(z - z_0)^n$  also converges in  $D_{z_0}(R)$  and that  $\sum_{n=0}^{+\infty} c_n(z - z_0)^n = \sum_{n=0}^{+\infty} a_n(z - z_0)^n \sum_{n=0}^{+\infty} b_n(z - z_0)^n$  for every  $z \in D_{z_0}(R)$ .

**4.6.11.** Let  $R$  be the radius of convergence of  $\sum_{n=1}^{+\infty} a_n(z - z_0)^n$ . If  $0 < R < +\infty$ , find the radii of convergence of  $\sum_{n=1}^{+\infty} n^k a_n(z - z_0)^n$ ,  $\sum_{n=1}^{+\infty} n! a_n(z - z_0)^n$ ,  $\sum_{n=1}^{+\infty} \frac{a_n}{n!} (z - z_0)^n$ .

**4.6.12.** Let  $k \in \mathbb{N}$ ,  $k \geq 2$ . Find the  $z$  at which  $\sum_{n=1}^{+\infty} \frac{z^{kn}}{n}$  converges.

**4.6.13.** Find the  $z$  at which  $\sum_{n=1}^{+\infty} z^{n!}$  converges.

**4.6.14.** Let  $0 < b < 1$ . Find the ring of convergence of  $\sum_{n=-\infty}^{+\infty} b^{n^2} z^n$ .

**4.6.15.** If  $s(z) = \sum_{n=0}^{+\infty} a_n(z - z_0)^n$  for every  $z \in D_{z_0}(R)$  and  $|a_1| \geq \sum_{n=2}^{+\infty} n|a_n|R^{n-1}$ , prove that  $s$  is one-to-one in  $D_{z_0}(R)$ .

## Chapter 5

# Local behaviour and basic properties of holomorphic functions.

### 5.1 The theorem of Cauchy for triangles.

Let  $\Delta$  be a closed triangular region. We write  $\oint_{\partial\Delta} f(z) dz$  to denote the curvilinear integral over a piecewise smooth curve  $\gamma$  with trajectory  $\gamma^* = \partial\Delta$  which describes the triangle  $\partial\Delta$  once and in the positive direction. For instance, if  $z_1, z_2, z_3$  are the vertices of the triangle in the order which agrees with the positive direction of  $\partial\Delta$ , then a valid curve is  $\gamma = [z_1, z_2] \dot{+} [z_2, z_3] \dot{+} [z_3, z_1]$ . Hence,  $\oint_{\partial\Delta} f(z) dz = \int_{[z_1, z_2]} f(z) dz + \int_{[z_2, z_3]} f(z) dz + \int_{[z_3, z_1]} f(z) dz$ .

Of course there are analogous statements for integrals  $\oint_{\partial R} f(z) dz$ , when  $R$  is a closed rectangular region or, more generally, a closed convex polygonal region.

**The theorem of Cauchy-Goursat.** *If  $f$  is holomorphic in an open set  $\Omega$  which contains the closed triangular region  $\Delta$ , then*

$$\oint_{\partial\Delta} f(z) dz = 0.$$

*Proof.* We write  $I = \oint_{\partial\Delta} f(z) dz$ , and we have to show that  $I = 0$ .

Let  $\Delta = \Delta(z_1, z_2, z_3)$  be the given closed triangular region with vertices  $z_1, z_2, z_3$  written in the order which agrees with the positive direction of  $\partial\Delta$ . We take the points  $w_3, w_1, w_2$ , which are the midpoints of the linear segments  $[z_1, z_2], [z_2, z_3], [z_3, z_1]$ , respectively. Then the closed triangular region  $\Delta(z_1, z_2, z_3)$  splits into the four closed triangular regions  $\Delta^{(1)} = \Delta(z_1, w_3, w_2)$ ,  $\Delta^{(2)} = \Delta(w_3, z_2, w_1)$ ,  $\Delta^{(3)} = \Delta(w_1, z_3, w_2)$  and  $\Delta^{(4)} = \Delta(w_3, w_1, w_2)$  and we define the corresponding curvilinear integrals:  $I^{(1)} = \oint_{\partial\Delta^{(1)}} f(z) dz$ ,  $I^{(2)} = \oint_{\partial\Delta^{(2)}} f(z) dz$ ,  $I^{(3)} = \oint_{\partial\Delta^{(3)}} f(z) dz$  and  $I^{(4)} = \oint_{\partial\Delta^{(4)}} f(z) dz$ . We analyse each of the four integrals into three integrals over the three linear segments of the corresponding triangle, we add the resulting twelve integrals and we observe the cancellations which occur between integrals over pairs of linear segments with opposite directions. We end up with six integrals over six successive linear segments which add up to give the three linear segments of the original triangle  $\partial\Delta$ . The result is  $I = I^{(1)} + I^{(2)} + I^{(3)} + I^{(4)}$ . This implies  $|I| \leq |I^{(1)}| + |I^{(2)}| + |I^{(3)}| + |I^{(4)}|$  and hence  $|I^{(j)}| \geq \frac{1}{4}|I|$  for at least one  $j$ . Now we take the corresponding closed triangular region  $\Delta^{(j)}$  and, for simplicity, we denote it  $\Delta_1$ . We also denote  $I_1$  the corresponding integral  $I^{(j)}$ . We have proved that there is a closed triangular region  $\Delta_1$  contained in the original  $\Delta$  such that, if  $I = \oint_{\partial\Delta} f(z) dz$  and  $I_1 = \oint_{\partial\Delta_1} f(z) dz$ , then  $|I_1| \geq \frac{1}{4}|I|$ . We also observe that  $\text{diam } \Delta_1 = \frac{1}{2} \text{diam } \Delta$ . We may continue inductively and produce a sequence of closed triangular regions  $\Delta_n$  and the corresponding sequence of curvilinear integrals  $I_n = \oint_{\partial\Delta_n} f(z) dz$  so that:

- (i)  $\Delta \supseteq \Delta_1 \supseteq \cdots \supseteq \Delta_n \supseteq \Delta_{n+1} \supseteq \cdots$ ,
- (ii)  $|I_n| \geq \frac{1}{4^n}|I|$ ,
- (iii)  $\text{diam } \Delta_n = \frac{1}{2^n} \text{diam } \Delta$ .

Now, (i), (iii) imply that there is a (unique) point  $z$  contained in all  $\Delta_n$ . In particular,  $z \in \Delta$  and hence  $f$  is differentiable at  $z$ . If we take an arbitrary  $\epsilon > 0$ , then there is  $\delta > 0$  so that  $|\frac{f(\zeta)-f(z)}{\zeta-z} - f'(z)| < \epsilon$  for every  $\zeta$  with  $0 < |\zeta - z| < \delta$ . Thus,

$$|f(\zeta) - f(z) - f'(z)(\zeta - z)| \leq \epsilon |\zeta - z| \quad \text{for every } \zeta \text{ with } |\zeta - z| < \delta. \quad (5.1)$$

Because of (iii), there is some large  $n$  so that  $\text{diam } \Delta_n < \delta$ . Since  $z \in \Delta_n$  and  $\text{diam } \Delta_n < \delta$ , we get  $|\zeta - z| \leq \text{diam } \Delta_n < \delta$  for every  $\zeta \in \partial\Delta_n \subseteq \Delta_n$  and now (5.1) and (iii) imply

$$|f(\zeta) - f(z) - f'(z)(\zeta - z)| \leq \epsilon |\zeta - z| \leq \epsilon \text{diam } \Delta_n = \frac{\epsilon}{2^n} \text{diam } \Delta \quad \text{for every } \zeta \in \partial\Delta_n.$$

Therefore,

$$\left| \oint_{\partial\Delta_n} (f(\zeta) - f(z) - f'(z)(\zeta - z)) d\zeta \right| \leq \frac{\epsilon}{2^n} \text{diam } \Delta l(\partial\Delta_n) \leq \frac{3\epsilon}{4^n} (\text{diam } \Delta)^2. \quad (5.2)$$

Since  $f(z) + f'(z)(\zeta - z)$  is a polynomial function of  $\zeta$ , we get  $\oint_{\partial\Delta_n} (f(z) + f'(z)(\zeta - z)) d\zeta = 0$  from example 4.5.1, and (5.2) becomes

$$|I_n| = \left| \oint_{\partial\Delta_n} f(\zeta) d\zeta \right| \leq \frac{3\epsilon}{4^n} (\text{diam } \Delta)^2.$$

Finally, (ii) implies  $|I| \leq 3\epsilon(\text{diam } \Delta)^2$  and since  $\epsilon > 0$  is arbitrary, we conclude that  $I = 0$ .  $\square$

## 5.2 Primitives and the theorem of Cauchy in convex regions.

**Proposition 5.1.** *If  $f$  is holomorphic in the convex region  $\Omega$ , then  $f$  has a primitive in  $\Omega$ .*

*Proof.* We fix  $z_0 \in \Omega$ . Then for every  $z \in \Omega$  the linear segment  $[z_0, z]$  is contained in  $\Omega$  and we define  $F(z) = \int_{[z_0, z]} f(\zeta) d\zeta$ . We shall prove that  $F$  is a primitive of  $f$  in  $\Omega$ . We take arbitrary  $z, w \in \Omega$  and consider the closed triangular region  $\Delta$  with vertices  $z_0, z, w$ . Since  $\Omega$  is convex,  $\Delta$  is contained in  $\Omega$  and the Cauchy-Goursat theorem implies  $\oint_{\partial\Delta} f(z) dz = 0$ , i.e.  $\int_{[z_0, z]} f(\zeta) d\zeta + \int_{[z, w]} f(\zeta) d\zeta + \int_{[w, z_0]} f(\zeta) d\zeta = 0$ . Therefore  $F(w) - F(z) = \int_{[z, w]} f(\zeta) d\zeta$  and hence

$$F(w) - F(z) - f(z)(w - z) = \int_{[z, w]} f(\zeta) d\zeta - f(z) \int_{[z, w]} d\zeta = \int_{[z, w]} (f(\zeta) - f(z)) d\zeta. \quad (5.3)$$

Since  $f$  is continuous, for every  $\epsilon > 0$  there is  $\delta > 0$  so that  $|f(\zeta) - f(z)| < \epsilon$  for every  $\zeta \in \Omega$  with  $|\zeta - z| < \delta$ . Taking  $w \in \Omega$  with  $|w - z| < \delta$  we automatically have  $|\zeta - z| < \delta$  for every  $\zeta \in [z, w]$  and (5.3) implies

$$|F(w) - F(z) - f(z)(w - z)| \leq \epsilon |w - z|.$$

Therefore,  $\left| \frac{F(w) - F(z)}{w - z} - f(z) \right| \leq \epsilon$  for every  $w$  with  $0 < |w - z| < \delta$  and hence  $F'(z) = f(z)$ .  $\square$

**The theorem of Cauchy in convex regions.** *If  $f$  is holomorphic in the convex region  $\Omega$ , then*

$$\oint_{\gamma} f(z) dz = 0$$

*for every piecewise smooth closed curve  $\gamma$  in  $\Omega$ .*

*Proof.* Direct from propositions 4.10 and 5.1.  $\square$

Now we shall describe a very useful technique to handle curvilinear integrals of holomorphic functions. Every piecewise smooth closed curve  $\gamma$  we shall refer to will be *visually simple*, for instance a circle or a triangle or a rectangle, and we shall be able to distinguish between the points *inside*  $\gamma$  and the points *outside*  $\gamma$ . We assume that  $\gamma$  surrounds every point inside it once and in the positive direction and that it does not surround the points outside it. The points inside  $\gamma$  form

the *region inside*  $\gamma$  and the points outside  $\gamma$  form the *region outside*  $\gamma$ . Then  $\gamma^*$  is the common boundary of the region inside  $\gamma$  and the region outside  $\gamma$ . We shall concentrate on two characteristic cases.

*First case.* Let  $f$  be holomorphic in the open set  $\Omega$  and let  $\gamma$  be a piecewise smooth closed curve in  $\Omega$ . We want to evaluate  $\oint_{\gamma} f(z) dz$ .

If  $\Omega$  is convex, then  $\oint_{\gamma} f(z) dz = 0$ . So let us assume that  $\Omega$  is not convex. To continue, we assume that the region inside  $\gamma$ , call it  $D$ , is contained in  $\Omega$ , and hence  $f$  is holomorphic in  $D$  as well as in  $\partial D = \gamma^*$ . Now our technique is the following. We split  $D$  into specific disjoint open sets  $E_1, \dots, E_m$  so that their boundaries  $\partial E_1, \dots, \partial E_m$  are trajectories of piecewise smooth closed curves  $\sigma_1, \dots, \sigma_m$ , so that  $\overline{D} = \overline{E_1} \cup \dots \cup \overline{E_m}$  and, finally, so that, when we analyse in an appropriate way each of  $\sigma_1, \dots, \sigma_m$  in successive subcurves and drop those subcurves which appear as pairs of opposite curves, the remaining subcurves can be summed up to give the original curve  $\gamma$ . The result is:

$$\oint_{\gamma} f(z) dz = \oint_{\sigma_1} f(z) dz + \dots + \oint_{\sigma_m} f(z) dz.$$

In fact we applied this technique in the proof of the theorem of Cauchy-Goursat.

Now, if the various  $E_1, \dots, E_m$  can be chosen so that each  $\overline{E_1}, \dots, \overline{E_m}$  is contained in a corresponding *convex* open subset of  $\Omega$ , then we conclude that

$$\oint_{\gamma} f(z) dz = \oint_{\sigma_1} f(z) dz + \dots + \oint_{\sigma_m} f(z) dz = 0 + \dots + 0 = 0.$$

*Second case.* Let  $f$  be holomorphic in the open set  $\Omega$  and let  $\gamma, \gamma_1, \dots, \gamma_n$  be  $n + 1$  piecewise smooth closed curves in  $\Omega$ . We want to relate  $\oint_{\gamma} f(z) dz, \oint_{\gamma_1} f(z) dz, \dots, \oint_{\gamma_n} f(z) dz$ .

We assume that the regions inside  $\gamma_1, \dots, \gamma_n$  are disjoint and that they are all contained in the region inside  $\gamma$ . Let us call  $D$  the *intermediate* region, i.e. the set consisting of the points which are inside  $\gamma$  and outside every  $\gamma_1, \dots, \gamma_n$ , i.e. the intersection of the region inside  $\gamma$  and the regions outside  $\gamma_1, \dots, \gamma_n$ . We further assume that  $D$  is a subset of  $\Omega$ , and hence  $f$  is holomorphic in  $D$  as well as in  $\partial D = \gamma^* \cup \gamma_1^* \cup \dots \cup \gamma_n^*$ . Now, here is the technique. We split  $D$  into specific disjoint open sets  $E_1, \dots, E_m$  so that their boundaries  $\partial E_1, \dots, \partial E_m$  are trajectories of piecewise smooth closed curves  $\sigma_1, \dots, \sigma_m$ , so that  $\overline{E} = \overline{E_1} \cup \dots \cup \overline{E_m}$  and, finally, so that, when we analyse in an appropriate way each of  $\sigma_1, \dots, \sigma_m$  in successive subcurves and drop those subcurves which appear as pairs of opposite curves, the remaining subcurves can be summed up to give  $\gamma$  as well as the opposites of  $\gamma_1, \dots, \gamma_n$ . The result is:

$$\oint_{\gamma} f(z) dz - \oint_{\gamma_1} f(z) dz - \dots - \oint_{\gamma_n} f(z) dz = \oint_{\sigma_1} f(z) dz + \dots + \oint_{\sigma_m} f(z) dz.$$

If the various  $E_1, \dots, E_m$  can be chosen so that each  $\overline{E_1}, \dots, \overline{E_m}$  is contained in a corresponding *convex* open subset of  $\Omega$ , then  $\oint_{\sigma_1} f(z) dz + \dots + \oint_{\sigma_m} f(z) dz = 0 + \dots + 0 = 0$  and hence

$$\oint_{\gamma} f(z) dz = \oint_{\gamma_1} f(z) dz + \dots + \oint_{\gamma_n} f(z) dz.$$

**Corollary 5.1.** *Let  $C, C_1, \dots, C_n$  be  $n + 1$  circles and let  $D, D_1, \dots, D_n$  be the corresponding open discs. Assume that  $\overline{D_1}, \dots, \overline{D_n}$  are disjoint and that they are all contained in  $D$ . Consider also the closed region  $M = \overline{D} \setminus (D_1 \cup \dots \cup D_n)$ . If  $f : \Omega \rightarrow \mathbb{C}$  is holomorphic in an open set  $\Omega$  which contains  $M$ , then  $\oint_C f(z) dz = \oint_{C_1} f(z) dz + \dots + \oint_{C_n} f(z) dz$ . Instead of circles we may consider rectangles or triangles or any combination of the three shapes.*

### Exercises.

**5.2.1.** Let  $\gamma_R$  be the piecewise smooth closed curve which is the sum of the linear segment  $[0, R]$ , the arc of the circle  $C_0(R)$  from  $R$  to  $Re^{i\frac{\pi}{4}}$  in the positive direction and the linear segment  $[Re^{i\frac{\pi}{4}}, 0]$ . Also, let  $\sigma_R$  be the curve which describes only the above arc from  $R$  to  $Re^{i\frac{\pi}{4}}$ .

(i) Prove that  $\int_{\sigma_R} e^{-z^2} dz \rightarrow 0$  when  $R \rightarrow +\infty$ .

(ii) Using  $\gamma_R$  appropriately together with the formula  $\int_0^{+\infty} e^{-t^2} dt = \frac{\sqrt{\pi}}{2}$ , prove the formulas for the so-called Fresnel integrals:  $\int_0^{+\infty} \sin t^2 dt = \int_0^{+\infty} \cos t^2 dt = \frac{\sqrt{\pi}}{2\sqrt{2}}$ .

**5.2.2.** Let  $y, R > 0$  and  $\gamma_{R,y}$  be the piecewise smooth closed curve which is the sum of the linear segments  $[-R, R]$ ,  $[R, R + iy]$ ,  $[R + iy, -R + iy]$  and  $[-R + iy, -R]$ .

(i) If  $y > 0$  is constant, prove that  $\int_{[R, R+iy]} e^{-z^2} dz \rightarrow 0$  and  $\int_{[-R+iy, -R]} e^{-z^2} dz \rightarrow 0$  when  $R \rightarrow +\infty$ .

(ii) Using  $\gamma_{R,y}$  appropriately, prove that  $\int_{-\infty}^{+\infty} e^{-(x+iy)^2} dx$  does not depend on  $y \in [0, +\infty)$ .

(iii) Using the formula  $\int_0^{+\infty} e^{-x^2} dt = \frac{\sqrt{\pi}}{2}$ , prove that  $\int_{-\infty}^{+\infty} e^{-x^2} \cos(2xy) dx = \sqrt{\pi} e^{-y^2}$  for every  $y \geq 0$  (and hence for every  $y \leq 0$  also). This identity is very important in harmonic analysis.

### 5.3 Cauchy's formulas for circles and infinite differentiability.

**Cauchy's formula for circles.** If  $f$  is holomorphic in an open set  $\Omega$  containing the closed disc  $\overline{D}_{z_0}(R)$ , then

$$f(z) = \frac{1}{2\pi i} \oint_{C_{z_0}(R)} \frac{f(\zeta)}{\zeta - z} d\zeta \quad \text{for every } z \in D_{z_0}(R).$$

*Proof.* Let  $z \in D_{z_0}(R)$ . We consider any open disc  $D_z(r)$  with  $r < R - |z - z_0|$ . Then  $\overline{D}_z(r) \subseteq D_{z_0}(R)$  and the function  $\frac{f(\zeta)}{\zeta - z}$  is holomorphic in the open set  $\Omega \setminus \{z\}$  which contains the closed region between the circles  $C_z(r)$  and  $C_{z_0}(R)$ . Corollary 5.1 implies

$$\oint_{C_{z_0}(R)} \frac{f(\zeta)}{\zeta - z} d\zeta = \oint_{C_z(r)} \frac{f(\zeta)}{\zeta - z} d\zeta. \quad (5.4)$$

Now, we have  $\oint_{C_z(r)} \frac{1}{\zeta - z} d\zeta = \int_0^{2\pi} \frac{1}{re^{it}} ire^{it} dt = 2\pi i$  and hence

$$\oint_{C_z(r)} \frac{f(\zeta)}{\zeta - z} d\zeta - 2\pi i f(z) = \oint_{C_z(r)} \frac{f(\zeta) - f(z)}{\zeta - z} d\zeta. \quad (5.5)$$

We take  $\epsilon > 0$ . Since  $f$  is continuous at  $z$ , there is  $\delta > 0$  so that  $|f(\zeta) - f(z)| < \epsilon$  for every  $\zeta \in \Omega$  with  $|\zeta - z| < \delta$ . Therefore, if  $r < \delta$ , (5.5) implies

$$\left| \oint_{C_z(r)} \frac{f(\zeta)}{\zeta - z} d\zeta - 2\pi i f(z) \right| \leq \frac{\epsilon}{r} 2\pi r = 2\pi \epsilon.$$

Since  $\epsilon$  is arbitrary, we conclude that  $\lim_{r \rightarrow 0} \oint_{C_z(r)} \frac{f(\zeta)}{\zeta - z} d\zeta = 2\pi i f(z)$ . Now, letting  $r \rightarrow 0$  in (5.4), we get  $\oint_{C_{z_0}(R)} \frac{f(\zeta)}{\zeta - z} d\zeta = 2\pi i f(z)$ .  $\square$

A particular instance of the formula of Cauchy is when we take  $z = z_0$ , the center of the circle  $C_{z_0}(R)$ . Using the parametric equation  $\zeta = z_0 + Re^{it}$ ,  $t \in [0, 2\pi]$ , we get

$$f(z_0) = \frac{1}{2\pi} \int_0^{2\pi} f(z_0 + Re^{it}) dt$$

and this is called **mean value property** of the holomorphic function  $f$ .

**Cauchy's formula for derivatives and circles.** If  $f$  is holomorphic in an open set  $\Omega$  containing the closed disc  $\overline{D}_{z_0}(R)$ , then  $f$  is infinitely many times differentiable at every  $z \in D_{z_0}(R)$  and

$$f^{(n)}(z) = \frac{n!}{2\pi i} \oint_{C_{z_0}(R)} \frac{f(\zeta)}{(\zeta - z)^{n+1}} d\zeta \quad \text{for every } z \in D_{z_0}(R) \text{ and every } n \in \mathbb{N}.$$

*Proof.* Proposition 4.12 says that  $\frac{1}{2\pi i} \oint_{C_{z_0}(R)} \frac{f(\zeta)}{\zeta - z} d\zeta$  is an infinitely many times differentiable function of  $z$  in the disc  $D_{z_0}(R)$ . On the other hand, Cauchy's formula says that this function coincides with the function  $f(z)$  in the same disc. Therefore  $f(z)$  is infinitely many times differentiable in  $D_{z_0}(R)$ . Moreover, the derivatives of  $f(z)$  are the same as the derivatives of  $\frac{1}{2\pi i} \oint_{C_{z_0}(R)} \frac{f(\zeta)}{\zeta - z} d\zeta$  and these are given by the formulas in proposition 4.12.  $\square$

**Example 5.3.1.** Let  $n \in \mathbb{N}$ . Then  $\oint_{C_{z_0}(R)} \frac{1}{(\zeta-z)^n} d\zeta = 0$  for every  $z \notin \overline{D}_{z_0}(R)$ . To see this we observe that the circle  $C_{z_0}(R)$  is contained in a slightly larger open disc  $D_{z_0}(R')$  which does not contain  $z$ : it is enough to take  $R < R' < |z - z_0|$ . Then the disc  $D_{z_0}(R')$  is a convex region and  $\frac{1}{(\zeta-z)^n}$  is a holomorphic function of  $\zeta$  in  $D_{z_0}(R')$ . Now the result is an application of the theorem of Cauchy in convex regions.

On the other hand, for every  $z \in D_{z_0}(R)$  we have  $\oint_{C_{z_0}(R)} \frac{1}{(\zeta-z)^n} d\zeta = 2\pi i$ , if  $n = 1$ , and  $\oint_{C_{z_0}(R)} \frac{1}{(\zeta-z)^n} d\zeta = 0$ , if  $n \geq 2$ . This is a simple application of Cauchy's formula (for a function and its derivatives) to the constant function 1. The special case  $z = z_0$  we have already seen in examples 2.3.9 and 4.2.2.

**Theorem 5.1.** *If  $f$  is holomorphic in the open set  $\Omega$ , then  $f$  is infinitely many times differentiable in  $\Omega$ .*

*Proof.* Let  $z_0 \in \Omega$ . We take a closed disc  $\overline{D}_{z_0}(R) \subseteq \Omega$  and then  $f$  is infinitely many times differentiable in  $D_{z_0}(R)$  and hence at  $z_0$ .  $\square$

It is time to recall the remark after theorem 4.1. The assumption of continuity of the derivative in theorem 4.1 is superfluous. The same we may say for the hypothesis in example 4.5.6 and in exercises 3.3.3 and 4.5.1.

**Cauchy's estimates.** *If  $f$  is holomorphic in an open set containing the closed disc  $\overline{D}_{z_0}(R)$  and if  $|f(\zeta)| \leq M$  for every  $\zeta \in C_{z_0}(R)$ , then*

$$|f^{(n)}(z_0)| \leq \frac{n!M}{R^n} \quad \text{for every } n \in \mathbb{N}.$$

*Proof.* Direct application of Cauchy's formulas.  $\square$

### Exercises.

**5.3.1.** Evaluate  $\oint_{C_0(r)} \frac{z^2+1}{z(z^2+4)} dz$  for  $0 < r < 2$  and for  $2 < r < +\infty$ .

**5.3.2.** If  $n \in \mathbb{N}$ , evaluate  $\oint_{C_0(1)} \frac{e^z}{z^n} dz$  and  $\int_0^{2\pi} e^{\cos \theta} \sin(n\theta - \sin \theta) d\theta$ ,  $\int_0^{2\pi} e^{\cos \theta} \cos(n\theta - \sin \theta) d\theta$ .

**5.3.3.** If  $n \in \mathbb{N}$ , evaluate  $\oint_{C_0(1)} \frac{e^{iz}}{z^n} dz$ ,  $\oint_{C_0(1)} \frac{\sin z}{z^n} dz$ ,  $\oint_{C_0(1)} \frac{e^z - e^{-z}}{z^n} dz$ ,  $\oint_{C_1(\frac{1}{2})} \frac{\text{Log } z}{(z-1)^n} dz$ .

**5.3.4.** Let  $f$  be holomorphic in  $\mathbb{C}$  and let  $|f(z)| \leq A + M|z|^n$  for every  $z$ . Prove that  $f^{(n+1)}(z) = 0$  for every  $z$  and that  $f$  is a polynomial function of degree  $\leq n$ .

**5.3.5.** Let the complex function  $f$  be continuous in  $\overline{D}_{z_0}(R)$  and holomorphic in  $D_{z_0}(R)$ . Prove that  $f(z) = \frac{1}{2\pi i} \oint_{C_{z_0}(R)} \frac{f(\zeta)}{\zeta-z} d\zeta$  for every  $z \in D_{z_0}(R)$ .

**5.3.6.** Let  $f$  be holomorphic in an open set containing the closed disc  $\overline{D}_{z_0}(R)$  and let  $0 < r < R$ . If  $|f(z)| \leq M$  for every  $z \in C_{z_0}(R)$ , find an upper bound for  $|f^{(n)}|$  in  $\overline{D}_{z_0}(r)$ , which depends only on  $n, r, R, M$  and not on  $f$  or  $z_0$ .

**5.3.7.** Let  $f$  be holomorphic in  $D_{z_0}(R)$ . If  $|f(z)| \leq \frac{1}{R-|z-z_0|}$  for every  $z \in D_{z_0}(R)$ , find the smallest possible upper bound for  $|f^{(n)}(z_0)|$ , which depends only on  $n, R$  and not on  $f$  or  $z_0$ .

**5.3.8.** Let  $f$  be holomorphic and bounded in  $\mathbb{D}$ . Prove that  $f(w) = \frac{1}{\pi} \iint_{\mathbb{D}} \frac{f(z)}{(1-zw)^2} dx dy$  for every  $w \in \mathbb{D}$ .

## 5.4 Morera's theorem.

Theorem 5.1 and proposition 4.10 imply the following corollary. *If the complex function  $f$  is continuous in the region  $\Omega \subseteq \mathbb{C}$  and if  $\oint_{\gamma} f(z) dz = 0$  for every piecewise smooth closed curve  $\gamma$  in  $\Omega$ , then  $f$  is holomorphic in  $\Omega$ .* Indeed, since  $\oint_{\gamma} f(z) dz = 0$  for every piecewise smooth closed curve  $\gamma$  in  $\Omega$ , we get that  $f$  has a primitive, say  $F$ , in  $\Omega$ . This means that  $F' = f$  in  $\Omega$  and hence  $F$  is holomorphic in  $\Omega$ . Therefore,  $F$  is infinitely many times differentiable in  $\Omega$  and then  $f$  is also infinitely many times differentiable in  $\Omega$ . In particular,  $f$  is holomorphic in  $\Omega$ .

The next theorem proves the same result with weaker assumptions.

**The theorem of Morera.** *If the complex function  $f$  is continuous in the open set  $\Omega \subseteq \mathbb{C}$  and if  $\oint_{\partial\Delta} f(z) dz = 0$  for every closed triangular region  $\Delta$  in  $\Omega$ , then  $f$  is holomorphic in  $\Omega$ .*

*Proof.* Let  $z_0 \in \Omega$ . We consider a disc  $D_{z_0}(R) \subseteq \Omega$ . This disc is a convex set and we have that  $\oint_{\partial\Delta} f(z) dz = 0$  for every closed triangular region  $\Delta$  in  $D_{z_0}(R)$ . Then the *proof* of proposition 5.1 applies, and we get that  $f$  has a primitive, say  $F$ , in  $D_{z_0}(R)$ . This means that  $F' = f$  in  $D_{z_0}(R)$  and hence  $F$  is holomorphic in  $D_{z_0}(R)$ . Therefore,  $F$  is infinitely many times differentiable in  $D_{z_0}(R)$  and  $f$  is also infinitely many times differentiable in  $D_{z_0}(R)$ . In particular,  $f$  is holomorphic in  $D_{z_0}(R)$  and hence at  $z_0$ .  $\square$

### Exercises.

**5.4.1.** If the complex function  $f$  is continuous in the open set  $\Omega$  and holomorphic in  $\Omega \setminus l$ , where  $l$  is a line, prove that  $f$  is holomorphic in  $\Omega$ .

## 5.5 Liouville's theorem. The fundamental theorem of algebra.

**The theorem of Liouville.** *If  $f$  is holomorphic and bounded in  $\mathbb{C}$ , then  $f$  is constant in  $\mathbb{C}$ .*

*Proof.* There is  $M \geq 0$  so that  $|f(z)| \leq M$  for every  $z$ . We take any  $z_0$  and apply Cauchy's estimate for  $n = 1$  with an arbitrary circle  $C_{z_0}(R)$  and we find that  $|f'(z_0)| \leq \frac{M}{R}$ . Letting  $R \rightarrow +\infty$ , we get  $f'(z_0) = 0$ . Since  $z_0$  is arbitrary, we conclude that  $f$  is constant.  $\square$

**Fundamental theorem of algebra.** *Every polynomial of degree  $\geq 1$  has at least one root in  $\mathbb{C}$ .*

*Proof.* Let  $p$  be a polynomial of degree  $\geq 1$  and assume that  $p$  has no root in  $\mathbb{C}$ .

We consider the function  $f = \frac{1}{p}$ , which is holomorphic in  $\mathbb{C}$ , and we see easily that it is also bounded in  $\mathbb{C}$ . Indeed, since  $\lim_{z \rightarrow \infty} p(z) = \infty$ , we have  $\lim_{z \rightarrow \infty} f(z) = 0$ , and hence there is  $R > 0$  so that  $|f(z)| \leq 1$  for every  $z$  with  $|z| > R$ . Since  $|f|$  is continuous in the compact disc  $\overline{D_0}(R)$ , there is  $M' \geq 0$  so that  $|f(z)| \leq M'$  for every  $z$  with  $|z| \leq R$ . Taking  $M = \max\{M', 1\}$ , we have that  $|f(z)| \leq M$  for every  $z$  and hence  $f$  is bounded.

Liouville's theorem implies that  $f$  and hence  $p$  is constant and we arrive at a contradiction.  $\square$

Having proved that a polynomial  $p$  has a root  $z_1$ , we may prove in a purely algebraic way that  $z - z_1$  is a factor of  $p$ , i.e. there is a polynomial  $p_1$  so that  $p(z) = (z - z_1)p_1(z)$  for every  $z$ . Continuing inductively, we conclude that, if  $n \geq 1$  is the degree of  $p$ , there are  $z_1, \dots, z_n$  so that

$$p(z) = c(z - z_1) \cdots (z - z_n) \quad \text{for every } z$$

where  $c$  is a constant. Thus, every polynomial  $p$  of degree  $n \geq 1$  has exactly  $n$  roots in  $\mathbb{C}$ .

### Exercises.

**5.5.1.** If  $f : \mathbb{C} \rightarrow \mathbb{C}$  is holomorphic in  $\mathbb{C}$  and  $\operatorname{Re} f$  is bounded in  $\mathbb{C}$ , prove that  $f$  is constant in  $\mathbb{C}$ .

**5.5.2.** We say that  $z, w$  are symmetric with respect to  $\mathbb{T}$  if either  $z = 0, w = \infty$  or  $z = \infty, w = 0$  or  $z, w \in \mathbb{C}, z = \frac{1}{\bar{w}}$ .

Let  $r = \frac{p}{q}$  be a non-constant rational function so that the polynomials  $p, q$  have no common root and so that  $r(z) \in \mathbb{T}$  for every  $z \in \mathbb{T}$ . Prove that, if  $a \in \mathbb{C} \setminus \{0\}$  is a root of  $p$  of multiplicity  $k$ , then  $b = \frac{1}{\bar{a}}$  is a root of  $q$  of multiplicity  $k$  and conversely. I.e. the roots of  $p$  and the roots of  $q$  form pairs of points symmetric with respect to  $\mathbb{T}$ . (In particular,  $p$  and  $q$  have the same degree.)

## 5.6 Maximum principle.

**Maximum principle.** Let  $f$  be holomorphic in the region  $\Omega \subseteq \mathbb{C}$  and  $M = \sup_{z \in \Omega} |f(z)|$ . If there is  $z_0 \in \Omega$  so that  $|f(z_0)| = M$ , then  $f$  is constant in  $\Omega$ .

*Proof.* We take any  $z \in \Omega$  for which  $|f(z)| = M$ . We consider an open disc  $D_z(R) \subseteq \Omega$  and any  $r$  with  $0 < r < R$ . The mean value property of  $f$  says that  $f(z) = \frac{1}{2\pi} \int_0^{2\pi} f(z + re^{it}) dt$ . Since  $|f(z + re^{it})| \leq M$  for every  $t \in [0, 2\pi]$ , we have

$$M = |f(z)| = \left| \frac{1}{2\pi} \int_0^{2\pi} f(z + re^{it}) dt \right| \leq \frac{1}{2\pi} \int_0^{2\pi} |f(z + re^{it})| dt \leq M.$$

Hence,  $\frac{1}{2\pi} \int_0^{2\pi} |f(z + re^{it})| dt = M$  and, since  $|f(z + re^{it})|$  is a continuous function of  $t$ , we get  $|f(z + re^{it})| = M$  for every  $t \in [0, 2\pi]$ . Now,  $r$  is arbitrary in the interval  $(0, R)$  and we find that  $|f(z + re^{it})| = M$  for every  $t \in [0, 2\pi]$  and every  $r \in (0, R)$ . So we get  $|f(w)| = M$  for every  $w \in D_z(R)$ . We proved that, if  $|f(z)| = M$  for a  $z \in \Omega$ , then this equality holds in a neighborhood of  $z$ . Now we define

$$B = \{z \in \Omega \mid |f(z)| = M\}, \quad C = \{z \in \Omega \mid |f(z)| < M\}$$

and it is clear that  $B \cup C = \Omega$ .

If  $z \in B$ , then  $|f(z)| = M$  and hence the same is true at every point in a neighborhood of  $z$ . Therefore  $z$  is not a limit point of  $C$ . Moreover, if  $z \in C$  then  $|f(z)| < M$  and, by the continuity of  $f$ , the same is true in a neighborhood of  $z$ . Hence  $z$  is not a limit point of  $B$ .

If both  $B$  and  $C$  are non-empty, then they form a decomposition of  $\Omega$ . But  $\Omega$  is connected and, since  $z_0 \in B$ , we get that  $C = \emptyset$ . Therefore,  $|f(z)| = M$  for every  $z \in \Omega$ .

Now we shall prove that  $f$  is constant in  $\Omega$ . If  $M = 0$ , then clearly  $f = 0$  in  $\Omega$ . So let us assume that  $M > 0$ . If  $u$  and  $v$  are the real and the imaginary part of  $f$ , then  $u^2 + v^2 = M^2$  in  $\Omega$  and hence  $u \frac{\partial u}{\partial x} + v \frac{\partial v}{\partial x} = 0$  and  $u \frac{\partial u}{\partial y} + v \frac{\partial v}{\partial y} = 0$  in  $\Omega$ . Using the C-R equations, we get  $u \frac{\partial u}{\partial x} + v \frac{\partial v}{\partial x} = 0$  and  $v \frac{\partial u}{\partial x} - u \frac{\partial v}{\partial x} = 0$  in  $\Omega$ . Viewing this as a system with unknowns  $\frac{\partial u}{\partial x}, \frac{\partial v}{\partial x}$ , we see that its determinant is  $u^2 + v^2 = M^2 > 0$ , and we find that  $\frac{\partial u}{\partial x} = 0$  and  $\frac{\partial v}{\partial x} = 0$  in  $\Omega$ . Therefore,  $f' = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = 0$  in  $\Omega$  and hence  $f$  is constant in the region  $\Omega$ .  $\square$

**Maximum principle.** Let the complex function  $f$  be holomorphic in the bounded region  $\Omega$  and continuous in  $\bar{\Omega}$ . Then either  $f$  is constant in  $\bar{\Omega}$  or  $|f|$  has a maximum value, say  $M$ , attained at a point of  $\partial\Omega$  and  $|f(z)| < M$  for every  $z \in \Omega$ . In every case,  $|f|$  has a maximum value which is attained at a point of  $\partial\Omega$ .

*Proof.* If  $f$  is constant in  $\bar{\Omega}$ , then  $|f|$  is also constant, say  $M$ , in  $\bar{\Omega}$ . Then, obviously,  $M$  is the maximum value of  $|f|$  and it is attained (everywhere and hence) at every point of  $\partial\Omega$ .

Now we assume that  $f$  is not constant in  $\bar{\Omega}$ . This implies easily that  $f$  is not constant in  $\Omega$  either. Now,  $|f|$  is continuous in the compact set  $\bar{\Omega}$  and hence attains its maximum value, say  $M$ , at some point  $z_0 \in \bar{\Omega}$ . I.e. we have  $|f(z_0)| = M$  and  $|f(z)| \leq M$  for every  $z \in \bar{\Omega}$ .

If any such  $z_0$  belongs to  $\Omega$ , then the previous maximum principle implies that  $f$  is constant in  $\Omega$  and we arrive at a contradiction. We conclude that  $z_0 \in \partial\Omega$  and  $|f(z)| < M$  for every  $z \in \Omega$ .  $\square$

*Exercise 5.6.3 refers to the case of an unbounded region  $\Omega$ .*



### Exercises.

**5.6.1.** Let  $f$  be holomorphic in the region  $\Omega \subseteq \mathbb{C}$  and let  $m = \inf_{z \in \Omega} |f(z)|$ . If there is  $z_0 \in \Omega$  so that  $|f(z_0)| = m$ , prove that either  $m = 0$  (and hence  $f(z_0) = 0$ ) or  $m > 0$  and then  $f$  is constant in  $\Omega$ . This is called **minimum principle**.

**5.6.2.** Let the complex function  $f$  be holomorphic in  $\mathbb{D}$  and continuous in  $\overline{\mathbb{D}}$ , let  $|f(z)| > 1$  for every  $z \in \mathbb{T}$  and  $f(0) = 1$ . Does  $f$  have a root in  $\mathbb{D}$ ?

**5.6.3.** State and prove the second maximum principle in the case of an *unbounded* region  $\Omega \subseteq \mathbb{C}$ . In this case we must include the point  $\infty$  in  $\overline{\Omega}$ .

**5.6.4.** Let  $f$  be holomorphic in the bounded region  $\Omega$  and  $\lim_{z \rightarrow \zeta} f(z) = 0$  for every  $\zeta \in \partial\Omega$ . Prove that  $f$  is constant 0 in  $\Omega$ .

In the case of an *unbounded* region  $\Omega \subseteq \mathbb{C}$ , we must include the point  $\infty$  in  $\partial\Omega$ .

**5.6.5.** Let  $f$  be holomorphic in the region  $\Omega \subseteq \mathbb{C}$  and  $K = \sup_{z \in \Omega} \operatorname{Re} f(z)$ . If there is  $z_0 \in \Omega$  so that  $\operatorname{Re} f(z_0) = K$ , prove that  $f$  is constant in  $\Omega$ .

**5.6.6.** Prove the fundamental theorem of algebra using the maximum principle.

**5.6.7.** Let  $f_n, f$  be holomorphic in the bounded region  $\Omega$  and continuous in  $\overline{\Omega}$ . If  $f_n \rightarrow f$  uniformly in  $\partial\Omega$ , prove that  $f_n \rightarrow f$  uniformly in  $\overline{\Omega}$ .

In the case of an *unbounded* region  $\Omega \subseteq \mathbb{C}$ , we must include the point  $\infty$  in  $\partial\Omega$ .

**5.6.8.** Let  $R$  be a square region with center  $z_0$ . Let  $f$  be holomorphic in  $R$  and continuous in  $\overline{R}$ . If  $|f(z)| \leq m$  for every  $z$  in one of the four sides of  $R$  and  $|f(z)| \leq M$  for every  $z$  in the other three sides of  $R$ , prove that  $|f(z_0)| \leq \sqrt[4]{mM^3}$ .

**5.6.9.** Let  $\Omega = \{x + iy \mid -\frac{\pi}{2} < y < \frac{\pi}{2}\}$  and  $f(z) = e^{e^z}$ . Then  $f$  is holomorphic in  $\Omega$  and continuous in  $\overline{\Omega} = \{x + iy \mid -\frac{\pi}{2} \leq y \leq \frac{\pi}{2}\}$ . Prove that  $|f(x - i\frac{\pi}{2})| = |f(x + i\frac{\pi}{2})| = 1$  for every  $x \in \mathbb{R}$  and that  $\lim_{x \rightarrow +\infty} f(x) = +\infty$ . Does this contradict the maximum principle?

**5.6.10.** Let the complex function  $f$  be holomorphic in the bounded region  $\Omega$  and continuous in  $\overline{\Omega}$ . If  $|f|$  is constant in  $\partial\Omega$ , prove that either  $f$  has at least one root in  $\Omega$  or  $f$  is constant in  $\Omega$ .

**5.6.11.** Let  $f$  be holomorphic in the bounded region  $\Omega$  and continuous in  $\overline{\Omega}$ . If  $\operatorname{Re} f = 0$  in  $\partial\Omega$ , prove that  $f$  is constant in  $\Omega$ .

**5.6.12.** (i) Let  $f$  be holomorphic and non-constant in the region  $\Omega \subseteq \mathbb{C}$ . For every  $\mu > 0$  prove that  $\{z \in \Omega \mid |f(z)| < \mu\} \cap \Omega = \{z \in \Omega \mid |f(z)| \leq \mu\}$ .

(ii) Let  $p$  be a polynomial of degree  $n \geq 1$ . Prove that for every  $\mu > 0$  the set  $\{z \mid |p(z)| < \mu\}$  has at most  $n$  connected components and each of them contains at least one root of  $p$ . How do these connected components behave when  $\mu \rightarrow 0+$  and when  $\mu \rightarrow +\infty$ ?

**5.6.13.** Let  $f$  be holomorphic and non-constant in the bounded region  $\Omega$ . If  $\limsup_{\Omega \ni z \rightarrow \zeta} |f(z)| \leq M$  for every  $\zeta \in \partial\Omega$ , prove that  $|f(z)| < M$  for every  $z \in \Omega$ .

In the case of an *unbounded* region  $\Omega \subseteq \mathbb{C}$ , we must include the point  $\infty$  in  $\partial\Omega$ .

**5.6.14.** Let the complex function  $f$  be holomorphic in the bounded region  $\Omega$  and continuous in  $\overline{\Omega}$ . If  $U$  is an open set so that  $\overline{U} \subseteq \Omega$ , prove that  $\sup_{z \in \partial U} |f(z)| \leq \sup_{z \in \partial\Omega} |f(z)|$ . If equality holds, prove that  $f$  is constant in  $\overline{\Omega}$ .

**5.6.15.** Let  $f$  be holomorphic in  $D_0(R_1, R_2)$  and  $a \in \mathbb{R}$ . Prove that  $|z|^a |f(z)|$  has no maximum value in  $D_0(R_1, R_2)$ , except if  $a \in \mathbb{Z}$  and there is  $c$  so that  $f(z) = cz^{-a}$  for every  $z \in D_0(R_1, R_2)$ .

**5.6.16. The three circles theorem of Hadamard.** Let  $f$  be holomorphic in  $D_{z_0}(R_1, R_2)$  and let  $M(r) = \max_{z \in C_{z_0}(r)} |f(z)|$  for  $R_1 < r < R_2$ . Prove that  $\ln M(r)$  is a convex function of  $\ln r$  in  $(R_1, R_2)$ . I.e. prove that, if  $R_1 < r_1 < r < r_2 < R_2$  and  $\ln r = (1-t)\ln r_1 + t\ln r_2$  for  $0 < t < 1$ , then  $\ln M(r) \leq (1-t)\ln M(r_1) + t\ln M(r_2)$ .

**5.6.17. The three lines theorem.** Let  $f$  be holomorphic and bounded in the vertical zone  $K = \{x + iy \mid X_1 < x < X_2\}$  and let  $M(x) = \sup_{y \in \mathbb{R}} |f(x + iy)|$  for  $X_1 < x < X_2$ . Prove that  $\ln M(x)$  is a convex function of  $x$  in  $(X_1, X_2)$ . I.e. prove that, if  $X_1 < x_1 < x < x_2 < X_2$  and  $x = (1-t)x_1 + tx_2$  for  $0 < t < 1$ , then  $\ln M(x) \leq (1-t)\ln M(x_1) + t\ln M(x_2)$ .

**5.6.18. The Phragmén-Lindelöf theorem.** Let  $f, \phi$  be holomorphic in the bounded region  $\Omega$  and let  $\phi$  be bounded in  $\Omega$  and have no root in  $\Omega$ . Let also  $A \cap B = \emptyset$  and  $A \cup B = \partial\Omega$ . If

(i)  $\limsup_{\Omega \ni z \rightarrow \zeta} |f(z)| \leq M$  for every  $\zeta \in A$  and

(ii)  $\limsup_{\Omega \ni z \rightarrow \zeta} |f(z)||\phi(z)|^\epsilon \leq M$  for every  $\zeta \in B$  and every  $\epsilon > 0$ ,

then prove that  $|f(z)| \leq M$  for every  $z \in \Omega$ . If, moreover,  $f$  is non-constant in  $\Omega$ , prove that  $|f(z)| < M$  for every  $z \in \Omega$ .

## 5.7 Taylor series and Laurent series.

**Proposition 5.2.** Let  $f$  be holomorphic in the open set  $\Omega$ ,  $z_0 \in \Omega$  and let  $D_{z_0}(R)$  be the largest disc with center  $z_0$  which is contained in  $\Omega$ . Then there is a unique power series  $\sum_{n=0}^{+\infty} a_n(z - z_0)^n$  so that

$$f(z) = \sum_{n=0}^{+\infty} a_n(z - z_0)^n \quad \text{for every } z \in D_{z_0}(R).$$

The coefficients are given by  $a_n = \frac{f^{(n)}(z_0)}{n!} = \frac{1}{2\pi i} \oint_{C_{z_0}(r)} \frac{f(\zeta)}{(\zeta - z_0)^{n+1}} d\zeta$  for  $0 < r < R$ .

*Proof.* We take  $z \in D_{z_0}(R)$ , and then  $|z - z_0| < R$ . If  $|z - z_0| < r < R$ , then  $z \in D_{z_0}(r)$  and, according to the formula of Cauchy, we have

$$f(z) = \frac{1}{2\pi i} \oint_{C_{z_0}(r)} \frac{f(\zeta)}{\zeta - z} d\zeta. \quad (5.6)$$

Now for every  $\zeta \in C_{z_0}(r)$  we have  $|\frac{z - z_0}{\zeta - z_0}| = \frac{|z - z_0|}{r} < 1$  and hence

$$\frac{1}{\zeta - z} = \frac{1}{(\zeta - z_0) - (z - z_0)} = \frac{1}{\zeta - z_0} \frac{1}{1 - \frac{z - z_0}{\zeta - z_0}} = \frac{1}{\zeta - z_0} \sum_{n=0}^{+\infty} \left(\frac{z - z_0}{\zeta - z_0}\right)^n.$$

The test of Weierstrass implies that  $\sum_{n=0}^{+\infty} \left(\frac{z - z_0}{\zeta - z_0}\right)^n$  converges, as a series of functions of  $\zeta$ , uniformly in  $C_{z_0}(r)$ . Indeed,  $|\frac{z - z_0}{\zeta - z_0}|^n = \left(\frac{|z - z_0|}{r}\right)^n$  for every  $\zeta \in C_{z_0}(r)$  and  $\sum_{n=0}^{+\infty} \left(\frac{|z - z_0|}{r}\right)^n < +\infty$ . So from (5.6) we have that

$$f(z) = \sum_{n=0}^{+\infty} \frac{1}{2\pi i} \oint_{C_{z_0}(r)} \frac{f(\zeta)}{(\zeta - z_0)^{n+1}} d\zeta (z - z_0)^n. \quad (5.7)$$

Now, we observe that the radius  $r$  has been chosen to satisfy the inequality  $|z - z_0| < r < R$  and hence the integrals  $\frac{1}{2\pi i} \oint_{C_{z_0}(r)} \frac{f(\zeta)}{(\zeta - z_0)^{n+1}} d\zeta$  depend *a priori* on  $z$ . But there are two reasons that these integrals actually do not depend on the value of  $r$  in the interval  $(0, R)$  and hence on  $z$ . The first reason is that from the formulas of Cauchy we get  $\frac{1}{2\pi i} \oint_{C_{z_0}(r)} \frac{f(\zeta)}{(\zeta - z_0)^{n+1}} d\zeta = \frac{f^{(n)}(z_0)}{n!}$  when  $0 < r < R$ . The second reason is that  $\frac{f(\zeta)}{(\zeta - z_0)^{n+1}}$  is holomorphic in  $D_{z_0}(R) \setminus \{z_0\}$ , and because of corollary 5.1, we have  $\frac{1}{2\pi i} \oint_{C_{z_0}(r_1)} \frac{f(\zeta)}{(\zeta - z_0)^{n+1}} d\zeta = \frac{1}{2\pi i} \oint_{C_{z_0}(r_2)} \frac{f(\zeta)}{(\zeta - z_0)^{n+1}} d\zeta$  when  $0 < r_1 < r_2 < R$ . We conclude from (5.7) that  $f(z) = \sum_{n=0}^{+\infty} a_n(z - z_0)^n$  for every  $z \in D_{z_0}(R)$ , where  $a_n = \frac{f^{(n)}(z_0)}{n!} = \frac{1}{2\pi i} \oint_{C_{z_0}(r)} \frac{f(\zeta)}{(\zeta - z_0)^{n+1}} d\zeta$  for  $0 < r < R$ .

Regarding uniqueness, assume that  $f(z) = \sum_{n=0}^{+\infty} b_n(z - z_0)^n$  for every  $z \in D_{z_0}(R)$ . Then, if  $0 < r < R$ , the series  $\sum_{n=0}^{+\infty} b_n(z - z_0)^n$  converges uniformly in  $C_{z_0}(r)$  and we get

$$\begin{aligned} 2\pi i a_k &= \oint_{C_{z_0}(r)} \frac{f(\zeta)}{(\zeta - z_0)^{k+1}} d\zeta = \oint_{C_{z_0}(r)} \frac{1}{(\zeta - z_0)^{k+1}} \sum_{n=0}^{+\infty} b_n(\zeta - z_0)^n d\zeta \\ &= \sum_{n=0}^{+\infty} b_n \oint_{C_{z_0}(r)} (\zeta - z_0)^{n-k-1} d\zeta = 2\pi i b_k. \end{aligned}$$

The last equality uses the calculation in example 4.2.2. Finally,  $b_k = a_k$  for every  $k$ .  $\square$

The power series  $\sum_{n=0}^{+\infty} a_n(z - z_0)^n$  with  $a_n = \frac{f^{(n)}(z_0)}{n!} = \frac{1}{2\pi i} \oint_{C_{z_0}(r)} \frac{f(\zeta)}{(\zeta - z_0)^{n+1}} d\zeta$  for  $0 < r < R$  is called **Taylor series** of  $f$  in the disc  $D_{z_0}(R)$ , the largest open disc with center  $z_0$  which is contained in the domain of holomorphy of  $f$ .

**Example 5.7.1.** The function  $f(z) = \frac{1}{1-z}$  is holomorphic in  $\mathbb{C} \setminus \{1\}$  and the largest open disc with center 0 which is contained in  $\mathbb{C} \setminus \{1\}$  is  $D_0(1)$ . To find the Taylor series of  $f$  in  $D_0(1)$  we calculate the derivatives  $f^{(n)}(z) = \frac{n!}{(1-z)^{n+1}}$  for every  $n \geq 0$ . Thus,  $a_n = \frac{f^{(n)}(0)}{n!} = 1$  for every  $n \geq 0$  and the Taylor series of  $f$  is  $\sum_{n=0}^{+\infty} z^n$ . I.e.  $\frac{1}{1-z} = \sum_{n=0}^{+\infty} z^n$  for every  $z \in D_0(1)$ . Of course, this is already known.

**Example 5.7.2.** The function  $f(z) = \frac{1}{1+z^2} = \frac{1}{(z+i)(z-i)}$  is holomorphic in the open set  $\mathbb{C} \setminus \{i, -i\}$  and the largest open disc with center 0 which is contained in  $\mathbb{C} \setminus \{i, -i\}$  is  $D_0(1)$ . To find the Taylor series of  $f$  in  $D_0(1)$  we calculate the derivatives of  $f$ . We write  $f(z) = -\frac{1}{2i} \left( \frac{1}{i-z} + \frac{1}{i+z} \right)$  and get  $f^{(n)}(z) = -\frac{1}{2i} \left( \frac{n!}{(i-z)^{n+1}} + (-1)^n \frac{n!}{(i+z)^{n+1}} \right)$  for every  $n \geq 0$ . Hence  $a_n = \frac{f^{(n)}(0)}{n!} = \frac{1+(-1)^n}{2i^n}$  for every  $n \geq 0$ . If  $n$  is odd, then  $a_n = 0$ . If  $n$  is even, then  $a_n = \frac{1}{i^n} = (-1)^{\frac{n}{2}}$  and the Taylor series of  $f$  is  $\sum_{k=0}^{+\infty} (-1)^k z^{2k}$ . I.e.  $\frac{1}{1+z^2} = \sum_{k=0}^{+\infty} (-1)^k z^{2k}$  for every  $z \in D_0(1)$ .

We may find the same formula if we use the Taylor series of  $\frac{1}{1-z}$ , i.e.  $\frac{1}{1-z} = \sum_{n=0}^{+\infty} z^n$ . We replace  $z$  with  $-z^2$  and find  $\frac{1}{1+z^2} = \sum_{n=0}^{+\infty} (-1)^n z^{2n}$ . From the moment that we have found *some* power series which coincides with our function in  $D_0(1)$ , then, because of uniqueness, *this* is the Taylor series of our function.

**Example 5.7.3.** The exponential function  $f(z) = e^z$  is holomorphic in  $\mathbb{C}$  and the largest open disc with center 0 which is contained in  $\mathbb{C}$  is  $D_0(+\infty) = \mathbb{C}$ . The derivatives of  $f$  are  $f^{(n)}(z) = e^z$  for every  $n \geq 0$  and the coefficients of the Taylor series of  $f$  are  $a_n = \frac{f^{(n)}(0)}{n!} = \frac{1}{n!}$  for every  $n \geq 0$ . Thus, the Taylor series of  $f$  is  $\sum_{n=0}^{+\infty} \frac{1}{n!} z^n$  and we have

$$e^z = \sum_{n=0}^{+\infty} \frac{1}{n!} z^n \quad \text{for every } z.$$

We have proven this identity differently in example 4.6.3.

**Example 5.7.4.** The function  $f(z) = \cos z$ , defined in exercise 4.2.6, is holomorphic in  $\mathbb{C}$  and the largest open disc with center 0 which is contained in  $\mathbb{C}$  is  $D_0(+\infty) = \mathbb{C}$ . The derivatives of  $f$  are  $f^{(n)}(z) = (-1)^{\frac{n}{2}} \cos z$  for even  $n$  and  $f^{(n)}(z) = (-1)^{\frac{n+1}{2}} \sin z$  for odd  $n$ . Therefore, the coefficients of the Taylor series are  $a_n = \frac{f^{(n)}(0)}{n!} = \frac{(-1)^{\frac{n}{2}}}{n!}$  for even  $n$  and  $a_n = \frac{f^{(n)}(0)}{n!} = 0$  for odd  $n$ . Thus, the Taylor series of  $f$  is  $\sum_{k=0}^{+\infty} \frac{(-1)^k}{(2k)!} z^{2k}$  and we have

$$\cos z = \sum_{k=0}^{+\infty} \frac{(-1)^k}{(2k)!} z^{2k} \quad \text{for every } z.$$

In the same manner we can prove that

$$\sin z = \sum_{k=1}^{+\infty} \frac{(-1)^{k-1}}{(2k-1)!} z^{2k-1} \quad \text{for every } z.$$

Another way to find the Taylor series of  $\cos z$  and  $\sin z$  is through the definitions of the two functions and the Taylor series of  $e^z$ . For instance:

$$\begin{aligned}\cos z &= \frac{e^{iz} + e^{-iz}}{2} = \frac{1}{2} \sum_{n=0}^{+\infty} \frac{1}{n!} (iz)^n + \frac{1}{2} \sum_{n=0}^{+\infty} \frac{1}{n!} (-iz)^n = \sum_{n=0}^{+\infty} \frac{i^n(1+(-1)^n)}{2n!} z^n \\ &= \sum_{k=0}^{+\infty} \frac{i^{2k}}{(2k)!} z^{2k} = \sum_{k=0}^{+\infty} \frac{(-1)^k}{(2k)!} z^{2k}.\end{aligned}$$

The power series we found coincides with the function  $\cos z$  in the largest open disc with center 0 which is contained in the domain of holomorphy of  $\cos z$  and, because of uniqueness, this is the Taylor series of  $\cos z$ .

**Example 5.7.5.** The function  $f(z) = -\text{Log}(1-z)$  is defined and holomorphic in  $\mathbb{C} \setminus [1, +\infty)$ . The largest disc with center 0 in  $\mathbb{C} \setminus [1, +\infty)$  is  $\mathbb{D}$ . The derivatives of  $f$  are  $f^{(n)}(z) = \frac{(n-1)!}{(1-z)^n}$  for every  $n \geq 1$ . Thus,  $a_0 = 0$  and  $a_n = \frac{f^{(n)}(0)}{n!} = \frac{1}{n}$  for every  $n \geq 1$  and the Taylor series of  $f$  is  $\sum_{n=1}^{+\infty} \frac{z^n}{n}$ . I.e.

$$-\text{Log}(1-z) = \sum_{n=1}^{+\infty} \frac{z^n}{n} \quad \text{for every } z \in \mathbb{D}.$$

**Proposition 5.3.** Let  $f$  be holomorphic in the open set  $\Omega$  and let  $D_{z_0}(R_1, R_2)$  be a largest open ring with center  $z_0$  which is contained in  $\Omega$ . Then there is a unique power series  $\sum_{-\infty}^{+\infty} a_n(z-z_0)^n$  so that

$$f(z) = \sum_{-\infty}^{+\infty} a_n(z-z_0)^n \quad \text{for every } z \in D_{z_0}(R_1, R_2).$$

The coefficients are given by  $a_n = \frac{1}{2\pi i} \oint_{C_{z_0}(r)} \frac{f(\zeta)}{(\zeta-z_0)^{n+1}} d\zeta$  for  $R_1 < r < R_2$ .

*Proof.* We take  $z \in D_{z_0}(R_1, R_2)$ , and then  $R_1 < |z-z_0| < R_2$ . We choose any  $r_1, r_2$  so that  $R_1 < r_1 < |z-z_0| < r_2 < R_2$ . Then  $z \in D_{z_0}(r_1, r_2)$  and

$$f(z) = \frac{1}{2\pi i} \oint_{C_{z_0}(r_2)} \frac{f(\zeta)}{\zeta-z} d\zeta - \frac{1}{2\pi i} \oint_{C_{z_0}(r_1)} \frac{f(\zeta)}{\zeta-z} d\zeta. \quad (5.8)$$

To prove (5.8), we consider an open disc  $D_z(r)$  with  $r < \min\{r_2 - |z-z_0|, |z-z_0| - r_1\}$ . Then  $\overline{D_z(r)} \subseteq D_{z_0}(r_1, r_2)$  and we apply corollary 5.1 to  $\frac{f(\zeta)}{\zeta-z}$ , which is a holomorphic function of  $\zeta$  in  $D_{z_0}(R_1, R_2) \setminus \{z\}$ . We get  $\oint_{C_{z_0}(r_2)} \frac{f(\zeta)}{\zeta-z} dz - \oint_{C_{z_0}(r_1)} \frac{f(\zeta)}{\zeta-z} dz = \oint_{C_z(r)} \frac{f(\zeta)}{\zeta-z} d\zeta$ . Now as in the proof of Cauchy's formula for circles, we have  $\lim_{r \rightarrow 0} \oint_{C_z(r)} \frac{f(\zeta)}{\zeta-z} d\zeta = 2\pi i f(z)$  and the proof of (5.8) is complete.

For every  $\zeta \in C_{z_0}(r_2)$  we have

$$\frac{1}{\zeta-z} = \frac{1}{(\zeta-z_0)-(z-z_0)} = \frac{1}{\zeta-z_0} \frac{1}{1-\frac{z-z_0}{\zeta-z_0}} = \frac{1}{\zeta-z_0} \sum_{n=0}^{+\infty} \left(\frac{z-z_0}{\zeta-z_0}\right)^n,$$

because  $|\frac{z-z_0}{\zeta-z_0}| = \frac{|z-z_0|}{r_2} < 1$ . Similarly, for every  $\zeta \in C_{z_0}(r_1)$  we have

$$\frac{1}{\zeta-z} = \frac{1}{(\zeta-z_0)-(z-z_0)} = -\frac{1}{z-z_0} \frac{1}{1-\frac{\zeta-z_0}{z-z_0}} = -\frac{1}{z-z_0} \sum_{n=0}^{+\infty} \left(\frac{\zeta-z_0}{z-z_0}\right)^n$$

because  $|\frac{\zeta-z_0}{z-z_0}| = \frac{r_1}{|z-z_0|} < 1$ . Exactly as in the proof of proposition 5.2, we see that these two series of functions converge uniformly and (5.8) implies

$$\begin{aligned}f(z) &= \sum_{n=0}^{+\infty} \frac{1}{2\pi i} \oint_{C_{z_0}(r_2)} \frac{f(\zeta)}{(\zeta-z_0)^{n+1}} d\zeta (z-z_0)^n \\ &\quad + \sum_{n=0}^{+\infty} \frac{1}{2\pi i} \oint_{C_{z_0}(r_1)} f(\zeta)(\zeta-z_0)^n d\zeta \frac{1}{(z-z_0)^{n+1}}.\end{aligned}$$

In the last series we change  $n+1$  to  $-n$  and get

$$\begin{aligned}f(z) &= \sum_{n=0}^{+\infty} \frac{1}{2\pi i} \oint_{C_{z_0}(r_2)} \frac{f(\zeta)}{(\zeta-z_0)^{n+1}} d\zeta (z-z_0)^n \\ &\quad + \sum_{-\infty}^{n=-1} \frac{1}{2\pi i} \oint_{C_{z_0}(r_1)} \frac{f(\zeta)}{(\zeta-z_0)^{n+1}} d\zeta (z-z_0)^n.\end{aligned} \quad (5.9)$$

Now,  $\frac{f(\zeta)}{(\zeta-z_0)^{n+1}}$  is holomorphic in  $D_{z_0}(R_1, R_2)$  and another application of corollary 5.1 implies that  $\oint_{C_{z_0}(r_1)} \frac{f(\zeta)}{(\zeta-z_0)^{n+1}} d\zeta = \oint_{C_{z_0}(r_2)} \frac{f(\zeta)}{(\zeta-z_0)^{n+1}} d\zeta$  for  $R_1 < r_1 < r_2 < R_2$ . Therefore the coefficients of both series in (5.9) do not depend on the values of  $r_1, r_2$ , and we replace both radii with any  $r$  with  $R_1 < r < R_2$ . We conclude that  $f(z) = \sum_{-\infty}^{+\infty} a_n(z-z_0)^n$  for every  $z \in D_{z_0}(R_1, R_2)$ , where  $a_n = \frac{1}{2\pi i} \oint_{C_{z_0}(r)} \frac{f(\zeta)}{(\zeta-z_0)^{n+1}} d\zeta$  for  $R_1 < r < R_2$ .

Regarding uniqueness, assume that  $f(z) = \sum_{-\infty}^{+\infty} b_n(z-z_0)^n$  for every  $z \in D_{z_0}(R_1, R_2)$ . We take any  $r$  with  $R_1 < r < R_2$ , and then  $\sum_{-\infty}^{+\infty} b_n(z-z_0)^n$  converges uniformly in  $C_{z_0}(r)$ . Then

$$\begin{aligned} 2\pi i a_k &= \oint_{C_{z_0}(r)} \frac{f(\zeta)}{(\zeta-z_0)^{k+1}} d\zeta = \oint_{C_{z_0}(r)} \frac{1}{(\zeta-z_0)^{k+1}} \sum_{-\infty}^{+\infty} b_n(\zeta-z_0)^n d\zeta \\ &= \sum_{-\infty}^{+\infty} b_n \oint_{C_{z_0}(r)} (\zeta-z_0)^{n-k-1} d\zeta = 2\pi i b_k \end{aligned}$$

and we get that  $b_k = a_k$  for every  $k$ .  $\square$

The power series  $\sum_{-\infty}^{+\infty} a_n(z-z_0)^n$  with  $a_n = \frac{1}{2\pi i} \oint_{C_{z_0}(r)} \frac{f(\zeta)}{(\zeta-z_0)^{n+1}} d\zeta$  for  $R_1 < r < R_2$  is called **Laurent series** of  $f$  in the ring  $D_{z_0}(R_1, R_2)$ , a largest open ring with center  $z_0$  which is contained in the domain of holomorphy of  $f$ .

**Example 5.7.6.** The function  $f(z) = \frac{1}{z}$  is holomorphic in  $\mathbb{C} \setminus \{0\}$ . The ring  $D_0(0, +\infty) = \mathbb{C} \setminus \{0\}$  is the largest open ring with center 0 which is contained in  $\mathbb{C} \setminus \{0\}$ . To find the Laurent series of  $f$  in  $D_0(0, +\infty)$  we evaluate the coefficients  $a_n$ . We take any  $r$  with  $0 < r < +\infty$ , and then we have  $a_n = \frac{1}{2\pi i} \oint_{C_0(r)} \frac{1/\zeta}{\zeta^{n+1}} d\zeta = \frac{1}{2\pi i} \oint_{C_0(r)} \frac{1}{\zeta^{n+2}} d\zeta$  for every  $n$ . If  $n \neq -1$ , then  $a_n = 0$  and, if  $n = -1$ , then  $a_{-1} = 1$ . Therefore, the Laurent series of  $f$  in  $D_0(0, +\infty)$  is  $\sum_{-\infty}^{+\infty} a_n z^n = z^{-1}$  and hence we have the obvious identity  $\frac{1}{z} = z^{-1}$  for every  $z \in D_0(0, +\infty)$ .

In the following examples we shall use the uniqueness of the Laurent series to find the Laurent series of certain functions without evaluating integrals: we find in an indirect way a power series which coincides with the function in a specific ring and then, because of uniqueness, this is the Laurent series of the function in the ring.

**Example 5.7.7.** The function  $f(z) = \frac{1}{1-z}$  is holomorphic in the open set  $\mathbb{C} \setminus \{1\}$ . We have seen that the largest open disc with center 0 which is contained in  $\mathbb{C} \setminus \{1\}$  is  $D_0(1)$  and that the Taylor series of  $f$  in this disc is  $\sum_{n=0}^{+\infty} z^n$ .

Another largest open ring with center 0 which is contained in  $\mathbb{C} \setminus \{1\}$  is  $D_0(1, +\infty)$ . To find the Laurent series of  $f$  in this ring, we may evaluate the coefficients  $a_n$  using their formulas with the integrals. But we can do something simpler. If  $z \in D_0(1, +\infty)$ , then  $|\frac{1}{z}| < 1$  and hence

$$\frac{1}{1-z} = -\frac{1}{z} \frac{1}{1-\frac{1}{z}} = -\frac{1}{z} \sum_{n=0}^{+\infty} \left(\frac{1}{z}\right)^n = -\sum_{-\infty}^{n=-1} z^n.$$

Because of uniqueness, the Laurent series of  $f$  in  $D_0(1, +\infty)$  is  $-\sum_{-\infty}^{n=-1} z^n$ .

**Example 5.7.8.** The function  $f(z) = \frac{1}{(z-1)(z-2)}$  is holomorphic in  $\mathbb{C} \setminus \{1, 2\}$ . There is a largest open disc and two largest open rings with center 0 which are contained in  $\mathbb{C} \setminus \{1, 2\}$ : the disc  $D_0(1)$  and the rings  $D_0(1, 2)$  and  $D_0(2, +\infty)$ . To find the corresponding Taylor and Laurent series we write  $f$  as a sum of simple fractions:  $f(z) = \frac{1}{z-2} - \frac{1}{z-1}$ .

If  $z \in D_0(1)$ , then  $|z| < 1$  and  $|\frac{z}{2}| < 1$ , and hence

$$f(z) = -\frac{1}{2} \frac{1}{1-\frac{z}{2}} + \frac{1}{1-z} = -\frac{1}{2} \sum_{n=0}^{+\infty} \left(\frac{z}{2}\right)^n + \sum_{n=0}^{+\infty} z^n = \sum_{n=0}^{+\infty} \left(1 - \frac{1}{2^{n+1}}\right) z^n.$$

Therefore, the Taylor series of  $f$  in  $D_0(1)$  is  $\sum_{n=0}^{+\infty} \left(1 - \frac{1}{2^{n+1}}\right) z^n$ .

If  $z \in D_0(1, 2)$ , then  $|\frac{1}{z}| < 1$  and  $|\frac{z}{2}| < 1$ , and hence

$$f(z) = -\frac{1}{2} \frac{1}{1-\frac{z}{2}} - \frac{1}{z} \frac{1}{1-\frac{1}{z}} = -\frac{1}{2} \sum_{n=0}^{+\infty} \left(\frac{z}{2}\right)^n - \frac{1}{z} \sum_{n=0}^{+\infty} \left(\frac{1}{z}\right)^n = -\sum_{-\infty}^{n=-1} z^n - \sum_{n=0}^{+\infty} \frac{1}{2^{n+1}} z^n.$$

Therefore, the Laurent series of  $f$  in  $D_0(1, 2)$  is  $-\sum_{n=-\infty}^{-1} z^n - \sum_{n=0}^{+\infty} \frac{1}{2^{n+1}} z^n$ .  
 If  $z \in D_0(2, +\infty)$ , then  $|\frac{1}{z}| < 1$  and  $|\frac{2}{z}| < 1$ , and hence

$$f(z) = \frac{1}{z} \frac{1}{1-\frac{2}{z}} - \frac{1}{z} \frac{1}{1-\frac{1}{z}} = \frac{1}{z} \sum_{n=0}^{+\infty} \left(\frac{2}{z}\right)^n - \frac{1}{z} \sum_{n=0}^{+\infty} \left(\frac{1}{z}\right)^n = \sum_{n=-\infty}^{-2} \left(\frac{1}{2^{n+1}} - 1\right) z^n.$$

Therefore, the Laurent series of  $f$  in  $D_0(2, +\infty)$  is  $\sum_{n=-\infty}^{-2} \left(\frac{1}{2^{n+1}} - 1\right) z^n$ .

**Example 5.7.9.** The function  $f(z) = e^{\frac{1}{z}}$  is holomorphic in  $\mathbb{C} \setminus \{0\}$ . Then  $D_0(0, +\infty) = \mathbb{C} \setminus \{0\}$  is the only largest open ring with center 0 which is contained in  $\mathbb{C} \setminus \{0\}$ . We find the Laurent series of  $f$  in  $D_0(0, +\infty)$  using the Taylor series of  $e^z$  in  $\mathbb{C}$ . In the identity  $e^z = \sum_{n=0}^{+\infty} \frac{1}{n!} z^n$  we replace  $z$  with  $\frac{1}{z}$  and we find  $e^{\frac{1}{z}} = \sum_{n=-\infty}^{-1} \frac{1}{(-n)!} z^n + 1$  for every  $z \neq 0$ . Therefore, the Laurent series of  $f$  in  $D_0(0, +\infty)$  is  $\sum_{n=-\infty}^{-1} \frac{1}{(-n)!} z^n + 1$ .

### Exercises.

**5.7.1.** Let  $0 < |a| < |b|$ . Find the three Laurent series with center 0, the two Laurent series with center  $a$  and the two Laurent series with center  $b$  of the function  $\frac{z}{(z-a)(z-b)}$ .

**5.7.2.** Find the Taylor series of  $\frac{1}{1+z^2}$  with center any  $a \in \mathbb{R}$ .

**5.7.3.** Find the Taylor series with center 1 of the holomorphic branch of  $z^{\frac{1}{2}}$  with value 1 at 1.

**5.7.4.** Let  $f$  be holomorphic in  $D_{z_0}(R)$  and let  $\sum_{n=0}^{+\infty} a_n(z-z_0)^n$  be the Taylor series of  $f$ .

(i) Prove that, if  $0 \leq r < R$ , then  $\frac{1}{2\pi} \int_0^{2\pi} |f(z_0 + re^{it})|^2 dt = \sum_{n=0}^{+\infty} |a_n|^2 r^{2n}$ .

(ii) If  $|f(z)| \leq M$  for every  $z \in D_{z_0}(R)$ , prove that  $\sum_{n=0}^{+\infty} |a_n|^2 R^{2n} \leq M^2$ .

(iii) If  $g$  is also holomorphic in  $D_{z_0}(R)$  with Taylor series  $\sum_{n=0}^{+\infty} b_n(z-z_0)^n$ , prove that, if  $0 \leq r < R$ , then  $\frac{1}{2\pi} \int_0^{2\pi} f(z_0 + re^{it}) \overline{g(z_0 + re^{it})} dt = \sum_{n=0}^{+\infty} a_n \overline{b_n} r^{2n}$ .

**5.7.5.** Let  $f$  be holomorphic in  $D_{z_0}(R_1, R_2)$ . Prove that there are functions  $f_1, f_2$  so that  $f_2$  is holomorphic in  $D_{z_0}(R_2)$  and  $f_1$  is holomorphic in  $D_{z_0}(R_1, +\infty) \cup \{\infty\}$  and so that  $f = f_1 + f_2$  in  $D_{z_0}(R_1, R_2)$ . Prove that, if  $f$  is bounded in  $D_{z_0}(R_1, R_2)$ , then  $f_1, f_2$  are bounded in  $D_{z_0}(R_1, R_2)$ .

**5.7.6.** Let  $f$  be holomorphic in  $D_0(R, +\infty)$ . Prove that  $f$  is holomorphic also at  $\infty$  if and only if the Laurent series of  $f$  in  $D_0(R, +\infty)$  is of the form  $\sum_{n=-\infty}^{-1} a_n z^n + a_0$ . Observe that  $f(\infty) = a_0$ .

**5.7.7.** Prove that  $\frac{1}{\cos z} = 1 + \sum_{k=1}^{+\infty} \frac{E_{2k}}{(2k)!} z^{2k}$  for  $|z| < \frac{\pi}{2}$ , where the numbers  $E_{2k}$  satisfy the recursive relations  $E_{2n} - \binom{2n}{2n-2} E_{2n-2} + \binom{2n}{2n-4} E_{2n-4} - \dots + (-1)^{n-1} \binom{2n}{2} E_2 + (-1)^n = 0$ . Evaluate  $E_2, E_4, E_6, E_8$ . The numbers  $E_{2k}$  are called Euler constants.

**5.7.8.** Let  $f$  be holomorphic in the horizontal zone  $\Omega = \{x + iy \mid A < y < B\}$  and periodic with period 1, i.e.  $f(z+1) = f(z)$  for every  $z \in \Omega$ .

(i) Prove that there are  $c_n$  so that  $f(z) = \sum_{n=-\infty}^{+\infty} c_n e^{2\pi i n z}$  for every  $z \in \Omega$  and find formulas for the coefficients  $c_n$ .

(ii) Prove that the series in (i) converges uniformly in every smaller zone  $\{x + iy \mid a < y < b\}$  with  $A < a < b < B$ .

**5.7.9.** (i) Prove that  $e^{\frac{w}{2}(z-\frac{1}{z})} = b_0(w) + \sum_{n=1}^{+\infty} b_n(w) \left(z^n + \frac{(-1)^n}{z^n}\right)$  for every  $z \neq 0$ , where  $b_n(w) = \frac{1}{\pi} \int_0^\pi \cos(nt - w \sin t) dt$  for  $n \in \mathbb{N}_0$ .

(ii) If  $m, n \in \mathbb{N}_0$ , prove that  $\frac{1}{2\pi i} \int_{C_0(1)} \frac{(z^2 \pm 1)^m}{z^{m+n+1}} dz = \begin{cases} \frac{(\pm 1)^p (n+2p)!}{p!(n+p)!} & \text{if } m = n + 2p, p \in \mathbb{N}_0 \\ 0, & \text{otherwise} \end{cases}$

(iii) The function  $b_n(w)$  is called Bessel function of the first kind. Find the Taylor series of  $b_n(w)$  with center 0.

**5.7.10.** Let  $I$  be an open interval in  $\mathbb{R}$ . The function  $f : I \rightarrow \mathbb{C}$  is called **real analytic** in  $I$  if for every  $t_0 \in I$  there are  $\epsilon > 0$  and  $a_n \in \mathbb{C}$ ,  $n \in \mathbb{N}_0$ , so that  $(t_0 - \epsilon, t_0 + \epsilon) \subseteq I$  and  $f(t) = \sum_{n=0}^{+\infty} a_n (t - t_0)^n$  for every  $t \in (t_0 - \epsilon, t_0 + \epsilon)$ .

Prove that, if  $f$  is real analytic in  $I$ , then there is an open set  $\Omega \subseteq \mathbb{C}$  so that  $I \subseteq \Omega$  and so that  $f$  can be extended as a function  $f : \Omega \rightarrow \mathbb{C}$  holomorphic in  $\Omega$ .

## 5.8 Roots and the principle of identity.

Let  $f$  be holomorphic in the open set  $\Omega$  and  $z_0 \in \Omega$ . We consider the largest open disc  $D_{z_0}(R)$  which is contained in  $\Omega$  and the Taylor series of  $f$  in this disc. Then

$$f(z) = \sum_{n=0}^{+\infty} a_n(z-z_0)^n = a_0 + a_1(z-z_0) + a_2(z-z_0)^2 + \cdots \quad \text{for every } z \in D_{z_0}(R).$$

We assume that  $z_0$  is a root of  $f$  or, equivalently, that  $a_0 = 0$  and we distinguish between two cases.

*First case:  $a_n = 0$  for every  $n$ .*

Then, obviously,  $f(z) = 0$  for every  $z \in D_{z_0}(R)$ , i.e.  $f$  is identically 0 in  $D_{z_0}(R)$ . Because of the formulas for  $a_n$ , the condition  $a_n = 0$  for every  $n$  is equivalent to  $f^{(n)}(z_0) = 0$  for every  $n$ .

*Second case:  $a_n \neq 0$  for at least one  $n$ .*

We consider the smallest  $n \geq 1$  with  $a_n \neq 0$  and let this be  $N$ . I.e.  $a_0 = a_1 = \dots = a_{N-1} = 0$  and  $a_N \neq 0$ . This is equivalent to  $f(z_0) = f^{(1)}(z_0) = \dots = f^{(N-1)}(z_0) = 0$  and  $f^{(N)}(z_0) \neq 0$ . Then we have  $f(z) = (z-z_0)^N \sum_{n=0}^{+\infty} a_{N+n}(z-z_0)^n$  for every  $z \in D_{z_0}(R)$ . The power series  $\sum_{n=0}^{+\infty} a_{N+n}(z-z_0)^n = a_N + a_{N+1}(z-z_0) + a_{N+2}(z-z_0)^2 + \dots$  converges in the disc  $D_{z_0}(R)$  and defines a function  $g$  holomorphic in  $D_{z_0}(R)$ . Then

$$f(z) = (z-z_0)^N g(z) \quad \text{for every } z \in D_{z_0}(R),$$

and thus  $g(z) = \frac{f(z)}{(z-z_0)^N}$  for every  $z \in D_{z_0}(R) \setminus \{z_0\}$ . We observe that  $\frac{f(z)}{(z-z_0)^N}$  is a holomorphic function in  $\Omega \setminus \{z_0\}$  and not only in  $D_{z_0}(R) \setminus \{z_0\}$ . Therefore, we may consider  $g$  as defined in  $\Omega \setminus \{z_0\}$  with the same formula:  $g(z) = \frac{f(z)}{(z-z_0)^N}$ . We also recall that  $g$  is defined, through its power series, at  $z_0$  and it is holomorphic in  $D_{z_0}(R) \subseteq \Omega$ . In fact its value at  $z_0$  is  $g(z_0) = a_N = \frac{f^{(N)}(z_0)}{N!}$ . Thus, the formula of  $g$ , as a function holomorphic in  $\Omega$ , can be written:

$$g(z) = \begin{cases} \frac{f(z)}{(z-z_0)^N}, & \text{if } z \in \Omega \setminus \{z_0\} \\ a_N = \frac{f^{(N)}(z_0)}{N!}, & \text{if } z = z_0 \end{cases} \quad (5.10)$$

Since  $g(z_0) = a_N \neq 0$  and since  $g$  is continuous at  $z_0$ , there is  $r$  with  $0 < r \leq R$  so that  $g(z) \neq 0$  for every  $z \in D_{z_0}(r)$ , and hence  $f(z) \neq 0$  for every  $z \in D_{z_0}(r) \setminus \{z_0\}$ .

Let  $f$  be holomorphic in the open set  $\Omega$ ,  $z_0 \in \Omega$  and  $\sum_{n=0}^{+\infty} a_n(z-z_0)^n$  be the Taylor series of  $f$  at  $z_0$ . Then we have three cases. (i) If  $a_n = 0$  for every  $n$ , then we say that  $z_0$  is a root of  $f$  of **infinite multiplicity**. (ii) If  $a_0 = a_1 = \dots = a_{N-1} = 0$  and  $a_N \neq 0$  for some  $N \geq 1$ , then we say that  $z_0$  is a root of  $f$  of **multiplicity  $N$** . (iii) In case  $f(z_0) = a_0 \neq 0$  we say that  $z_0$  is a root of  $f$  of **multiplicity 0**.

We saw that, if  $z_0$  is a root of  $f$  of infinite multiplicity, then  $f$  is identically 0 in the largest disc with center  $z_0$  which is contained in the domain of holomorphy of  $f$ . If  $z_0$  is a root of  $f$  of finite multiplicity, then there is some disc  $D_{z_0}(r)$  which contains no other root of  $f$  besides  $z_0$  and hence we say that the root  $z_0$  is **isolated**. Moreover, if the multiplicity of  $z_0$  is  $N$ , then the function  $g(z) = \frac{f(z)}{(z-z_0)^N}$ , which is holomorphic in  $\Omega \setminus \{z_0\}$ , can be defined at  $z_0$  as  $g(z_0) = a_N = \frac{f^{(N)}(z_0)}{N!}$  and then it is holomorphic in  $\Omega$ . In other words, we can factorize  $(z-z_0)^N$  from  $f(z)$ , i.e. we can write  $f(z) = (z-z_0)^N g(z)$  with a function  $g$  holomorphic in  $\Omega$ . This is a striking generalization of the analogous factorization for polynomials: if  $z_0$  is a root of the polynomial  $p(z)$  of multiplicity  $N$ , then we can write  $p(z) = (z-z_0)^N q(z)$ , where  $q(z)$  is another polynomial.

**Example 5.8.1.** The function  $e^{z^3} - 1$  is holomorphic in  $\mathbb{C}$  and its Taylor series with center 0 is  $\sum_{n=1}^{+\infty} \frac{1}{n!} z^{3n}$ . Therefore,  $e^{z^3} - 1 = z^3 \sum_{n=1}^{+\infty} \frac{1}{n!} z^{3(n-1)} = z^3 \sum_{n=0}^{+\infty} \frac{1}{(n+1)!} z^{3n} = z^3 g(z)$  for every  $z$ , where  $g$  is the function defined by the power series  $\sum_{n=0}^{+\infty} \frac{1}{(n+1)!} z^{3n}$ . Now  $g$  is holomorphic in  $\mathbb{C}$  with  $g(0) = 1 \neq 0$ , hence 0 is a root of  $e^{z^3} - 1$  of multiplicity 3.

**Lemma 5.1.** *If  $f$  is holomorphic in the region  $\Omega$  and if  $z_0 \in \Omega$  is a root of  $f$  of infinite multiplicity, then  $f$  is identically 0 in  $\Omega$ .*

*Proof.*  $f$  is identically 0 in some disc with center  $z_0$ . We define

$$B = \{z \in \Omega \mid f \text{ is identically 0 in some disc with center } z\}$$

and the complementary set  $C = \Omega \setminus B$ . Obviously,  $B \cup C = \Omega$  and  $B \neq \emptyset$ , since  $z_0 \in B$ .

If  $z \in B$ , then  $f$  is identically 0 in some disc  $D_z(r)$ , and if we take any  $w \in D_z(r)$ , then  $f$  is identically 0 in some small disc  $D_w(r') \subseteq D_z(r)$ . Thus every  $w \in D_z(r)$  belongs to  $B$ , i.e.  $D_z(r) \subseteq B$  and  $z$  is not a limit point of  $C$ .

Now, let  $z \in C$ . Then  $f$  is identically 0 in no disc with center  $z$ , and hence  $z$  is not a root of infinite multiplicity of  $f$ . Therefore, there is a disc  $D_z(r)$  in which the only possible root of  $f$  is its center  $z$ . Then this disc contains no  $w \in B$  and  $z$  is not a limit point of  $B$ .

Thus, none of  $B, C$  contains a limit point of the other. Since  $B \neq \emptyset$ , we must have  $C = \emptyset$ , otherwise  $B, C$  would form a decomposition of  $\Omega$ . Hence  $\Omega = B$  and  $f$  is identically 0 in  $\Omega$ .  $\square$

**Principle of identity.** *If  $f$  is holomorphic in the region  $\Omega$  and if the roots of  $f$  have an accumulation point in  $\Omega$ , then  $f$  is identically 0 in  $\Omega$ .*

*Proof.* Suppose that there is a sequence  $(z_n)$  of roots of  $f$  so that  $z_n \rightarrow z$  with  $z \in \Omega$  and  $z_n \neq z$  for every  $n$ . Since  $f$  is continuous at  $z$  and  $z_n \rightarrow z$ , we have  $0 = f(z_n) \rightarrow f(z)$  and hence  $f(z) = 0$ . If  $z$  is a root of finite multiplicity of  $f$ , then there would be some disc  $D_z(r)$  in which the only root of  $f$  is its center  $z$ . This is wrong, since  $D_z(r)$  contains, after some index, all roots  $z_n$  and these are different from  $z$ . Therefore,  $z$  is a root of infinite multiplicity of  $f$ , and lemma 5.1 implies that  $f$  is identically 0 in  $\Omega$ .  $\square$

Lemma 5.1 and the principle of identity can be stated for a non-connected open set  $\Omega$ . Then the result of lemma 5.1 holds in the connected component of  $\Omega$  which contains the root of infinite multiplicity  $z_0$  and the result of the principle of identity holds in the connected component of  $\Omega$  which contains the accumulation point of the roots of  $f$ .

Instead of speaking only about the roots of  $f$ , i.e. the solutions of the equation  $f(z) = 0$ , we may state our results for the solutions of the equation  $f(z) = w$  for any fixed  $w$ . The results are the same as before. We just consider the function  $g(z) = f(z) - w$ , and then the solutions of  $f(z) = w$  are the same as the roots of  $g$ . For instance, if  $z_0$  is a solution of  $f(z) = w$  of infinite multiplicity, then  $f$  is constant  $w$  in some disc  $D_{z_0}(R)$  and, if  $z_0$  is a solution of  $f(z) = w$  of finite multiplicity  $N$ , then in some disc  $D_{z_0}(r)$  the function  $f$  takes the value  $w$  only at the center  $z_0$ . Then lemma 5.1 says that, if  $f$  is holomorphic in the region  $\Omega$  and  $z_0$  is a solution of  $f(z) = w$  of infinite multiplicity, then  $f$  is constant  $w$  in  $\Omega$ . And the principle of identity says that, if  $f$  is holomorphic in the region  $\Omega$  and the solutions of  $f(z) = w$  have an accumulation point in  $\Omega$ , then  $f$  is constant  $w$  in  $\Omega$ .

The principle of identity has another equivalent form.

**Principle of identity.** *If  $f$  is holomorphic in the region  $\Omega$  and if some compact  $K \subseteq \Omega$  contains infinitely many roots of  $f$ , then  $f$  is identically 0 in  $\Omega$ .*

*Proof.* Let us assume the previous principle of identity and let us suppose that some compact  $K \subseteq \Omega$  contains infinitely many roots of  $f$ . Then there is a sequence  $(z_n)$  of roots of  $f$  in  $K$  with distinct terms. Since  $K$  is compact, there is a subsequence  $(z_{n_k})$  so that  $z_{n_k} \rightarrow z$  for some  $z \in K$ . But then  $z \in \Omega$  is an accumulation point of roots of  $f$  and hence  $f$  is identically 0 in  $\Omega$ .

Conversely, let us assume the present form of the principle of identity and let us suppose that the roots of  $f$  have an accumulation point in  $\Omega$ . Then there is a sequence  $(z_n)$  of roots of  $f$  so that  $z_n \rightarrow z$  with  $z \in \Omega$  and  $z_n \neq z$  for every  $n$ . We take a compact disc  $\bar{D}_z(r) \subseteq \Omega$  and then this disc contains infinitely many of the roots  $z_n$ . Hence  $f$  is identically 0 in  $\Omega$ .  $\square$



**Example 5.8.2.** Assume that there is  $f$  holomorphic in  $\mathbb{C}$  so that  $f(\frac{1}{n}) = \frac{n}{n+1}$  for every  $n \in \mathbb{N}$ . We write  $f(\frac{1}{n}) = \frac{1}{1+\frac{1}{n}}$  and compare the functions  $f(z)$  and  $\frac{1}{1+z}$ . Both are holomorphic in  $\mathbb{C} \setminus \{-1\}$  and their difference  $f(z) - \frac{1}{1+z}$  has roots at the points  $\frac{1}{n}$  which have 0 as their accumulation point. Since  $0 \in \mathbb{C} \setminus \{-1\}$  and  $\mathbb{C} \setminus \{-1\}$  is connected, we have that  $f(z) - \frac{1}{1+z}$  is identically 0 in this set, i.e.  $f(z) = \frac{1}{1+z}$  for every  $z \neq -1$ . Since we assume that  $f$  is holomorphic at  $-1$ , we get  $\lim_{z \rightarrow -1} \frac{1}{1+z} = \lim_{z \rightarrow -1} f(z) = f(-1)$  and we arrive at a contradiction.

**Example 5.8.3.** Assume that there is some  $f$  holomorphic in  $\mathbb{C} \setminus \{0\}$  so that  $f(x) = \sqrt{x}$  for every  $x \in (0, +\infty)$  or even for every  $x$  in some subinterval  $(a, b)$  of  $(0, +\infty)$ . We consider the continuous branch  $g$  of  $z^{\frac{1}{2}}$  in the open set  $A = \mathbb{C} \setminus (-\infty, 0]$  which has value 1 at  $z = 1$ . The function  $g$  is given by

$$g(z) = \sqrt{r} e^{i\frac{\theta}{2}} \quad \text{for } z = re^{i\theta} \text{ with } r > 0 \text{ and } -\pi < \theta < \pi.$$

We see that  $f(x) = \sqrt{x} = g(x)$  for every  $x \in (a, b)$ . Hence  $f - g$  is holomorphic in the region  $A$  and has roots at all points of  $(a, b)$ . We conclude that  $f - g$  is identically 0 in  $A$ . I.e.

$$f(z) = \sqrt{r} e^{i\frac{\theta}{2}} \quad \text{for } z = re^{i\theta} \text{ with } r > 0 \text{ and } -\pi < \theta < \pi.$$

Since  $f$  is holomorphic in  $\mathbb{C} \setminus \{0\}$ , it is continuous at every point of  $(-\infty, 0)$ , e.g. at  $-1$ . We take points  $z = re^{i\theta}$  converging to  $-1$  from the upper halfplane. This means that  $r \rightarrow 1$  and  $\theta \rightarrow \pi^-$ . Then we have  $f(-1) = \lim_{r \rightarrow 1, \theta \rightarrow \pi^-} \sqrt{r} e^{i\frac{\theta}{2}} = e^{i\frac{\pi}{2}} = i$ . Now we take points  $z = re^{i\theta}$  converging to  $-1$  from the lower halfplane. This means that  $r \rightarrow 1$  and  $\theta \rightarrow -\pi^+$ . Then we have  $f(-1) = \lim_{r \rightarrow 1, \theta \rightarrow -\pi^+} \sqrt{r} e^{i\frac{\theta}{2}} = e^{-i\frac{\pi}{2}} = -i$ . We arrive at a contradiction.

### Exercises.

**5.8.1.** Let  $f$  be holomorphic in the disc  $D_{z_0}(R)$  and let  $z_0$  be a root of multiplicity  $N \geq 1$  of  $f$ . If  $F$  is a primitive of  $f$  in  $D_{z_0}(R)$  and  $F(z_0) = w_0$ , which is the multiplicity of  $z_0$  as a solution of  $F(z) = w_0$ ?

**5.8.2.** Is there any  $f$  holomorphic in  $\mathbb{C}$  which satisfies one of the following?

- (i)  $f(\frac{1}{n}) = (-1)^n$  for every  $n \in \mathbb{N}$ .
- (ii)  $f(\frac{1}{n}) = \frac{1+(-1)^n}{n}$  for every  $n \in \mathbb{N}$ .
- (iii)  $f(\frac{1}{2k}) = f(\frac{1}{2k+1}) = \frac{1}{k}$  for every  $k \in \mathbb{N}$ .

**5.8.3.** Is there any  $f$  holomorphic in  $\mathbb{C} \setminus \{0\}$  so that  $f(x) = |x|$  for every  $x \in \mathbb{R} \setminus \{0\}$ ?

**5.8.4.** Let  $f, g$  be holomorphic in the region  $\Omega$  and  $0 \in \Omega$ . If  $f, g$  have no root in  $\Omega$  and if  $f'(\frac{1}{n})/f(\frac{1}{n}) = g'(\frac{1}{n})/g(\frac{1}{n})$  for every  $n \in \mathbb{N}$ , what do you conclude about  $f, g$ ?

**5.8.5.** Let  $f, g$  be holomorphic in the region  $\Omega$ . If  $fg = 0$  in  $\Omega$ , prove that either  $f = 0$  in  $\Omega$  or  $g = 0$  in  $\Omega$ .

**5.8.6.** Let  $f, g$  be holomorphic in the region  $\Omega$ . If  $\bar{f}g$  is holomorphic in  $\Omega$ , prove that either  $g = 0$  in  $\Omega$  or  $f$  is constant in  $\Omega$ .

**5.8.7.** (i) Let the region  $\Omega$  be symmetric with respect to  $\mathbb{R}$ , i.e.  $\bar{z} \in \Omega$  for every  $z \in \Omega$ . If  $\Omega \neq \emptyset$ , prove that  $\Omega \cap \mathbb{R} \neq \emptyset$ . Let also  $f$  be holomorphic in  $\Omega$  and assume that  $f(z) \in \mathbb{R}$  for every  $z \in \Omega \cap \mathbb{R}$ . Prove that  $f(\bar{z}) = \overline{f(z)}$  for every  $z \in \Omega$ .

(ii) Let the region  $\Omega \subseteq \mathbb{C} \setminus \{0\}$  be symmetric with respect to  $\mathbb{T}$ , i.e.  $\frac{1}{\bar{z}} \in \Omega$  for every  $z \in \Omega$ . If  $\Omega \neq \emptyset$ , prove that  $\Omega \cap \mathbb{T} \neq \emptyset$ . Let also  $f$  be holomorphic in  $\Omega$  and assume that  $f(z) \in \mathbb{T}$  for every  $z \in \Omega \cap \mathbb{T}$ . Prove that  $f(\frac{1}{\bar{z}}) = \frac{1}{f(z)}$  for every  $z \in \Omega$ .

(iii) Let  $f$  be holomorphic in  $\mathbb{C}$  and let  $f(z) \in \mathbb{T}$  for every  $z \in \mathbb{T}$ . Prove that there is  $c$  with  $|c| = 1$  and  $n \in \mathbb{N}_0$  so that  $f(z) = cz^n$  for every  $z$ .

**5.8.8.** (i) Let  $z_0 \in \mathbb{D}$  and  $T(z) = \frac{z-z_0}{1-\bar{z}_0z}$  for  $z \in \bar{\mathbb{D}}$ . Prove that  $T$  is holomorphic in  $\mathbb{D}$  and continuous in  $\bar{\mathbb{D}}$ . Also prove that  $T(z) \in \mathbb{D}$  for every  $z \in \mathbb{D}$  and that  $T(z) \in \mathbb{T}$  for every  $z \in \mathbb{T}$ .

(ii) Let  $z_1, \dots, z_n \in \mathbb{D}$  and  $|c| = 1$  and  $B(z) = c \prod_{k=1}^n \frac{z-z_k}{1-\bar{z}_kz}$  for  $z \in \bar{\mathbb{D}}$ . Prove that  $B$  is holomorphic in  $\mathbb{D}$  and continuous in  $\bar{\mathbb{D}}$ . Also prove that  $B(z) \in \mathbb{D}$  for every  $z \in \mathbb{D}$  and that  $B(z) \in \mathbb{T}$  for every  $z \in \mathbb{T}$ .

(iii) Prove the converse of (ii). I.e. let  $f$  be holomorphic in  $\mathbb{D}$  and continuous in  $\bar{\mathbb{D}}$  and let  $f(z) \in \mathbb{D}$  for every  $z \in \mathbb{D}$  and  $f(z) \in \mathbb{T}$  for every  $z \in \mathbb{T}$ . If  $f$  is non-constant, prove that there is  $n \in \mathbb{N}$  and  $z_1, \dots, z_n \in \mathbb{D}$  and  $c$  with  $|c| = 1$  so that  $f(z) = c \prod_{k=1}^n \frac{z-z_k}{1-\bar{z}_kz}$  for every  $z \in \bar{\mathbb{D}}$ .

**5.8.9.** Let  $f, g$  be holomorphic in  $\mathbb{C}$  and  $|f(z)| \leq |g(z)|$  for every  $z$ . Prove that there is  $\mu$  so that  $f(z) = \mu g(z)$  for every  $z$ .

**5.8.10.** Let  $f$  be holomorphic in  $\mathbb{D}$ . Prove that there is a sequence  $(z_n)$  in  $\mathbb{D}$  so that  $|z_n| \rightarrow 1$  and  $(f(z_n))$  is bounded.

**5.8.11.** Many of the results of this section hold also for the point  $\infty$ .

(i) Let  $\Omega \subseteq \mathbb{C}$  be an open set containing some ring  $D_0(R, +\infty)$  and let  $f$  be holomorphic in  $\Omega \cup \{\infty\}$ . Then, according to exercise 5.7.6, the Laurent series of  $f$  in  $D_0(R, +\infty)$  is of the form  $\sum_{n=-\infty}^{-1} a_n z^n + a_0$  and also  $f(\infty) = a_0$ .

If  $a_n = 0$  for every  $n \leq 0$ , we say that  $\infty$  is a root of  $f$  of multiplicity  $+\infty$ , and in this case prove that  $f$  is identically 0 in the connected component of  $\Omega$  which contains  $D_0(R, +\infty)$ .

If  $a_0 = a_{-1} = \dots = a_{-N+1} = 0$  and  $a_{-N} \neq 0$ , we say that  $\infty$  is a root of  $f$  of multiplicity  $N$ , and in this case prove that  $\infty$  is an isolated root of  $f$ , i.e. there is some  $r \geq R$  so that  $f$  has no root in  $D_0(r, +\infty)$ .

Of course, if  $a_0 \neq 0$ , we say that  $\infty$  is a root of  $f$  of multiplicity 0.

If  $\infty$  is an accumulation point of roots of  $f$ , prove that  $f$  is identically 0 in the connected component of  $\Omega$  which contains  $D_0(R, +\infty)$ .

Prove that  $\infty$  is a root of  $f$  of multiplicity  $N$  if and only if 0 is a root of  $g$  of multiplicity  $N$ , where  $g$  is defined by  $g(w) = f(\frac{1}{w})$ .

(ii) Let  $r = \frac{p}{q}$  be a rational function and let  $n$  be the degree of the polynomial  $p$  and  $m$  be the degree of the polynomial  $q$ . If  $n \leq m$ , prove that  $\infty$  is a root of  $r$  of multiplicity  $m - n$ .

## 5.9 Isolated singularities.

We say that  $z_0$  is an **isolated singularity** of  $f$  if there is some disc  $D_{z_0}(R)$  so that  $f$  is holomorphic in  $D_{z_0}(R) \setminus \{z_0\}$ . Then  $f$  has a Laurent series in  $D_{z_0}(0, R) = D_{z_0}(R) \setminus \{z_0\}$ . I.e.

$$f(z) = \sum_{n=-\infty}^{+\infty} a_n (z - z_0)^n \quad \text{for every } z \in D_{z_0}(R) \setminus \{z_0\}.$$

Now we have three cases. (i) If  $a_n = 0$  for every  $n < 0$ , then we say that  $z_0$  is a **removable singularity** of  $f$ . (ii) If  $a_n \neq 0$  for at least one  $n < 0$  and there are only finitely many  $n < 0$  such that  $a_n \neq 0$ , then we say that  $z_0$  is a **pole** of  $f$ . (iii) If  $a_n \neq 0$  for infinitely many  $n < 0$ , then we say that  $z_0$  is an **essential singularity** of  $f$ .

Let us start with the case of a removable singularity  $z_0$ . Then  $f(z) = \sum_{n=0}^{+\infty} a_n (z - z_0)^n$  for every  $z \in D_{z_0}(R) \setminus \{z_0\}$ . The power series  $\sum_{n=0}^{+\infty} a_n (z - z_0)^n$  converges at every  $z \in D_{z_0}(R)$  and defines a holomorphic function in  $D_{z_0}(R)$  with value  $a_0$  at  $z_0$ . The function  $f$  may not be defined at  $z_0$  or it may be defined at  $z_0$  with a value  $f(z_0)$  either equal to  $a_0$  or not equal to  $a_0$ . Now, in any case, we define (or redefine)  $f$  at  $z_0$  to be  $f(z_0) = a_0$ . Then we have  $f(z) = \sum_{n=0}^{+\infty} a_n (z - z_0)^n$  for every  $z \in D_{z_0}(R)$  and  $f$  becomes holomorphic in  $D_{z_0}(R)$ .

We summarize. *If  $z_0 \in \Omega$  is a removable singularity of  $f$ , then  $f$  can be defined (or redefined) appropriately at  $z_0$  so that it becomes holomorphic in a disc with center  $z_0$ . The Laurent series of  $f$  at  $z_0$  reduces to a power series of first type and this power series is the Taylor series of the (extended)  $f$  in a disc with center  $z_0$ .*

Here is a useful test to decide if an isolated singularity is removable without calculating the Laurent series of the function.

**Riemann's criterion.** *Let  $z_0$  be an isolated singularity of  $f$ . If  $\lim_{z \rightarrow z_0} (z - z_0)f(z) = 0$ , then  $z_0$  is a removable singularity of  $f$ .*

*Proof.* Let  $f(z) = \sum_{-\infty}^{+\infty} a_n(z - z_0)^n$  for every  $z \in D_{z_0}(R) \setminus \{z_0\}$ . We take any  $\epsilon > 0$  and then there is  $\delta > 0$  so that  $|z - z_0||f(z)| \leq \epsilon$  for every  $z \in D_{z_0}(R)$  with  $0 < |z - z_0| < \delta$ . Now, we consider any  $r$  with  $0 < r < \min\{\delta, R, 1\}$  and any  $n < 0$ . Then we have

$$|a_n| = \left| \frac{1}{2\pi i} \oint_{C_{z_0}(r)} \frac{f(\zeta)}{(\zeta - z_0)^{n+1}} d\zeta \right| \leq \frac{1}{2\pi} \frac{\epsilon}{r^{n+2}} 2\pi r = \epsilon r^{-n-1} = \epsilon r^{|n|-1} \leq \epsilon.$$

Since  $\epsilon > 0$  is arbitrary, we get  $a_n = 0$  for every  $n < 0$  and  $z_0$  is a removable singularity of  $f$ .  $\square$

In the case of an isolated singularity  $z_0$  for  $f$ , sometimes we know that the  $\lim_{z \rightarrow z_0} f(z)$  exists and it is finite or that  $f$  is bounded close to  $z_0$ . In both cases we have that  $\lim_{z \rightarrow z_0} (z - z_0)f(z) = 0$  is satisfied and we conclude that  $z_0$  is a removable singularity of  $f$ .

**Example 5.9.1.** The function  $f(z) = \frac{z^2 - 3z + 2}{z - 2}$  is holomorphic in  $\mathbb{C} \setminus \{2\}$ . Since  $\lim_{z \rightarrow 2} f(z) = 1$ , the point 2 is a removable singularity of  $f$ . If we define  $f(2) = 1$ , then  $f$ , now defined in  $\mathbb{C}$ , is holomorphic in  $\mathbb{C}$ . In fact, the extended  $f$  is the simple function  $z - 1$  in  $\mathbb{C}$ .

Now we consider the case of a pole  $z_0$  of  $f$ . Let  $\sum_{-\infty}^{+\infty} a_n(z - z_0)^n$  be the Laurent series of  $f$  in the ring  $D_{z_0}(R) \setminus \{z_0\}$  and then there is a largest  $m \geq 1$  so that  $a_{-m} \neq 0$ . Let  $N$  be this largest  $m$ . Then we say that  $z_0$  is a pole of  $f$  of **order  $N$**  or of **multiplicity  $N$**  and we have

$$f(z) = \frac{a_{-N}}{(z - z_0)^N} + \cdots + \frac{a_{-1}}{z - z_0} + \sum_{n=0}^{+\infty} a_n(z - z_0)^n \quad \text{for every } z \in D_{z_0}(R) \setminus \{z_0\}$$

with  $a_{-N} \neq 0$ . We may write this as  $f(z) = \frac{1}{(z - z_0)^N} \sum_{n=0}^{+\infty} a_{n-N}(z - z_0)^n$  for every  $z \in D_{z_0}(R) \setminus \{z_0\}$ . Since the power series  $\sum_{n=0}^{+\infty} a_{n-N}(z - z_0)^n$  converges in the disc  $D_{z_0}(R)$ , it defines a function  $g$  holomorphic in  $D_{z_0}(R)$  and we have

$$f(z) = \frac{g(z)}{(z - z_0)^N} \quad \text{for every } z \in D_{z_0}(R) \setminus \{z_0\}.$$

Observe that  $g(z_0) = a_{-N} \neq 0$ .

It is easy to prove the converse. Suppose there is a  $g$  holomorphic in some disc  $D_{z_0}(R)$  so that  $g(z_0) \neq 0$  and  $f(z) = \frac{g(z)}{(z - z_0)^N}$  for every  $z \in D_{z_0}(R) \setminus \{z_0\}$ . Let  $\sum_{n=0}^{+\infty} b_n(z - z_0)^n$  be the Taylor series of  $g$  and then we have  $f(z) = \frac{b_0}{(z - z_0)^N} + \cdots + \frac{b_{N-1}}{z - z_0} + \sum_{n=0}^{+\infty} b_{n+N}(z - z_0)^n$  for  $z \in D_{z_0}(R) \setminus \{z_0\}$ . The last power series is the Laurent series of  $f$  in  $D_{z_0}(R) \setminus \{z_0\}$  and since  $b_0 = g(z_0) \neq 0$ , we have that  $z_0$  is a pole of  $f$  of order  $N$ .

Here are some more comments. Since  $g(z_0) \neq 0$  and  $g$  is continuous at  $z_0$ , we have that  $g$  does not vanish at any point of some disc  $D_{z_0}(r)$  with  $0 < r \leq R$ . Then  $h(z) = \frac{1}{g(z)}$  is holomorphic in  $D_{z_0}(r)$  and  $\frac{1}{f(z)} = (z - z_0)^N h(z)$  for every  $z \in D_{z_0}(r) \setminus \{z_0\}$ . Therefore,  $z_0$  is a removable singularity of  $\frac{1}{f}$ . Moreover, if we define  $\frac{1}{f}$  to take the value 0 at  $z_0$ , then we have  $\frac{1}{f}(z) = (z - z_0)^N h(z)$  for every  $z \in D_{z_0}(r)$  and, since  $h(z_0) \neq 0$ , then  $z_0$  is a root of the extended  $\frac{1}{f}$  of multiplicity  $N$ . It is easy to prove in a similar way the converse, and we conclude that  $z_0$  is a pole of  $f$  of order  $N$  if and only if it is a root of  $\frac{1}{f}$  of multiplicity  $N$ .

**Example 5.9.2.** Many times we meet functions of the form  $f = \frac{p}{q}$ , where  $p, q$  are holomorphic in a neighborhood of  $z_0$ . For instance, if  $p, q$  are polynomials, then  $f$  is a rational function.

Let  $z_0$  be a root of  $p$  and  $q$  of multiplicity  $M \geq 0$  and  $N \geq 0$ , respectively. In this case we saw that there are holomorphic functions  $p_1$  and  $q_1$  in a neighborhood  $D_{z_0}(R)$  of  $z_0$  so that  $p(z) = (z - z_0)^M p_1(z)$  and  $q(z) = (z - z_0)^N q_1(z)$  for every  $z \in D_{z_0}(R)$  and also  $p_1(z_0) \neq 0$  and

$q_1(z_0) \neq 0$ . (Of course we consider the case that none of  $p, q$  is identically 0.) Then there is  $r$  with  $0 < r \leq R$  so that  $p_1(z) \neq 0$  and  $q_1(z) \neq 0$  for every  $z \in D_{z_0}(r)$ , and then we have

$$f(z) = \frac{p(z)}{q(z)} = (z - z_0)^{M-N} \frac{p_1(z)}{q_1(z)} = (z - z_0)^{M-N} g(z) \quad \text{for every } z \in D_{z_0}(r) \setminus \{z_0\},$$

where the function  $g = \frac{p_1}{q_1}$  is holomorphic in  $D_{z_0}(r)$  and  $g(z_0) = \frac{p_1(z_0)}{q_1(z_0)} \neq 0$ . Now we have two cases. If  $M \geq N$ , then  $z_0$  is a removable singularity of  $f$ , and  $f$  (after we extend it appropriately at  $z_0$ ) is holomorphic at  $z_0$  and  $z_0$  is a root of  $f$  of multiplicity  $M - N$ . If  $M < N$ , then  $z_0$  is a pole of order  $N - M$  of  $f$ .

Here are some concrete instances of this example.

**Example 5.9.3.** The function  $f(z) = \frac{z^2 - 3z + 2}{(z-2)^2}$  is holomorphic in  $\mathbb{C} \setminus \{2\}$ .

Since  $z^2 - 3z + 2 = (z - 2)(z - 1)$ , we have  $f(z) = \frac{z-1}{z-2}$  for  $z \neq 2$ . The function  $g(z) = z - 1$  is holomorphic in  $\mathbb{C}$  and  $g(2) = 1 \neq 0$ . Therefore, 2 is a pole of  $f$  of order 1.

**Example 5.9.4.** The function  $f(z) = \frac{e^z - 1}{z^3}$  is holomorphic in  $\mathbb{C} \setminus \{0\}$ .

The Taylor series of  $e^z - 1$  with center 0 is  $z + \frac{1}{2!}z^2 + \frac{1}{3!}z^3 + \dots$ . Hence  $e^z - 1 = zg(z)$  with  $g(z) = 1 + \frac{1}{2!}z + \frac{1}{3!}z^2 + \dots$ . The function  $g$  is holomorphic in  $\mathbb{C}$  and  $g(0) = 1 \neq 0$  and we have  $f(z) = \frac{g(z)}{z^2}$  for  $z \neq 0$ . Therefore, 0 is a pole of  $f$  of order 2.

**Example 5.9.5.** The function  $\cot z = \frac{\cos z}{\sin z}$  is holomorphic in  $\mathbb{C} \setminus \{k\pi \mid k \in \mathbb{Z}\}$ .

The points  $k\pi, k \in \mathbb{Z}$ , are isolated singularities of  $\cot z$  and we shall prove that they are all poles of order 1. We fix  $k \in \mathbb{Z}$ . The Taylor series of  $\sin z$  with center  $k\pi$  results from the Taylor series of  $\sin z$  with center 0, as follows

$$\begin{aligned} \sin z &= \sin((z - k\pi) + k\pi) = \cos k\pi \sin(z - k\pi) = (-1)^k \sin(z - k\pi) \\ &= (-1)^k \left( (z - k\pi) - \frac{1}{3!}(z - k\pi)^3 + \dots \right) = (-1)^k (z - k\pi) - \frac{(-1)^k}{3!}(z - k\pi)^3 + \dots \end{aligned}$$

Therefore,  $\sin z = (z - k\pi)q_1(z)$  for every  $z$ , where the function  $q_1$  is holomorphic in  $\mathbb{C}$  with  $q_1(k\pi) = (-1)^k$ . Hence,  $\cot z = \frac{\cos z}{(z - k\pi)q_1(z)} = \frac{g(z)}{z - k\pi}$  with  $g(z) = \frac{\cos z}{q_1(z)}$  and  $g$  is holomorphic in the disc  $D_{k\pi}(\pi)$  and  $g(k\pi) = \frac{\cos k\pi}{q_1(k\pi)} = 1$ . Therefore,  $k\pi$  is a pole of  $\cot z$  of order 1.

(Observe that  $D_{k\pi}(\pi)$  is the largest open disc with center  $k\pi$  which is contained in the domain of holomorphy of  $g$  because it is the largest open disc with center  $k\pi$  which does not contain any root of  $q_1$ . This is true because  $q_1(z) = \frac{\sin z}{z - k\pi}$  vanishes at every  $l\pi$  with  $l \in \mathbb{Z}, l \neq k$ .)

The Laurent series of  $\cot z$  in  $D_{k\pi}(0, \pi)$  is  $\cot z = \frac{1}{z - k\pi} + g'(k\pi) + \frac{1}{2}g''(k\pi)(z - k\pi) + \dots$ .

For the determination of poles there is a criterion similar to the criterion of Riemann for removable singularities.

**Proposition 5.4.** Let  $z_0$  be an isolated singularity of  $f$ . Then  $z_0$  is a pole of  $f$  if and only if  $\lim_{z \rightarrow z_0} f(z) = \infty$ .

*Proof.* There is a disc  $D_{z_0}(R)$  so that  $f$  is holomorphic in  $D_{z_0}(R) \setminus \{z_0\}$ .

If  $z_0$  is a pole of order  $N$  of  $f$ , then we saw that there is a function  $g$  holomorphic in  $D_{z_0}(R)$  so that  $g(z_0) \neq 0$  and  $f(z) = \frac{g(z)}{(z - z_0)^N}$  for every  $z \in D_{z_0}(R) \setminus \{z_0\}$ . This implies  $\lim_{z \rightarrow z_0} f(z) = \infty$ .

Conversely, let  $\lim_{z \rightarrow z_0} f(z) = \infty$ . Then there is  $r$  with  $0 < r \leq R$  so that  $f(z) \neq 0$  for every  $z \in D_{z_0}(r) \setminus \{z_0\}$  and then the function  $h = \frac{1}{f}$  is holomorphic in  $D_{z_0}(r) \setminus \{z_0\}$ . Since  $\lim_{z \rightarrow z_0} h(z) = \lim_{z \rightarrow z_0} \frac{1}{f(z)} = 0$ , the criterion of Riemann implies that  $z_0$  is a removable singularity of  $h$ . Therefore, we may define  $h$  appropriately at  $z_0$  so that it becomes holomorphic in  $D_{z_0}(r)$ : we set  $h(z_0) = \lim_{z \rightarrow z_0} h(z) = 0$ . It is clear that  $z_0$  is the only root of (the extended)  $h$  in  $D_{z_0}(r)$  and, if  $N$  is the multiplicity of this root, then  $h(z) = (z - z_0)^N h_1(z)$  where  $h_1$  is holomorphic in  $D_{z_0}(r)$  and has no root in  $D_{z_0}(r)$ . Thus, the function  $g = \frac{1}{h_1}$  is holomorphic in  $D_{z_0}(r)$  and, clearly, has no root in  $D_{z_0}(r)$ . Now we have altogether that  $f(z) = \frac{g(z)}{(z - z_0)^N}$  for every  $z \in D_{z_0}(r) \setminus \{z_0\}$  with  $g(z_0) \neq 0$ , and so  $z_0$  is a pole of  $f$  of order  $N$ .  $\square$

There is one more test for the case of a pole which also determines the exact order of the pole.

**Proposition 5.5.** *Let  $z_0$  be an isolated singularity of  $f$ . Then  $z_0$  is a pole of  $f$  of order  $N \geq 1$  if and only if the  $\lim_{z \rightarrow z_0} (z - z_0)^N f(z)$  exists and it is finite and  $\neq 0$ .*

*Proof.* If  $z_0$  is a pole of  $f$  of order  $N$ , then we repeat the beginning of the proof of proposition 5.4 and we get that  $\lim_{z \rightarrow z_0} (z - z_0)^N f(z) = \lim_{z \rightarrow z_0} g(z) = g(z_0) \neq 0$ .

Conversely, let  $\lim_{z \rightarrow z_0} (z - z_0)^N f(z)$  be finite and  $\neq 0$ . Riemann's criterion implies that the function  $g(z) = (z - z_0)^N f(z)$ , which is holomorphic in some ring  $D_{z_0}(R) \setminus \{z_0\}$ , can be extended at  $z_0$  by setting  $g(z_0) = \lim_{z \rightarrow z_0} g(z) = \lim_{z \rightarrow z_0} (z - z_0)^N f(z) \neq 0$ , and the extended  $g$  is holomorphic in  $D_{z_0}(R)$ . Therefore, there is a  $g$  holomorphic in  $D_{z_0}(R)$  with  $g(z_0) \neq 0$  so that  $f(z) = \frac{g(z)}{(z - z_0)^N}$  for every  $z \in D_{z_0}(R) \setminus \{z_0\}$  and  $z_0$  is a pole of  $f$  of order  $N$ .  $\square$

Finally, for the case of an essential singularity we have the following result.

**Proposition 5.6.** *Let  $z_0$  be an isolated singularity of  $f$ . Then  $z_0$  is an essential singularity of  $f$  if and only if the  $\lim_{z \rightarrow z_0} f(z)$  does not exist.*

*Proof.* By the criterion of Riemann,  $z_0$  is a removable singularity if and only if the  $\lim_{z \rightarrow z_0} f(z)$  exists and it is finite. Proposition 5.4 says that  $z_0$  is a pole if and only if  $\lim_{z \rightarrow z_0} f(z) = \infty$ .  $\square$

**Example 5.9.6.** In example 5.7.9 we saw that  $\sum_{-\infty}^{n=-1} \frac{1}{(-n)!} z^n + 1$  is the Laurent series of  $e^{\frac{1}{z}}$  in  $D_0(0, +\infty)$ . Hence 0 is an essential singularity of  $e^{\frac{1}{z}}$ .

Therefore, the  $\lim_{z \rightarrow 0} e^{\frac{1}{z}}$  does not exist. We can see this without proving first that 0 is an essential singularity of  $e^{\frac{1}{z}}$ . In fact, proving that the  $\lim_{z \rightarrow 0} e^{\frac{1}{z}}$  does not exist is another way to see that 0 is an essential singularity of  $e^{\frac{1}{z}}$ . Indeed, if  $z = x$  tends to 0 on the positive  $x$ -semiaxis, then  $|e^{\frac{1}{z}}| = e^{\frac{1}{x}} \rightarrow +\infty$ , and hence  $e^{\frac{1}{z}} \rightarrow \infty$ . If  $z = x$  tends to 0 on the negative  $x$ -semiaxis, then  $|e^{\frac{1}{z}}| = e^{\frac{1}{x}} \rightarrow 0$ , and hence  $e^{\frac{1}{z}} \rightarrow 0$ . Thus, the  $\lim_{z \rightarrow 0} e^{\frac{1}{z}}$  does not exist.

Let  $z_0$  be an isolated singularity of  $f$  and let  $\sum_{-\infty}^{+\infty} a_n (z - z_0)^n$  be the Laurent series of  $f$  in the ring  $D_{z_0}(0, R) = D_{z_0}(R) \setminus \{z_0\}$ . Then  $\sum_{-\infty}^{n=-1} a_n (z - z_0)^n$  is called the **singular part** of the Laurent series of  $f$  or, simply, the singular part of  $f$  at  $z_0$ . Also,  $\sum_{n=0}^{+\infty} a_n (z - z_0)^n$  is called the **regular part** of the Laurent series of  $f$  or, simply, the regular part of  $f$  at  $z_0$ .

We have seen that in the case of a removable singularity  $z_0$  the singular part of  $f$  at  $z_0$  is zero and the Laurent series of  $f$  at  $z_0$  consists only of its regular part. In the case of a pole  $z_0$  of  $f$  of order  $N$  the singular part at  $z_0$  is a finite sum of the form  $\sum_{n=1}^N \frac{a_{-n}}{(z - z_0)^n}$  with  $a_{-N} \neq 0$ . In this case the singular part is a *rational function* whose denominator is  $(z - z_0)^N$ . In the case of an essential singularity  $z_0$  the singular part at  $z_0$  has infinitely many terms.

If we subtract from  $f$  its singular part at its singularity  $z_0$ , then we get

$$f(z) - \sum_{-\infty}^{n=-1} a_n (z - z_0)^n = \sum_{n=0}^{+\infty} a_n (z - z_0)^n,$$

which is a power series of first type and hence converges in the disc  $D_{z_0}(R)$ , including the center  $z_0$ . Therefore,  $z_0$  is a removable singularity of the function  $F(z) = f(z) - \sum_{-\infty}^{n=-1} a_n (z - z_0)^n$  and if we define  $F$  to have value  $F(z_0) = a_0$  at  $z_0$ , then this function is holomorphic in  $D_{z_0}(R)$ .

We shall now establish the well known *analysis of a rational function into a sum of simple fractions*.

**Proposition 5.7.** *Let  $r = \frac{p}{q}$  be a rational function. We assume that the polynomials  $p, q$  have no common roots (and hence no common factors), that the degree of  $p$  is  $n$ , the degree of  $q$  is  $m$  and that  $z_1, \dots, z_k$  are the roots of  $q$  with corresponding multiplicities  $m_1, \dots, m_k$ . Then*

$$r(z) = p_1 \left( \frac{1}{z - z_1} \right) + \dots + p_k \left( \frac{1}{z - z_k} \right) + p_0(z),$$

where  $p_1, \dots, p_k$  are polynomials without constant terms and of degrees  $m_1, \dots, m_k$ , respectively, and  $p_0$  is either the null polynomial, if  $n < m$ , or a polynomial of degree  $n - m$ , if  $n \geq m$ .

*Proof.* We saw in example 5.9.2 that each  $z_j$  is a pole of  $r$  of degree  $m_j$ . Then the singular part of  $r$  at  $z_j$  has the form  $\sum_{l=1}^{m_j} \frac{a_{-l}}{(z-z_0)^l}$  with  $a_{-m_j} \neq 0$ . This can be written  $p_j(\frac{1}{z-z_j})$ , where  $p_j$  is the polynomial  $p_j(z) = \sum_{l=1}^{m_j} a_{-l}z^l$  without constant term and of degree  $m_j$ . We subtract from  $r$  all its singular parts at  $z_1, \dots, z_k$  and we form the function

$$p_0(z) = r(z) - \left( p_1\left(\frac{1}{z-z_1}\right) + \dots + p_k\left(\frac{1}{z-z_k}\right) \right).$$

This function is a rational function defined in  $\mathbb{C} \setminus \{z_1, \dots, z_k\}$  and its only *possible* poles are the points  $z_1, \dots, z_k$ . We observe, though, that every  $z_j$  is a removable singularity of  $r(z) - p_j(\frac{1}{z-z_j})$  and that each of  $p_1(\frac{1}{z-z_1}), \dots, p_k(\frac{1}{z-z_k})$ , besides  $p_j(\frac{1}{z-z_j})$ , is holomorphic at  $z_j$ . Thus, every  $z_j$  is a removable singularity of  $p_0$ . In other words, the rational function  $p_0$  has *no poles* and hence it is a polynomial. Now, we have the identity

$$r(z) = p_1\left(\frac{1}{z-z_1}\right) + \dots + p_k\left(\frac{1}{z-z_k}\right) + p_0(z)$$

and we consider two cases. If  $n < m$ , then  $\lim_{z \rightarrow \infty} r(z) = 0$  and, since  $\lim_{z \rightarrow \infty} p_j(\frac{1}{z-z_j}) = 0$  for every  $j$ , we have that  $\lim_{z \rightarrow \infty} p_0(z) = 0$ . Thus,  $p_0$  is the null polynomial. If  $n \geq m$ , then  $c = \lim_{z \rightarrow \infty} \frac{r(z)}{z^{n-m}}$  is a complex number  $\neq 0$ . Since  $\lim_{z \rightarrow \infty} p_j(\frac{1}{z-z_j})/z^{n-m} = 0$  for every  $j$ , we have that  $\lim_{z \rightarrow \infty} \frac{p_0(z)}{z^{n-m}} = c \neq 0$ . Thus the polynomial  $p_0$  has degree  $n - m$ .  $\square$

### Exercises.

**5.9.1.** Is 0 an isolated singularity of  $\frac{1}{\sin(1/z)}$ ?

**5.9.2.** Find the isolated (non-removable) singularities of:  $\frac{1}{z^2+5z+6}$ ,  $\frac{1}{(z^2-1)^2}$ ,  $\frac{e^z-1}{z}$ ,  $\frac{e^z-1}{z^3}$ ,  $\frac{z^2}{\sin z}$ ,  $\frac{1}{\sin z}$ ,  $\tan z$ ,  $\frac{1}{\sin^2 z}$ ,  $e^z + e^{1/z}$ ,  $\frac{1}{e^z-1}$ . Which of the singularities are poles and what is their order?

**5.9.3.** Find the initial four terms of the Laurent series at 0 of the functions:  $\cot z$ ,  $\frac{1}{\sin z}$ ,  $\frac{z}{\sin^2 z}$ ,  $\frac{1}{e^z-1}$ .

**5.9.4.** Prove that an isolated singularity of  $f$  cannot be a pole of  $e^f$ .

**5.9.5.** Let  $z_0$  be an isolated singularity of  $f$ , which is not constant in any neighborhood of  $z_0$ . If there is  $s \in \mathbb{R}$  so that  $\lim_{z \rightarrow z_0} |z - z_0|^s |f(z)| \in [0, +\infty]$ , prove that  $z_0$  is either a removable singularity or a pole of  $f$  and that there is  $m \in \mathbb{Z}$  so that

$$\lim_{z \rightarrow z_0} |z - z_0|^s |f(z)| \begin{cases} = 0, & \text{if } s > m \\ = +\infty, & \text{if } s < m \\ \in (0, +\infty), & \text{if } s = m \end{cases}$$

**5.9.6.** Let  $f$  be holomorphic in  $D_{z_0}(R) \setminus \{z_0\}$  and let either  $\operatorname{Re} f$  or  $\operatorname{Im} f$  be bounded either from above or from below in  $D_{z_0}(R) \setminus \{z_0\}$ . Prove that  $z_0$  is a removable singularity of  $f$ .

**5.9.7.** Let  $f$  be holomorphic in  $D_0(R) \setminus \{z_0\}$ , where  $R > 1$  and  $|z_0| = 1$ , and let  $z_0$  be a pole of  $f$ . If  $f(z) = \sum_{n=0}^{+\infty} a_n z^n$  is the Taylor series of  $f$  in  $D_0(1)$ , prove that  $\frac{a_n}{a_{n+1}} \rightarrow z_0$ .

**5.9.8.** Let  $\Omega$  be a region so that every point of  $\Omega$  is either a point of holomorphy or an isolated singularity of  $f$ . If the roots of  $f$  have an accumulation point in  $\Omega$ , which is not an essential singularity of  $f$ , prove that  $f$  is identically 0 in  $\Omega$ .

**5.9.9.** (i) Let  $z_0$  be an essential singularity of  $f$  and let  $w \in \mathbb{C}$ . Prove that for every  $r > 0$  the function  $\frac{1}{f-w}$  is *not* bounded in  $D_{z_0}(r) \setminus \{z_0\}$ .

(ii) Prove the **Casorati-Weierstrass theorem**. If  $z_0$  is an essential singularity of  $f$ , then for every  $w$  there is a sequence  $(z_n)$  with  $z_n \rightarrow z_0$  and  $z_n \neq z_0$  for every  $n$  so that  $f(z_n) \rightarrow w$ .

**5.9.10.** (i) Prove that every  $2k\pi i$ ,  $k \in \mathbb{Z}$ , is a pole of  $\frac{1}{e^z - 1}$  of order 1.

(ii) Prove that  $\frac{1}{e^z - 1} = \frac{1}{z} - \frac{1}{2} + \sum_{k=1}^{+\infty} (-1)^{k-1} \frac{B_k}{(2k)!} z^{2k-1}$  for  $|z| < 2\pi$ , where the numbers  $B_k$  satisfy the recursive relations  $\frac{1}{(2k+1)!} - \frac{1}{2(2k)!} + \sum_{\nu=1}^k \frac{(-1)^{\nu-1} B_\nu}{(2\nu)!(2k-2\nu+1)!} = 0$ ,  $k \geq 1$ . Evaluate  $B_1, B_2, B_3$ . The numbers  $B_k$  are called Bernoulli constants.

**5.9.11.** Look at exercises 5.7.6 and 5.8.11. We shall extend what we said in this section to the case of the point  $\infty$ .

(i) We say that  $\infty$  is an isolated singularity of  $f$  if  $f$  is holomorphic in some ring  $D_0(R, +\infty)$ . Let  $\sum_{-\infty}^{+\infty} a_n z^n$  be the Laurent series of  $f$  in this ring. If  $a_n = 0$  for every  $n \geq 1$ , then we say that  $\infty$  is a removable singularity of  $f$ . If  $a_n \neq 0$  for at least one  $n \geq 1$  and for only finitely many  $n \geq 1$ , then we say that  $\infty$  is a pole of  $f$ . Finally, if  $a_n \neq 0$  for infinitely many  $n \geq 1$ , then we say that  $\infty$  is an essential singularity of  $f$ .

Prove that  $\infty$  is a removable singularity of  $f$  if and only if  $\lim_{z \rightarrow \infty} \frac{f(z)}{z} = 0$ .

Prove that  $\infty$  is a pole of  $f$  if and only if  $\lim_{z \rightarrow \infty} f(z) = \infty$ .

Let  $\infty$  be a pole of  $f$  and let  $N$  be the largest  $n \geq 1$  with  $a_n \neq 0$ . Then we say that  $\infty$  is a pole of  $f$  of order  $N$ . Prove that  $\infty$  is a pole of  $f$  of order  $N$  if and only if there is a  $g$  holomorphic in  $D_0(R, +\infty) \cup \{\infty\}$  so that  $g(\infty) \neq 0$  and  $f(z) = z^N g(z)$  for every  $z \in D_0(R, +\infty)$ . Moreover, prove that  $\infty$  is a pole of  $f$  of order  $N$  if and only if the  $\lim_{z \rightarrow \infty} \frac{f(z)}{z^N}$  exists and it is finite and  $\neq 0$ . Prove that  $\infty$  is an essential singularity of  $f$  if and only if the  $\lim_{z \rightarrow \infty} f(z)$  does not exist.

(ii) Let  $r = \frac{p}{q}$  be a rational function and let  $n$  be the degree of the polynomial  $p$  and  $m$  be the degree of the polynomial  $q$ . Prove that  $\infty$  is a removable singularity of  $r$  if  $m \geq n$  and that it is a pole of  $r$  of order  $n - m$  if  $n > m$ . In particular, a polynomial  $p$  of degree  $n \geq 1$  has a pole of order  $n$  at  $\infty$ .

(iii) What kind of an isolated singularity is  $\infty$  for the functions  $e^z$ ,  $e^{\frac{1}{z}}$ ,  $z^2 e^{\frac{1}{z}}$ ,  $\sin z$ ,  $\sin \frac{1}{z}$ ,  $z^5 \sin \frac{1}{z}$ ?

(iv) What kind of an isolated singularity is  $\infty$  for any holomorphic branch of  $(z^2 - 1)^{\frac{1}{2}}$  in the region  $\mathbb{C} \setminus [-1, 1]$ ? (For the existence of such a branch look at exercise 4.4.9.)

(v) Is  $\infty$  an isolated singularity of  $\frac{1}{\sin z}$  or of  $\tan z$ ?

## 5.10 The open mapping theorem.

**Open mapping theorem.** *If  $f$  is holomorphic and not constant in the region  $\Omega$ , then  $f(U)$  is open for every open  $U \subseteq \Omega$ .*

*Proof.* Let  $U \subseteq \Omega$  be open. We shall prove that  $f(U)$  is also open, i.e. that every  $w_0 \in f(U)$  is an interior point of  $f(U)$ .

Since  $w_0 \in f(U)$  there is some  $z_0 \in U$  so that  $f(z_0) = w_0$ . Since  $U$  is open, there is  $r > 0$  so that  $\overline{D}_{z_0}(r) \subseteq U$ . Since  $f$  is non-constant in  $\Omega$ , the solution  $z_0$  of the equation  $f(z) = w_0$  is isolated. Therefore, we may take  $r$  small enough so that  $f(z) = w_0$  has no solution in  $\overline{D}_{z_0}(r)$  except  $z_0$ . This means that  $f(z) \neq w_0$  for every  $z \in \overline{D}_{z_0}(r) \setminus \{z_0\}$ . In particular,  $f(z) \neq w_0$  for every  $z \in C_{z_0}(r)$ . Now, the real function  $|f(z) - w_0|$  is continuous and the circle  $C_{z_0}(r)$  is compact. Therefore, the restriction of  $|f(z) - w_0|$  in  $C_{z_0}(r)$  attains a minimum value at some point of  $C_{z_0}(r)$  which is a *positive number*. We denote  $\epsilon$  this minimum and we have:

$$\epsilon = \min_{z \in C_{z_0}(r)} |f(z) - w_0| > 0. \quad (5.11)$$

Now, we consider any  $w \notin f(\overline{D}_{z_0}(r))$ . Again, the real function  $|f(z) - w|$  is continuous and the disc  $\overline{D}_{z_0}(r)$  is compact. Therefore the restriction of  $|f(z) - w|$  in  $\overline{D}_{z_0}(r)$  attains a minimum value at some point of  $\overline{D}_{z_0}(r)$  which is positive. But now we can say more: the function  $\frac{1}{f(z) - w}$  is *holomorphic* in  $D_{z_0}(r)$  and continuous in  $\overline{D}_{z_0}(r)$ . The second version of the maximum principle implies that the function  $\frac{1}{|f(z) - w|}$ , restricted in  $\overline{D}_{z_0}(r)$ , attains its maximum value at the boundary  $C_{z_0}(r)$ . Equivalently, the function  $|f(z) - w|$ , restricted in  $\overline{D}_{z_0}(r)$ , attains its minimum value at

the boundary  $C_{z_0}(r)$ . Since one of the values of  $|f(z) - w|$  in  $\overline{D}_{z_0}(r)$  is  $|f(z_0) - w| = |w_0 - w|$ , we get

$$|w_0 - w| \geq \min_{z \in C_{z_0}(r)} |f(z) - w|. \quad (5.12)$$

Now, we also have  $|f(z) - w| \geq |f(z) - w_0| - |w_0 - w|$  and, using (5.11) and (5.12), we get

$$|w_0 - w| \geq \min_{z \in C_{z_0}(r)} |f(z) - w| \geq \min_{z \in C_{z_0}(r)} |f(z) - w_0| - |w_0 - w| = \epsilon - |w_0 - w|.$$

Thus  $|w_0 - w| \geq \frac{\epsilon}{2}$  and we have proved that any  $w \notin f(\overline{D}_{z_0}(r))$  satisfies  $|w_0 - w| \geq \frac{\epsilon}{2}$ . This implies that every  $w \in D_{w_0}(\frac{\epsilon}{2})$  belongs to  $f(\overline{D}_{z_0}(r))$ . Hence  $D_{w_0}(\frac{\epsilon}{2}) \subseteq f(\overline{D}_{z_0}(r)) \subseteq f(U)$  and so  $w_0$  is an interior point of  $f(U)$ .  $\square$

### Exercises.

**5.10.1.** Prove the first maximum principle using the open mapping theorem.

## 5.11 Local mapping properties.

**Proposition 5.8.** *Let  $f$  be holomorphic in the open set  $\Omega$  and let  $z_0 \in \Omega$  with  $f'(z_0) \neq 0$ . Then there is an open set  $U \subseteq \Omega$  containing  $z_0$  so that  $W = f(U)$  is an open set containing  $w_0 = f(z_0)$  and so that the function  $f : U \rightarrow W$  is one-to-one. Moreover, the function  $f^{-1} : W \rightarrow U$  is holomorphic in  $W$ .*

*Proof.* We consider the Taylor series  $\sum_{n=0}^{+\infty} a_n(z - z_0)^n$  of  $f$  in its disc of convergence  $D_{z_0}(R) \subseteq \Omega$ . We know that the differentiated series  $\sum_{n=1}^{+\infty} n a_n(z - z_0)^{n-1}$  converges absolutely in the same disc, i.e.  $\sum_{n=1}^{+\infty} n |a_n| |z - z_0|^{n-1} < +\infty$  for every  $z \in D_{z_0}(R)$ . Thus,  $\sum_{n=1}^{+\infty} n |a_n| (z - z_0)^{n-1}$  converges in  $D_{z_0}(R)$  and defines a continuous function in this disc. In particular, we have that  $\lim_{z \rightarrow z_0} \sum_{n=1}^{+\infty} n |a_n| (z - z_0)^{n-1} = |a_1|$  or, equivalently,  $\lim_{z \rightarrow z_0} \sum_{n=2}^{+\infty} n |a_n| (z - z_0)^{n-1} = 0$ . Since  $a_1 = f'(z_0) \neq 0$ , there is a small enough  $r > 0$  so that

$$\sum_{n=2}^{+\infty} n |a_n| r^{n-1} < |a_1|.$$

We shall see now that  $f : D_{z_0}(r) \rightarrow \mathbb{C}$  is one-to-one. Assume that this is not the case and that there are  $z_1, z_2 \in D_{z_0}(r)$  so that  $z_1 \neq z_2$  and  $f(z_1) = f(z_2)$ . Then  $\sum_{n=0}^{+\infty} a_n(z_1 - z_0)^n = \sum_{n=0}^{+\infty} a_n(z_2 - z_0)^n$  and hence

$$a_1 = \sum_{n=2}^{+\infty} a_n((z_1 - z_0)^{n-1} + (z_1 - z_0)^{n-2}(z_2 - z_0) + \cdots + (z_1 - z_0)(z_2 - z_0)^{n-2} + (z_2 - z_0)^{n-1}).$$

This implies  $|a_1| \leq \sum_{n=2}^{+\infty} n |a_n| r^{n-1}$  and we arrive at a contradiction.

Since  $f'$  is continuous at  $z_0$  and  $f'(z_0) \neq 0$ , by taking a smaller  $r > 0$  if necessary, we may suppose that  $f'(z) \neq 0$  for every  $z \in D_{z_0}(r)$ . Now we take  $U = D_{z_0}(r)$ . From the open mapping theorem we have that the set  $W = f(U) = f(D_{z_0}(r))$  is open. We have already proved that  $f : U \rightarrow W$  is one-to-one and hence the inverse mapping  $f^{-1} : W \rightarrow U$  is defined. Now it is easy to see that this inverse mapping is continuous in  $W$ . Indeed let  $w \in W$ . Then there is (a unique)  $z \in U$  so that  $f(z) = w$ . We take any  $\epsilon > 0$  small enough so that  $D_z(\epsilon) \subseteq U$ . Then the set  $f(D_z(\epsilon))$  is open and contains  $w$ . Hence there is  $\delta > 0$  so that  $D_w(\delta) \subseteq f(D_z(\epsilon))$ . Then for every  $w' \in D_w(\delta)$  the (unique)  $z' \in U$  which satisfies  $f(z') = w'$  is contained in  $D_z(\epsilon)$ . This says that for every  $w' \in W$  with  $|w' - w| < \delta$  we have  $|f^{-1}(w') - f^{-1}(w)| = |z' - z| < \epsilon$  and the function  $f^{-1} : W \rightarrow U$  is continuous at every  $w \in W$ . Now, proposition 3.4 implies that  $f^{-1} : W \rightarrow U$  is holomorphic in  $W$ .  $\square$

**Theorem 5.2.** *Let  $f$  be holomorphic in the region  $\Omega$  and let  $z_0 \in \Omega$  and  $w_0 = f(z_0)$ . Let  $z_0$  be a solution of  $f(z) = w_0$  of multiplicity  $N$ . Then there is an open set  $U \subseteq \Omega$  containing  $z_0$  so that  $W = f(U)$  is an open set containing  $w_0 = f(z_0)$  and so that the function  $f : U \rightarrow W$  is  $N$ -to-one.*



*Proof.* We know that there is a disc  $D_{z_0}(R)$  and a function  $g$  holomorphic in  $D_{z_0}(R)$  so that  $f(z) - w_0 = (z - z_0)^N g(z)$  for every  $z \in D_{z_0}(R)$  and  $g(z_0) \neq 0$ . By the continuity of  $g$  we have that there is  $r \leq R$  so that  $g(z) \neq 0$  for every  $z \in D_{z_0}(r)$ . Then the function  $\frac{f'}{g}$  is holomorphic in  $D_{z_0}(r)$  and the theorem of Cauchy in convex regions implies that  $\oint_{\gamma} \frac{g'(z)}{g(z)} dz = 0$  for every closed curve  $\gamma$  in  $D_{z_0}(r)$ . Now, theorem 4.1 implies that there is a holomorphic branch of  $\log g$  in  $D_{z_0}(r)$  and then example 4.4.4 says that there is a holomorphic branch of  $g^{1/N}$  in  $D_{z_0}(r)$ . I.e. there is a function  $\phi$  holomorphic in  $D_{z_0}(r)$  so that  $\phi(z)^N = g(z)$  for every  $z \in D_{z_0}(r)$ . Now we consider the function  $h(z) = (z - z_0)\phi(z)$ . This is holomorphic in  $D_{z_0}(r)$  and we have that  $f(z) - w_0 = h(z)^N$  for every  $z \in D_{z_0}(r)$ . Moreover,  $h'(z_0) = \phi(z_0) \neq 0$ . Proposition 5.8, applied to  $h$ , implies that there is an open set  $U_0 \subseteq D_{z_0}(r)$  containing  $z_0$  so that  $W_0 = h(U_0)$  is an open set containing  $h(z_0) = 0$  and so that the function  $h : U_0 \rightarrow W_0$  is one-to-one. Now, we consider a disc  $D_0(r_0) \subseteq W_0$  and the open set  $U = h^{-1}(D_0(r_0)) \subseteq U_0$ . Then  $h : U \rightarrow D_0(r_0)$  is holomorphic in  $U$ , onto  $D_0(r_0)$  and one-to-one in  $U$ . Moreover, we have that  $f(z) - w_0 = h(z)^N$  for every  $z \in U$ . Since the  $N$ -th power  $w = \zeta^N$  maps the disc  $D_0(r_0)$  onto the disc  $D_0(r_0^N)$  and in an  $N$ -to-one manner, we conclude that  $f : U \rightarrow W$  is  $N$ -to-one, where  $W$  is the disc  $D_{w_0}(r_0^N)$ .  $\square$

In the proof of theorem 5.2 if we take any linear segment  $[w_0, w]$  in the disc  $D_{w_0}(r_0^N)$ , where  $w$  is a point of the circle  $C_{w_0}(r_0^N)$ , then, through the mapping  $w = w_0 + \zeta^N$ , this linear segment corresponds to  $N$  linear segments  $[0, z_1], \dots, [0, z_N]$  in the disc  $D_0(r_0)$ , where  $z_1, \dots, z_N$  are  $N$  points on the circle  $C_0(r_0)$ . These  $N$  linear segments form  $N$  successive angles at 0 all equal to  $\frac{2\pi}{N}$ . Now the one-to-one function  $h^{-1} : D_0(r_0) \rightarrow U$  maps these linear segments onto  $N$  curves  $\gamma_1, \dots, \gamma_N$  with common initial endpoint  $z_0$  and  $N$  corresponding final endpoints on  $\partial U$ . Since  $h'(z_0) \neq 0$ , the conformality of  $h$  at  $z_0$  implies that  $\gamma_1, \dots, \gamma_N$  form  $N$  successive angles at  $z_0$  all equal to  $\frac{2\pi}{N}$ . The  $N$  successive “angular” regions  $U_1, \dots, U_N$  in  $U$  between the curves  $\gamma_1, \dots, \gamma_N$  are mapped by  $h$  onto the corresponding successive angular regions  $A_1, \dots, A_N$  in  $D_0(r_0)$  between the linear segments  $[0, z_1], \dots, [0, z_N]$  and these are then mapped by the mapping  $w = w_0 + \zeta^N$  onto the same region  $B = D_{w_0}(r_0^N) \setminus [w_0, w]$ . We conclude that  $f$ , which is the composition of the two mappings, maps each of  $U_1, \dots, U_N$  in  $U$  onto  $B$  and in an one-to-one manner.

### Exercises.

**5.11.1.** Let  $f$  be holomorphic in  $D_0(R)$ ,  $f'(0) \neq 0$  and  $n \in \mathbb{N}$ . Prove that there is  $r > 0$  and there is  $g$  holomorphic in  $D_0(r)$  so that  $f(z^n) = f(0) + (g(z))^n$  for every  $z \in D_0(r)$ .

**5.11.2.** Let  $\Omega_1, \Omega_2$  be two regions, let  $f : \Omega_1 \rightarrow \Omega_2$  and  $g : \Omega_2 \rightarrow \mathbb{C}$  be non-constant functions and let  $h = g \circ f$ .

(i) If  $f, h$  are holomorphic in  $\Omega_1$ , is  $g$  holomorphic in  $\Omega_2$ ?

(ii) If  $g, h$  are holomorphic in  $\Omega_2, \Omega_1$ , respectively, is  $f$  holomorphic in  $\Omega_1$ ?

**5.11.3.** If  $f$  is holomorphic and one-to-one in  $\mathbb{C}$ , prove that there are  $a \neq 0$  and  $b$  so that  $f(z) = az + b$  for every  $z$ .

## Chapter 6

# Global behaviour of holomorphic functions.

### 6.1 Index of a closed curve with respect to a point.

Let  $A \subseteq \mathbb{C}$  and  $g : A \rightarrow \mathbb{C} \setminus \{0\}$  be continuous in  $A$ . We say that the function  $h$  is a **continuous branch of arg  $g$**  in  $A$  if  $h$  is continuous in  $A$  and for every  $w \in A$  we have that  $h(w)$  is an element of  $\arg g(w)$  or, equivalently,

$$g(w) = |g(w)|e^{ih(w)} \quad \text{for every } w \in A.$$

We recall the notion of a continuous branch of  $\log g$ . We say that  $f$  is a continuous branch of  $\log g$  if  $f$  is continuous in  $A$  and  $f(w)$  is an element of  $\log g(w)$  or, equivalently,

$$e^{f(w)} = g(w) \quad \text{for every } w \in A.$$

**Proposition 6.1.** *Let  $A \subseteq \mathbb{C}$  and  $g : A \rightarrow \mathbb{C} \setminus \{0\}$  be continuous in  $A$ . Then there is a one-to-one correspondence between continuous branches of  $\log g$  and continuous branches of  $\arg g$  in  $A$ .*

*Proof.* If  $h$  is a continuous branch of  $\arg g$  in  $A$ , then the function

$$f = \ln |g| + ih \tag{6.1}$$

is a continuous branch of  $\log g$  in  $A$ . Indeed, it is clear that  $f$  is continuous in  $A$  and also that  $e^{f(w)} = e^{\ln |g(w)|} e^{ih(w)} = |g(w)| e^{ih(w)} = g(w)$  for every  $w \in A$ .

Conversely, if  $f$  is a continuous branch of  $\log g$  in  $A$ , then  $h$ , defined through (6.1), is a continuous branch of  $\arg g$  in  $A$ . Indeed,  $h$  is continuous in  $A$  and also  $e^{ih(w)} = e^{f(w)} e^{-\ln |g(w)|} = \frac{g(w)}{|g(w)|}$  and hence  $g(w) = |g(w)| e^{ih(w)}$  for every  $w \in A$ .  $\square$

In other words, relation (6.1) says that, if we have a continuous branch  $f$  of  $\log g$  in  $A$ , then the imaginary part  $h$  of  $f$  is a continuous branch of  $\arg g$  in  $A$ . Conversely, if we have a continuous branch  $h$  of  $\arg g$  in  $A$ , then the function  $f$  with imaginary part  $h$  and real part  $\ln |g|$  is a continuous branch of  $\log g$  in  $A$ .

The next result is analogous to proposition 4.6 and their proofs are almost identical.

**Proposition 6.2.** *Let  $g : A \rightarrow \mathbb{C} \setminus \{0\}$  be continuous in  $A \subseteq \mathbb{C}$ .*

(i) *If  $h_1$  is a continuous branch of  $\arg g$  in  $A$  and  $h_2 - h_1 = k2\pi$  in  $A$ , where  $k$  is a fixed integer, then  $h_2$  is also a continuous branch of  $\arg g$  in  $A$ .*

(ii) *If, moreover,  $A$  is connected and  $h_1, h_2$  are continuous branches of  $\arg g$  in  $A$ , then  $h_2 - h_1 = k2\pi$  in  $A$ , where  $k$  is a fixed integer. In particular, if  $h_1(w_0) = h_2(w_0)$  for some  $w_0 \in A$ , then  $h_1 = h_2$  in  $A$ .*

**Proposition 6.3.** Let  $g_1, g_2 : A \rightarrow \mathbb{C} \setminus \{0\}$  be continuous in  $A \subseteq \mathbb{C}$ .

(i) If  $f_1, f_2$  are continuous branches of  $\log g_1, \log g_2$  in  $A$ , then  $f_1 + f_2$  is a continuous branch of  $\log(g_1 g_2)$  in  $A$ .

(ii) If  $h_1, h_2$  are continuous branches of  $\arg g_1, \arg g_2$  in  $A$ , then  $h_1 + h_2$  is a continuous branch of  $\arg(g_1 g_2)$  in  $A$ .

*Proof.* (i)  $f_1 + f_2$  is continuous in  $A$  and also  $e^{f_1(w)+f_2(w)} = e^{f_1(w)}e^{f_2(w)} = g_1(w)g_2(w)$  for every  $w \in A$ .

(ii) Just as in (i). □

Proposition 6.4 is the first existence result of this section:  $A$  is an interval in  $\mathbb{R}$ .

**Proposition 6.4.** Let  $g : [a, b] \rightarrow \mathbb{C} \setminus \{0\}$  be continuous in the interval  $[a, b]$ . Then there is a continuous branch of  $\log g$  and a continuous branch of  $\arg g$  in  $[a, b]$ .

*Proof.* It is enough to prove the existence of a continuous branch of  $\log g$ .

Since  $g$  is continuous in  $[a, b]$ , there is  $\epsilon > 0$  so that  $|g(t)| \geq \epsilon$  for every  $t \in [a, b]$ . Now,  $g$  is also uniformly continuous in  $[a, b]$  and hence there is  $\delta > 0$  so that  $|g(t') - g(t'')| < \epsilon$  for every  $t', t'' \in [a, b]$  with  $|t' - t''| < \delta$ . We take successive points  $a = t_0 < t_1 < \dots < t_{n-1} < t_n = b$  so that  $t_k - t_{k-1} < \delta$  for every  $k = 1, \dots, n$ . Then, for every  $k = 1, \dots, n$ , the set  $\{g(t) \mid t \in [t_{k-1}, t_k]\}$  is contained in the disc  $D_{g(t_k)}(\epsilon)$ . Since  $|g(t_k)| \geq \epsilon$ , the disc  $D_{g(t_k)}(\epsilon)$  does not contain 0 and hence a continuous branch of  $\log$  is defined in this disc. Then example 4.3.3 says that there is a continuous branch, say  $f_k$ , of  $\log g$  in  $[t_{k-1}, t_k]$ .

Now,  $f_1$  is a continuous branch of  $\log g$  in  $[t_0, t_1]$  and  $f_2$  is a continuous branch of  $\log g$  in  $[t_1, t_2]$ . Then  $f_2(t_1) - f_1(t_1) = m2\pi i$  for some  $m \in \mathbb{Z}$ . We replace the function  $f_2$  with the function  $f_2 - m2\pi i$  and the new function  $f_2$  is also a continuous branch of  $\log g$  in  $[t_1, t_2]$  with  $f_2(t_1) = f_1(t_1)$ . Working with the (new) function  $f_2$  and the function  $f_3$  which is a continuous branch of  $\log g$  in  $[t_2, t_3]$ , we see as before that  $f_3(t_2) - f_2(t_2) = l2\pi i$  for some  $l \in \mathbb{Z}$ . We replace the function  $f_3$  with the function  $f_3 - l2\pi i$  and the new function  $f_3$  is also a continuous branch of  $\log g$  in  $[t_2, t_3]$  with  $f_3(t_2) = f_2(t_2)$ . We continue inductively and finally we end up with continuous branches  $f_k$  of  $\log g$  in  $[t_{k-1}, t_k]$  for every  $k = 1, \dots, n$  so that  $f_k(t_k) = f_{k+1}(t_k)$  for every  $k = 1, \dots, n-1$ . Therefore, the function  $f : [a, b] \rightarrow \mathbb{C}$  which is defined to be equal to  $f_k$  in the corresponding interval  $[t_{k-1}, t_k]$  for every  $k = 1, \dots, n$  is continuous in  $[a, b]$ . Moreover,  $f$  is a continuous branch of  $\log g$  in every  $[t_{k-1}, t_k]$  and hence in  $[a, b]$ . □

We consider any curve  $\gamma : [a, b] \rightarrow \mathbb{C} \setminus \{z_0\}$ . Then the function  $\gamma - z_0 : [a, b] \rightarrow \mathbb{C} \setminus \{0\}$  is continuous in  $[a, b]$  and, according to proposition 6.4, there is a continuous branch  $f$  of  $\log(\gamma - z_0)$  and a continuous branch  $h$  of  $\arg(\gamma - z_0)$  in  $[a, b]$  related by

$$f = \ln |\gamma - z_0| + ih. \quad (6.2)$$

Then the functions  $f + k2\pi i$  and  $h + k2\pi$ , where  $k$  is an arbitrary, but constant, integer, are also continuous branches of  $\log(\gamma - z_0)$  and  $\arg(\gamma - z_0)$  in  $[a, b]$ . Moreover, since  $[a, b]$  is connected, these are all the continuous branches of  $\log(\gamma - z_0)$  and  $\arg(\gamma - z_0)$  in  $[a, b]$ .

Now, let  $h$  be any continuous branch of  $\arg(\gamma - z_0)$  in  $[a, b]$ . We observe that the expression  $h(b) - h(a)$  is independent of the particular choice of  $h$ . Indeed, if  $h_1$  is another continuous branch of  $\arg(\gamma - z_0)$  in  $[a, b]$ , then there is a constant integer  $k$  so that  $h_1 = h + k2\pi$  in  $[a, b]$  and then we have  $h_1(b) - h_1(a) = (h(b) + k2\pi) - (h(a) + k2\pi) = h(b) - h(a)$ .

Now, the expression

$$\Delta \arg(\gamma - z_0) = h(b) - h(a)$$

is called **total increment of argument** or **total increment of angle** over the curve  $\gamma$  with respect to  $z_0$ .

Let us consider the important special case when the curve  $\gamma : [a, b] \rightarrow \mathbb{C} \setminus \{z_0\}$  is closed, i.e. when  $\gamma(b) = \gamma(a)$ . This implies  $\gamma(b) - z_0 = \gamma(a) - z_0$ , and hence  $\ln |\gamma(b) - z_0| = \ln |\gamma(a) - z_0|$ .

It also implies that  $h(b)$  and  $h(a)$  differ by some integer multiple of  $2\pi$ : indeed, both  $h(b), h(a)$  are values of  $\arg(\gamma(b) - z_0) = \arg(\gamma(a) - z_0)$ . Therefore the expression  $\Delta \arg(\gamma - z_0) = h(b) - h(a)$  is an integer multiple of  $2\pi$ . Then the integer

$$n(\gamma; z_0) = \frac{\Delta \arg(\gamma - z_0)}{2\pi}$$

is called **rotation number** or **index** of  $\gamma$  with respect to  $z_0$ .

It is easy to see the geometric content of the notion of rotation number or index of a closed curve  $\gamma$  with respect to  $z_0$ . When  $t$  increases in  $[a, b]$ , the angle  $h(t)$  of  $\gamma(t) - z_0$  varies continuously from  $h(a)$  to  $h(b)$ . Since  $\Delta \arg(\gamma - z_0) = h(b) - h(a) = k2\pi$  for a certain integer  $k$ , the number  $n(\gamma; z_0) = \frac{\Delta \arg(\gamma - z_0)}{2\pi} = \frac{h(b) - h(a)}{2\pi} = k$  shows the numbers of complete rotations of the continuously moving point  $\gamma(t)$  around  $z_0$ .

**Proposition 6.5.** *Let  $\gamma_1$  and  $\gamma_2$  be closed curves in  $\mathbb{C} \setminus \{z_0\}$  with the same endpoints. Then  $\gamma_1 + \gamma_2$  is defined and it is also a closed curve in  $\mathbb{C} \setminus \{z_0\}$  and also  $n(\gamma_1 + \gamma_2; z_0) = n(\gamma_1; z_0) + n(\gamma_2; z_0)$ .*

*Proof.* Let  $\gamma_1 : [a, b] \rightarrow \mathbb{C} \setminus \{z_0\}$  and  $\gamma_2 : [b, c] \rightarrow \mathbb{C} \setminus \{z_0\}$  be the two curves and  $h_1 : [a, b] \rightarrow \mathbb{R}$  and  $h_2 : [b, c] \rightarrow \mathbb{R}$  be continuous branches of  $\arg(\gamma_1 - z_0)$  and  $\arg(\gamma_2 - z_0)$ . We may redefine  $h_2$  by adding to it an appropriate integer multiple of  $2\pi$  so that  $h_2(b) = h_1(b)$ . Then the function  $h : [a, c] \rightarrow \mathbb{R}$  which equals  $h_1$  in  $[a, b]$  and  $h_2$  in  $[b, c]$  is a continuous branch of  $\log((\gamma_1 + \gamma_2) - z_0)$  in  $[a, c]$ . Therefore  $h(c) - h(a) = h(c) - h(b) + h(b) - h(a) = h_2(c) - h_2(b) + h_1(b) - h_1(a)$  and hence  $n(\gamma_1 + \gamma_2; z_0) = n(\gamma_1; z_0) + n(\gamma_2; z_0)$ .  $\square$

**Proposition 6.6.** *Let  $\gamma_1, \gamma_2$  be closed curves in  $\mathbb{C} \setminus \{z_0\}$  so that each is a reparametrization of the other. Then  $n(\gamma_2; z_0) = n(\gamma_1; z_0)$ .*

*Proof.* Let  $\gamma_1 : [a, b] \rightarrow \mathbb{C} \setminus \{z_0\}$  and  $\gamma_2 : [c, d] \rightarrow \mathbb{C} \setminus \{z_0\}$  be the two curves and  $\sigma : [c, d] \rightarrow [a, b]$  be the change of parameter so that  $\gamma_2 = \gamma_1 \circ \sigma$ . If  $h$  is a continuous branch of  $\arg(\gamma_1 - z_0)$  in  $[a, b]$ , then  $h \circ \sigma$  is a continuous branch of  $\arg(\gamma_2 - z_0)$  in  $[c, d]$ . Indeed, from  $e^{ih(t)} = \frac{\gamma_1(t) - z_0}{|\gamma_1(t) - z_0|}$  for every  $t \in [a, b]$  we get  $e^{ih(\sigma(s))} = \frac{\gamma_1(\sigma(s)) - z_0}{|\gamma_1(\sigma(s)) - z_0|} = \frac{\gamma_2(s) - z_0}{|\gamma_2(s) - z_0|}$  for every  $s \in [c, d]$ . Therefore, from  $h(\sigma(d)) - h(\sigma(c)) = h(b) - h(a)$  we get  $n(\gamma_2; z_0) = n(\gamma_1; z_0)$ .  $\square$

**Proposition 6.7.** *Let  $\gamma$  be a closed curve in  $\mathbb{C} \setminus \{z_0\}$ . Then  $n(\neg\gamma; z_0) = -n(\gamma; z_0)$ .*

*Proof.* Let  $\gamma, \neg\gamma : [a, b] \rightarrow \mathbb{C} \setminus \{z_0\}$  be the two curves. Then  $\neg\gamma(t) = \gamma(a + b - t)$  for every  $t \in [a, b]$ . If  $h$  is a continuous branch of  $\arg(\gamma - z_0)$  in  $[a, b]$ , then the function  $k(t) = h(a + b - t)$  is a continuous branch of  $\arg(\neg\gamma - z_0)$  in  $[a, b]$ . Indeed, from  $e^{ih(t)} = \frac{\gamma(t) - z_0}{|\gamma(t) - z_0|}$  for every  $t \in [a, b]$  we get  $e^{ik(t)} = e^{ih(a+b-t)} = \frac{\gamma(a+b-t) - z_0}{|\gamma(a+b-t) - z_0|} = \frac{\neg\gamma(t) - z_0}{|\neg\gamma(t) - z_0|}$  for every  $t \in [a, b]$ . Therefore, from  $k(b) - k(a) = h(a) - h(b)$  we get  $n(\neg\gamma; z_0) = -n(\gamma; z_0)$ .  $\square$

**Proposition 6.8.** *Let  $\gamma$  be a closed curve in  $\mathbb{C} \setminus \{z_0\}$ . If a continuous branch of  $\log$  is defined in the set  $\gamma^* - z_0 \subseteq \mathbb{C} \setminus \{0\}$ , then  $n(\gamma; z_0) = 0$ .*

*Proof.* We shall apply the argument of example 4.3.3. Let  $\gamma : [a, b] \rightarrow \mathbb{C} \setminus \{z_0\}$  be the closed curve and  $q$  be a continuous branch of  $\log$  in  $\gamma^* - z_0 \subseteq \mathbb{C} \setminus \{0\}$ . Then the function  $f = q \circ (\gamma - z_0)$  is a continuous branch of  $\log(\gamma - z_0)$  in  $[a, b]$ . Now, since the curve  $\gamma$  is closed, we have  $\gamma(b) = \gamma(a)$  and hence  $f(b) = q(\gamma(b) - z_0) = q(\gamma(a) - z_0) = f(a)$ . According to (6.2), the imaginary part  $h$  of  $f$  is a continuous branch of  $\arg(\gamma - z_0)$  in  $[a, b]$ . From  $f(b) = f(a)$  we get  $h(b) = h(a)$  and hence  $n(\gamma; z_0) = \frac{h(b) - h(a)}{2\pi} = 0$ .  $\square$

**Example 6.1.1.** We consider the set  $A = \mathbb{C} \setminus l$ , where  $l$  is any halfline with vertex  $z_0$ . We know that a holomorphic branch of  $\log(z - z_0)$  exists in  $A$  and hence  $n(\gamma; z_0) = 0$  for every closed curve  $\gamma$  in  $A$ . This is geometrically obvious: *since  $\gamma$  is in  $A$ , it does not intersect the halfline  $l$  with vertex  $z_0$ , and hence it cannot make any complete rotation around  $z_0$ .*

**Proposition 6.9.** Let  $\gamma_1, \gamma_2 : [a, b] \rightarrow \mathbb{C} \setminus \{z_0\}$  be closed curves such that  $|\gamma_1(t) - \gamma_2(t)| < |\gamma_2(t) - z_0|$  for every  $t \in [a, b]$ . Then  $n(\gamma_1; z_0) = n(\gamma_2; z_0)$ .

*Proof.* From  $|\gamma_1(t) - \gamma_2(t)| < |\gamma_2(t) - z_0|$  we get

$$\left| \frac{\gamma_1(t) - z_0}{\gamma_2(t) - z_0} - 1 \right| < 1 \quad \text{for every } t \in [a, b].$$

Now, we apply again the argument of example 4.3.3. We consider the function  $g : [a, b] \rightarrow D_1(1)$  with  $g(t) = \frac{\gamma_1(t) - z_0}{\gamma_2(t) - z_0}$  for every  $t \in [a, b]$ . Let  $q$  be a continuous branch of  $\log$  in  $D_1(1)$ . Then  $f = q \circ g$  is a continuous branch of  $\log g$  in  $[a, b]$ . Since the curves  $\gamma_1, \gamma_2$  are closed, we have that  $g(b) = g(a)$  and hence  $f(b) = q(g(b)) = q(g(a)) = f(a)$ . According to (6.2), the imaginary part  $h$  of  $f$  is a continuous branch of  $\arg g$  in  $[a, b]$ . From  $f(b) = f(a)$  we get  $h(b) = h(a)$ .

Now let  $h_2$  be a continuous branch of  $\arg(\gamma_2 - z_0)$  in  $[a, b]$ . Since,  $\gamma_1 - z_0 = (\gamma_2 - z_0)g$  in  $[a, b]$ , proposition 6.3 implies that  $h_1 = h_2 + h$  is a continuous branch of  $\arg(\gamma_1 - z_0)$  in  $[a, b]$ . Therefore,  $h_1(b) - h_1(a) = h_2(b) - h_2(a) + h(b) - h(a) = h_2(b) - h_2(a)$  and hence  $n(\gamma_1; z_0) = n(\gamma_2; z_0)$   $\square$

For every curve  $\gamma$  in  $\mathbb{C} \setminus \{z_0\}$  we may consider the translated curve  $\gamma_{z_0} = \gamma - z_0$  in  $\mathbb{C} \setminus \{0\}$ . It is clear that  $n(\gamma; z_0) = n(\gamma_{z_0}; 0)$ .

**Proposition 6.10.** Let  $\gamma$  be a closed curve. Then the integer-valued function  $n(\gamma; \cdot) : \mathbb{C} \setminus \gamma^* \rightarrow \mathbb{Z}$  is constant in every connected component of the open set  $\mathbb{C} \setminus \gamma^*$ . We also have that  $n(\gamma; z) = 0$  for every  $z$  in the unbounded connected component of  $\mathbb{C} \setminus \gamma^*$ .

*Proof.* Let  $\gamma : [a, b] \rightarrow \mathbb{C}$  be the curve and let  $z \in \mathbb{C} \setminus \gamma^*$ . Then there is some disc  $D_z(r)$  contained in  $\mathbb{C} \setminus \gamma^*$  and hence  $|w - z| < r \leq |\gamma(t) - z|$  for every  $t \in [a, b]$  and every  $w \in D_z(r)$ .

We take any  $w \in D_z(r)$  and we consider the translated curves  $\gamma_z = \gamma - z$  and  $\gamma_w = \gamma - w$ . Then we have that  $|\gamma_w(t) - \gamma_z(t)| = |w - z| < |\gamma_z(t)|$  for every  $t \in [a, b]$ . Proposition 6.9 implies that  $n(\gamma_w; 0) = n(\gamma_z; 0)$  and hence  $n(\gamma; w) = n(\gamma; z)$ .

We proved that the function  $n(\gamma; \cdot) : \mathbb{C} \setminus \gamma^* \rightarrow \mathbb{Z}$  is locally constant: for every  $z \in \mathbb{C} \setminus \gamma^*$  there is a disc  $D_z(r)$  so that  $n(\gamma; w) = n(\gamma; z)$  for every  $w \in D_z(r)$ . Of course, this implies that  $n(\gamma; \cdot)$  is continuous in  $\mathbb{C} \setminus \gamma^*$ . Now, let  $\Omega$  be a connected component of  $\mathbb{C} \setminus \gamma^*$ . Since  $n(\gamma; \cdot)$  is continuous and integer-valued in the connected set  $\Omega$ , it is constant in  $\Omega$ .

Since  $\gamma^*$  is bounded, it is contained in some disc  $\overline{D}_0(R)$ . Then the connected ring  $D_0(R, +\infty)$  is contained in  $\mathbb{C} \setminus \gamma^*$  and hence it is contained in (exactly) one of the connected components, call it  $\Omega$ , of  $\mathbb{C} \setminus \gamma^*$ . Then  $\Omega$  is the only unbounded connected component of  $\mathbb{C} \setminus \gamma^*$  and we shall prove that  $n(\gamma; z) = 0$  for every  $z \in \Omega$ . Let  $c$  be the constant value of  $n(\gamma; z)$  in  $\Omega$ . We take any  $z_0 \in D_0(R, +\infty)$  and then obviously there is a halfline  $l$  with vertex  $z_0$  which does not intersect the disc  $\overline{D}_0(R)$  and hence it does not intersect  $\gamma^*$  either. From example 6.1.1 we have that  $c = n(\gamma; z_0) = 0$ .  $\square$

Proposition 6.10 says that if  $z_1, z_2$  are in the same connected component of the complement of the trajectory of the closed curve  $\gamma$ , then the number of complete rotations of  $\gamma$  around  $z_1$  is equal to the number of complete rotations of  $\gamma$  around  $z_2$ .

Let  $\gamma$  be a closed curve and  $z \notin \gamma^*$ . We say that  $\gamma$  **surrounds**  $z$  if  $n(\gamma; z) \neq 0$ .

Now we consider the case of a piecewise smooth curve  $\gamma$ , and initially we do not assume that  $\gamma$  is closed. Then there is a succession of points  $a = t_0 < t_1 < \dots < t_{n-1} < t_n = b$  so that  $\gamma$  is continuously differentiable in every  $[t_{k-1}, t_k]$ . We consider an arbitrary fixed  $z_0 \notin \gamma^*$  and we define  $f(t) = \int_a^t \frac{\gamma'(s)}{\gamma(s) - z_0} ds$  for  $t \in [a, b]$ . Then  $f$  is continuous in  $[a, b]$  and differentiable at every point of continuity of  $\frac{\gamma'}{\gamma - z_0}$ . So in every  $(t_{k-1}, t_k)$  we have  $f'(t) = \frac{\gamma'(t)}{\gamma(t) - z_0}$  and hence  $\frac{d}{dt}((\gamma(t) - z_0)e^{-f(t)}) = \gamma'(t)e^{-f(t)} - (\gamma(t) - z_0)f'(t)e^{-f(t)} = 0$ . Thus,  $(\gamma(t) - z_0)e^{-f(t)}$  is constant in each  $(t_{k-1}, t_k)$  with a constant value which *a priori* depends on  $k$ . Since this function is continuous in  $[a, b]$ , it is constant in  $[a, b]$ . Hence there is  $c \in \mathbb{C}$  so that  $(\gamma(t) - z_0)e^{-f(t)} = c$

for every  $t \in [a, b]$ . Since  $c \neq 0$ , there is  $d \in \mathbb{C}$  so that  $e^d = c$ , and thus  $e^{f(t)+d} = \gamma(t) - z_0$  for every  $t \in [a, b]$ . Now we redefine  $f$  by adding to it the constant  $d$ , i.e. we write

$$f(t) = \int_a^t \frac{\gamma'(s)}{\gamma(s)-z_0} ds + d \quad \text{for every } t \in [a, b] \quad (6.3)$$

and we have  $e^{f(t)} = \gamma(t) - z_0$  for every  $t \in [a, b]$ . In other words, the function  $f$  is a *continuous branch* of  $\log(\gamma - z_0)$  in  $[a, b]$ . Now, the real part of  $f$  is  $\ln|\gamma - z_0|$  and, if we denote  $h$  the imaginary part of  $f$ , then  $h$  is a continuous branch of  $\arg(\gamma - z_0)$  in  $[a, b]$ .

We have that  $f(b) - f(a) = \int_a^b \frac{\gamma'(s)}{\gamma(s)-z_0} ds = \int_\gamma \frac{1}{\zeta - z_0} d\zeta$ . Since  $\Delta \arg(\gamma - z_0) = h(b) - h(a) = \text{Im}(f(b) - f(a))$ , we get

$$\Delta \arg(\gamma - z_0) = \text{Im} \int_\gamma \frac{1}{\zeta - z_0} d\zeta.$$

In the case of a closed curve we have the following result.

**Proposition 6.11.** *Let  $\gamma$  be a piecewise smooth closed curve in  $\mathbb{C} \setminus \{z_0\}$ . Then*

$$n(\gamma; z_0) = \frac{1}{2\pi i} \oint_\gamma \frac{1}{\zeta - z_0} d\zeta.$$

*Proof.* Using the notation of the previous discussion, we have  $n(\gamma; z_0) = \frac{\Delta \arg(\gamma - z_0)}{2\pi} = \frac{h(b) - h(a)}{2\pi}$ . Since  $\gamma$  is closed, we have that  $\ln|\gamma(b) - z_0| = \ln|\gamma(a) - z_0|$  and from (6.2) and (6.3) we get that  $n(\gamma; z_0) = \frac{f(b) - f(a)}{2\pi i} = \frac{1}{2\pi i} \int_a^b \frac{\gamma'(s)}{\gamma(s) - z_0} ds = \frac{1}{2\pi i} \oint_\gamma \frac{1}{\zeta - z_0} d\zeta$ .  $\square$

**Example 6.1.2.** We take  $n \in \mathbb{Z}$  and consider the closed curve  $\gamma : [0, 2\pi] \rightarrow \mathbb{C} \setminus \{z_0\}$  with parametric equation  $\gamma(t) = z_0 + re^{int}$ . It is clear that, if  $n \neq 0$  and  $t$  increases in the interval  $[0, 2\pi]$ , then  $\gamma(t)$  describes  $|n|$  times the circle  $C_{z_0}(r)$  in the positive direction, if  $n > 0$ , and in the negative direction, if  $n < 0$ . In the case  $n = 0$ , then  $\gamma(t)$  is constant and describes  $|n| = 0$  times the circle  $C_{z_0}(r)$ . All these agree with the result of the calculation:

$$n(\gamma; z_0) = \frac{1}{2\pi i} \oint_\gamma \frac{1}{\zeta - z_0} d\zeta = \frac{1}{2\pi i} \int_0^{2\pi} \frac{1}{re^{int}} rin e^{int} dt = n.$$

Since  $\gamma^* = C_{z_0}(r)$ , the complement of  $\gamma^*$  has two connected components: the disc  $D_{z_0}(r)$  and the ring  $D_{z_0}(r, +\infty)$ . Thus,  $n(\gamma; z) = n(\gamma; z_0) = n$  when  $z \in D_{z_0}(r)$ . Also, since  $D_{z_0}(r, +\infty)$  is the unbounded component of the complement of  $\gamma^*$ , we have that  $n(\gamma; z) = 0$  when  $z \in D_{z_0}(r, +\infty)$ .

**Proposition 6.12.** *Let  $\Omega \subseteq \mathbb{C} \setminus \{z_0\}$  be a region. Then a holomorphic branch of  $\log(z - z_0)$  exists in  $\Omega$  if and only if  $n(\gamma; z_0) = 0$  for every closed curve  $\gamma$  in  $\Omega$ .*

*Proof.* If  $n(\gamma; z_0) = 0$  for every closed curve  $\gamma$  in  $\Omega$ , then  $n(\gamma; z_0) = 0$  for every piecewise smooth closed curve  $\gamma$  in  $\Omega$ . Now, theorem 4.1 applied to  $g(z) = z - z_0$  implies that a holomorphic branch of  $\log(z - z_0)$  exists in  $\Omega$ . The converse is a corollary of proposition 6.8.  $\square$

Of course example 6.1.1 is relevant to proposition 6.12.

**Cauchy's formula for derivatives and closed curves in convex regions.** *If  $f$  is holomorphic in the convex region  $\Omega$  and  $\gamma$  is a piecewise smooth closed curve in  $\Omega$ , then for all  $n \in \mathbb{N}_0$  we have*

$$n(\gamma; z)f^{(n)}(z) = \frac{n!}{2\pi i} \oint_\gamma \frac{f(\zeta)}{(\zeta - z)^{n+1}} d\zeta \quad \text{for every } z \in \Omega \setminus \gamma^*.$$

*Proof.* The function  $F(\zeta) = \frac{f(\zeta) - f(z)}{\zeta - z}$  is holomorphic in  $\Omega \setminus \{z\}$ . Since  $z$  is a root of  $f(\zeta) - f(z)$ , the singularity  $z$  of  $F$  is removable. So we may define  $F$  at  $z$  with  $F(z) = \lim_{\zeta \rightarrow z} \frac{f(\zeta) - f(z)}{\zeta - z} = f'(z)$  and then  $F$  becomes holomorphic in  $\Omega$ . Now we apply the theorem of Cauchy in convex regions and get  $\oint_\gamma \frac{f(\zeta) - f(z)}{\zeta - z} d\zeta = \oint_\gamma F(\zeta) d\zeta = 0$  for every  $z \in \Omega \setminus \gamma^*$ . This implies

$$\oint_\gamma \frac{f(\zeta)}{\zeta - z} d\zeta = f(z) \oint_\gamma \frac{1}{\zeta - z} d\zeta = f(z) 2\pi i n(\gamma; z) \quad (6.4)$$

for every  $z \in \Omega \setminus \gamma^*$ . This is the result of the statement in the case  $n = 0$ .

Now, we consider a small disc  $D_z(r) \subseteq \mathbb{C} \setminus \gamma^*$ . This is possible, since  $z$  belongs to the open set  $\mathbb{C} \setminus \gamma^*$ . The disc  $D_z(r)$  is connected and hence it is contained in one connected component of  $\mathbb{C} \setminus \gamma^*$ . Therefore, the index  $n(\gamma; w)$  is a constant function of  $w$  in  $D_z(r)$ , i.e.  $n(\gamma; w) = n(\gamma; z)$  for every  $w \in D_z(r)$ . This implies that all derivatives of  $n(\gamma; w)$  vanish at  $z$  and so when we differentiate (6.4) we get  $n! \oint_{\gamma} \frac{f(\zeta)}{(\zeta-z)^{n+1}} d\zeta = f^{(n)}(z)2\pi i n(\gamma; z)$  for every  $n \geq 1$ .  $\square$

A particular instance of the last result is Cauchy's formula for derivatives and circles. Indeed, when the curve  $\gamma$  describes the circle  $C_{z_0}(R)$  once in the positive direction we have  $n(\gamma; z) = 1$  for all  $z \in D_{z_0}(R)$ . We originally proved the result in the case of a circle, using corollary 5.1. We now have a "new" proof using that  $z$  is a removable singularity of  $\frac{f(\zeta)-f(z)}{\zeta-z}$ . We have also introduced the notion of the index of a closed curve. This new proof together with the introduction of the notion of index allows us to generalize the case of a circle to the case of a more general piecewise smooth closed curve. There is still a restriction in the sense that the curve has to be contained in a *convex* region in which the function is holomorphic. This is because our proof is based on Cauchy's theorem in *convex* regions. In this chapter we shall replace this restriction on the region with a restriction on the curve.

### Exercises.

**6.1.1.** (i) Consider closed curves  $\gamma_1, \gamma_2$  and  $z$  not on their trajectories. Assume that there are successive points  $w_1^{(1)}, \dots, w_n^{(1)}, w_{n+1}^{(1)} = w_1^{(1)}$  of  $\gamma_1^*$  and successive points  $w_1^{(2)}, \dots, w_n^{(2)}, w_{n+1}^{(2)} = w_1^{(2)}$  of  $\gamma_2^*$  and curves  $\sigma_1, \dots, \sigma_n, \sigma_{n+1} = \sigma_1$  so that every  $\sigma_j$  goes from  $w_j^{(1)}$  to  $w_j^{(2)}$  and so that, for each  $j = 1, \dots, n$ , the part of  $\gamma_1$  between  $w_j^{(1)}, w_{j+1}^{(1)}$ , the part of  $\gamma_2$  between  $w_j^{(2)}, w_{j+1}^{(2)}$ ,  $\sigma_j$  and  $\sigma_{j+1}$  are all in a convex subregion  $D_j$  of  $\mathbb{C} \setminus \{z\}$ . Prove that  $n(\gamma_1; z) = n(\gamma_2; z)$ .

(ii) Take a point  $z$  and two halflines  $l, m$  with vertex  $z$ . Let  $A \in l, A \neq z$  and  $B \in m, B \neq z$ . Consider any curve  $\gamma_1$  from  $A$  to  $B$  in one of the two angular regions defined by  $l, m$  and any curve  $\gamma_2$  from  $B$  to  $A$  in the second angular region defined by  $l, m$ . Consider the closed curve  $\gamma = \gamma_1 + \gamma_2$ . Using appropriately a small circle with center  $z$ , prove that  $n(\gamma; z) = \pm 1$ .

**6.1.2.** If  $\gamma_1, \gamma_2$  are closed curves in  $\mathbb{C} \setminus \{0\}$  then  $\gamma_1\gamma_2$  is a closed curve in  $\mathbb{C} \setminus \{0\}$ . Prove that  $\Delta \arg(\gamma_1\gamma_2) = \Delta \arg \gamma_1 + \Delta \arg \gamma_2$ .

**6.1.3.** Let  $F \subseteq \mathbb{C}$  be closed and connected,  $\pm 1 \in F$  and  $\Omega = \mathbb{C} \setminus F$ . Prove that there is a holomorphic branch of  $\log \frac{z-1}{z+1}$  in  $\Omega$ . Prove also that there is a holomorphic branch of  $(z^2 - 1)^{1/2}$  in  $\Omega$ .

## 6.2 Homotopy.

Let  $\gamma_0, \gamma_1 : [a, b] \rightarrow \mathbb{C}$  be two curves. We say that  $\gamma_1$  is **homotopic** to  $\gamma_0$  if there is a continuous function  $F : [a, b] \times [0, 1] \rightarrow \mathbb{C}$  so that  $F(t, 0) = \gamma_0(t)$  and  $F(t, 1) = \gamma_1(t)$  for every  $t \in [a, b]$ . The function  $F$  is called a **homotopy** from  $\gamma_0$  to  $\gamma_1$ .

For each  $s \in [0, 1]$  the function  $\gamma_s : [a, b] \rightarrow \mathbb{C}$ , given by  $\gamma_s(t) = F(t, s)$  for  $t \in [a, b]$ , is continuous and hence it is a curve. We shall call it *intermediate curve* between  $\gamma_0$  and  $\gamma_1$ .

Since  $[a, b] \times [0, 1]$  is compact, the homotopy  $F$  is uniformly continuous. Thus for every  $\epsilon > 0$  there is  $\delta > 0$  so that  $|F(t', s') - F(t'', s'')| < \epsilon$  when  $\sqrt{(t' - t'')^2 + (s' - s'')^2} < \delta$ . Therefore, if  $|s' - s''| < \delta$  then we have  $|\gamma_{s'}(t) - \gamma_{s''}(t)| < \epsilon$  for every  $t \in [a, b]$ , i.e. the curves  $\gamma_{s'}$  and  $\gamma_{s''}$  are uniformly close. We see that *when  $s$  increases in  $[0, 1]$  the curves  $\gamma_s$  form a continuously varying family of curves, starting with  $\gamma_0$  and ending with  $\gamma_1$* . To be more precise, we have a mapping  $[0, 1] \ni s \mapsto \gamma_s \in C([a, b])$ , which is continuous from  $[0, 1]$  with the euclidean distance to  $C([a, b])$  with the uniform distance:

$$|s' - s''| < \delta \quad \Rightarrow \quad \sup_{t \in [a, b]} |\gamma_{s'}(t) - \gamma_{s''}(t)| < \epsilon.$$

If all curves  $\gamma_s$  are closed, i.e. if  $F(a, s) = F(b, s)$  for every  $s \in [0, 1]$ , then we say that  $F$  is a **homotopy with closed intermediate curves**. If all curves  $\gamma_s$  have the same initial endpoint and the same final endpoint, i.e. if  $F(a, s)$  is constant and  $F(b, s)$  is constant for  $s \in [0, 1]$ , then we say that  $F$  is a **homotopy with fixed endpoints**.

If all curves  $\gamma_s$  are in the same set  $A$ , then we say that  $F$  is a homotopy in  $A$ .

We may define a relation between curves in a set  $A$ : we write  $\gamma_0 \equiv \gamma_1$  if there is a homotopy in  $A$  from  $\gamma_0$  to  $\gamma_1$ . It is easy to see that this is an equivalence relation:

(i) Every curve  $\gamma : [a, b] \rightarrow A$  is homotopic to itself through the homotopy  $F : [a, b] \times [0, 1] \rightarrow A$  given by  $F(t, s) = \gamma(t)$ .

(ii) If  $F : [a, b] \times [0, 1] \rightarrow A$  is a homotopy from  $\gamma_0$  to  $\gamma_1$ , i.e. if  $F(t, 0) = \gamma_0(t)$  and  $F(t, 1) = \gamma_1(t)$  for  $t \in [a, b]$ , then the function  $G : [a, b] \times [0, 1] \rightarrow A$  given by  $G(t, s) = F(t, 1-s)$  is a homotopy from  $\gamma_1$  to  $\gamma_0$ . In fact  $G$  is continuous and  $G(t, 0) = \gamma_1(t)$  and  $G(t, 1) = \gamma_0(t)$  for  $t \in [a, b]$ .

(iii) If  $F : [a, b] \times [0, 1] \rightarrow A$  is a homotopy from  $\gamma_0$  to  $\gamma_1$ , i.e. if  $F(t, 0) = \gamma_0(t)$  and  $F(t, 1) = \gamma_1(t)$  for  $t \in [a, b]$ , and if  $G : [a, b] \times [0, 1] \rightarrow A$  is a homotopy from  $\gamma_1$  to  $\gamma_2$ , i.e. if  $G(t, 0) = \gamma_1(t)$  and  $G(t, 1) = \gamma_2(t)$  for  $t \in [a, b]$ , then  $H : [a, b] \times [0, 1] \rightarrow A$ , given by

$$H(t, s) = \begin{cases} F(t, 2s), & t \in [a, b], s \in [0, \frac{1}{2}] \\ G(t, 2s - 1), & t \in [a, b], s \in [\frac{1}{2}, 1] \end{cases}$$

is a homotopy from  $\gamma_0$  to  $\gamma_2$ . Indeed,  $H$  is continuous and  $H(t, 0) = \gamma_0(t)$  and  $H(t, 1) = \gamma_2(t)$  for  $t \in [a, b]$ .

Furthermore, the previous argument shows that the relation of homotopy with closed intermediate curves and the relation of homotopy with fixed endpoints are both equivalence relations.

**Example 6.2.1.** If the set  $A$  is convex, every two curves in  $A$  are homotopic in  $A$ . Indeed, let  $\gamma_0, \gamma_1 : [a, b] \rightarrow A$  be two curves in  $A$ . Since  $\gamma_0(t), \gamma_1(t) \in A$  and  $A$  is convex, the linear segment  $[\gamma_0(t), \gamma_1(t)]$  is contained in  $A$ . Now, if we define  $F : [a, b] \times [0, 1] \rightarrow \mathbb{C}$  by

$$F(t, s) = (1-s)\gamma_0(t) + s\gamma_1(t),$$

then  $F$  is continuous and all its values are in  $A$ . Moreover,  $F(t, 0) = \gamma_0(t)$  and  $F(t, 1) = \gamma_1(t)$  for  $t \in [a, b]$ . Therefore,  $F$  is a homotopy in  $A$  from  $\gamma_0$  to  $\gamma_1$ . It is easy to see that, if  $\gamma_0$  and  $\gamma_1$  are closed, then all intermediate curves are closed. Also, if  $\gamma_0$  and  $\gamma_1$  have the same initial endpoint and the same final endpoint, then all intermediate curves have the same initial endpoint and the same final endpoint.

**Proposition 6.13.** *Let  $\gamma_0, \gamma_1$  be two closed curves in  $\mathbb{C} \setminus \{z\}$ . If there is a homotopy in  $\mathbb{C} \setminus \{z\}$ , with closed intermediate curves, between  $\gamma_0$  and  $\gamma_1$ , then  $n(\gamma_0; z) = n(\gamma_1; z)$ .*

*Proof.* Let  $F : [a, b] \times [0, 1] \rightarrow \mathbb{C} \setminus \{z\}$  be a homotopy with closed intermediate curves, between  $\gamma_0$  and  $\gamma_1$ . Since  $F$  is continuous and  $[a, b] \times [0, 1]$  is compact and  $F$  does not take the value  $z$ , there is  $\epsilon > 0$  so that  $|F(t, s) - z| \geq \epsilon$  for every  $t \in [a, b]$  and  $s \in [0, 1]$ . Also, since  $F$  is uniformly continuous, there is  $\delta > 0$  so that  $|s' - s''| < \delta$  implies  $|\gamma_{s'}(t) - \gamma_{s''}(t)| < \epsilon$  for every  $t \in [a, b]$ , where  $\gamma_s$  is the intermediate curve corresponding to  $s \in [0, 1]$ . Then we have  $|\gamma_{s'}(t) - \gamma_{s''}(t)| < |\gamma_{s''}(t) - z|$  for every  $t \in [a, b]$  and proposition 6.9 implies that  $n(\gamma_{s'}; z) = n(\gamma_{s''}; z)$ . Now we take successive points  $0 = s_0 < s_1 < \dots < s_{n-1} < s_n = 1$  so that  $s_k - s_{k-1} < \delta$  for every  $k = 1, \dots, n$ . Then we have  $n(\gamma_{s_{k-1}}; z) = n(\gamma_{s_k}; z)$  for every  $k = 1, \dots, n$  and hence  $n(\gamma_0; z) = n(\gamma_1; z)$ .  $\square$

**Proposition 6.14.** *Let  $f$  be holomorphic in the open set  $\Omega$ .*

(i) *If  $\gamma_0, \gamma_1$  are piecewise smooth curves in  $\Omega$  with the same initial endpoint and the same final endpoint and if there is a homotopy in  $\Omega$ , with fixed endpoints, between  $\gamma_0$  and  $\gamma_1$ , then  $\int_{\gamma_0} f(z) dz = \int_{\gamma_1} f(z) dz$ .*

(ii) *If  $\gamma_0, \gamma_1$  are piecewise smooth closed curves in  $\Omega$  and if there is a homotopy in  $\Omega$ , with closed intermediate curves, between  $\gamma_0$  and  $\gamma_1$ , then  $\oint_{\gamma_0} f(z) dz = \oint_{\gamma_1} f(z) dz$ .*



*Proof.* (i) Let  $F : [a, b] \times [0, 1] \rightarrow \Omega$  be the homotopy in  $\Omega$  from  $\gamma_0$  to  $\gamma_1$ . Then the subset  $F([a, b] \times [0, 1])$  of  $\Omega$  is compact and there is  $\epsilon > 0$  so that

$$|z - w| \geq \epsilon \quad \text{for every } z \in F([a, b] \times [0, 1]) \text{ and every } w \in \Omega^c. \quad (6.5)$$

Since  $F$  is uniformly continuous, there is  $\delta > 0$  so that

$$|F(t', s') - F(t'', s'')| < \epsilon \quad \text{if } |t' - t''| < \delta \text{ and } |s' - s''| < \delta. \quad (6.6)$$

We take points  $a = t_0 < t_1 < \dots < t_{n-1} < t_n = b$  and  $0 = s_0 < s_1 < \dots < s_{m-1} < s_m = 1$  so that  $t_k - t_{k-1} < \delta$  and  $s_l - s_{l-1} < \delta$  for all  $k$  and  $l$ . Then (6.5) and (6.6) imply that every rectangle  $[t_{k-1}, t_k] \times [s_{l-1}, s_l]$  is mapped by  $F$  in the disc  $D_{F(t_{k-1}, s_{l-1})}(\epsilon)$  which is contained in  $\Omega$ . Since  $f$  is holomorphic in this disc, its curvilinear integral over any closed curve in this disc is equal to 0.

Now we denote  $\gamma_{0,k}$  and  $\gamma_{1,k}$  the restrictions of  $\gamma_0$  and  $\gamma_1$  in  $[t_{k-1}, t_k]$ . We also denote  $\sigma_{k,l}$  the linear segment  $[F(t_{k-1}, s_l), F(t_k, s_l)]$  for  $k = 1, \dots, n$  and  $l = 1, \dots, m-1$ . Finally, we denote  $\rho_{k,l}$  the linear segment  $[F(t_k, s_{l-1}), F(t_k, s_l)]$  for  $k = 0, \dots, n$  and  $l = 1, \dots, m$ . Then for every  $k = 1, \dots, n$  we have

$$\begin{aligned} \int_{\gamma_{0,k}} f(z) dz - \int_{\sigma_{k,1}} f(z) dz &= \int_{\rho_{k-1,1}} f(z) dz - \int_{\rho_{k,1}} f(z) dz \\ \int_{\sigma_{k,l-1}} f(z) dz - \int_{\sigma_{k,l}} f(z) dz &= \int_{\rho_{k-1,l}} f(z) dz - \int_{\rho_{k,l}} f(z) dz \quad \text{for } l = 2, \dots, m-1 \\ \int_{\sigma_{k,m-1}} f(z) dz - \int_{\gamma_{1,k}} f(z) dz &= \int_{\rho_{k-1,m}} f(z) dz - \int_{\rho_{k,m}} f(z) dz. \end{aligned}$$

Adding these  $m$  equalities and then adding for  $k = 1, \dots, n$  and considering cancellations, we find

$$\int_{\gamma_0} f(z) dz - \int_{\gamma_1} f(z) dz = \sum_{l=1}^m \int_{\rho_{0,l}} f(z) dz - \sum_{l=1}^m \int_{\rho_{n,l}} f(z) dz. \quad (6.7)$$

Since all intermediate curves have the same initial endpoint and the same final endpoint, we see that all linear segments  $\rho_{0,l}$  and  $\rho_{n,l}$  are single point sets and hence all integrals in the right side of (6.7) are equal to 0. Thus,  $\int_{\gamma_0} f(z) dz = \int_{\gamma_1} f(z) dz$ .

(ii) Since all intermediate curves are closed, we have  $F(a, s) = F(b, s)$  for every  $s \in [0, 1]$ . Therefore, for each  $l$  the linear segments  $\rho_{0,l}$  and  $\rho_{n,l}$  coincide and again the right side of (6.7) is equal to 0. Thus,  $\oint_{\gamma_0} f(z) dz = \oint_{\gamma_1} f(z) dz$ .  $\square$

### Exercises.

**6.2.1.** Let  $A$  be arcwise connected and  $\gamma_1(t) = z_1$  and  $\gamma_2(t) = z_2$  be two constant curves in  $A$ . If a curve  $\gamma$  is homotopic in  $A$  to  $\gamma_1$ , prove that  $\gamma$  is homotopic in  $A$  to  $\gamma_2$ .

**6.2.2.** If  $\gamma$  is a closed curve in  $\mathbb{C} \setminus \{0\}$ , prove that  $\gamma$  is homotopic in  $\mathbb{C} \setminus \{0\}$  to a closed curve whose trajectory is contained in the circle  $\mathbb{T}$ .

**6.2.3.** (i) Let  $f$  be continuous in  $\overline{D_0}(R)$ . We define  $\gamma(t) = f(Re^{it})$  for every  $t \in [0, 2\pi]$ . Prove that, if  $n(\gamma; w) \neq 0$ , then  $w \in f(D_0(R))$ . I.e.  $\{w \mid w \text{ is surrounded by } \gamma\} \subseteq f(D_0(R))$ .

(ii) Using the result of (i), prove the fundamental theorem of algebra.

**6.2.4.** Let  $p \in A$  and let  $\mathcal{M}_p(A)$  be the set of all closed curves with both of their endpoints at  $p$ . If  $\gamma_1, \gamma_2 \in \mathcal{M}_p(A)$ , then clearly  $\gamma_1 + \gamma_2 \in \mathcal{M}_p(A)$ . Also, if  $\gamma \in \mathcal{M}_p(A)$ , then  $-\gamma \in \mathcal{M}_p(A)$ .

(i) Prove that the relation of homotopy in  $A$  with closed intermediate curves and fixed endpoints ( $= p$ ) is an equivalence relation in  $\mathcal{M}_p(A)$ . The set of all equivalence classes is denoted  $\mathcal{H}_p(A) = \{[\gamma] \mid \gamma \in \mathcal{M}_p(A)\}$ .

(ii) If  $\gamma, \gamma_1, \gamma_2 \in \mathcal{M}_p(A)$ , we define  $[\gamma_1] + [\gamma_2] = [\gamma_1 + \gamma_2]$  and  $-[\gamma] = [-\gamma]$ . Prove that these are well-defined and that  $\mathcal{H}_p(A)$  with these operations is a group, whose neutral element is  $[\gamma_p]$ ,

where  $\gamma_p$  is the constant curve  $p$ .

(iii) If  $A$  is arcwise connected, prove that for every  $p, q \in A$  the groups  $\mathcal{H}_p(A)$  and  $\mathcal{H}_q(A)$  are isomorphic. In this case we write  $\mathcal{H}(A)$ . (See exercise 6.2.1.)

(iv) Prove that  $\mathcal{H}(\mathbb{C}) \cong \{0\}$ ,  $\mathcal{H}(\mathbb{C} \setminus \{0\}) \cong \mathbb{Z}$ ,  $\mathcal{H}(\mathbb{T}) \cong \mathbb{Z}$ .

**6.2.5.** Let  $z_1, z_2, z_3, w_1, w_2, w_3$  be distinct points. Is it possible to join every  $z_k$  with every  $w_j$  with simple curves  $\gamma_{kj}$  whose trajectories are mutually disjoint?

### 6.3 Combinatorial results for curves and square nets.

**Lemma 6.1.** *Let  $\Sigma = \{\sigma_1, \dots, \sigma_n\}$  be a set of curves (not necessarily closed) and let  $A = \{a_1, \dots, a_m\}$  be the set of their endpoints ( $m \leq 2n$ ). We assume that for every point of  $A$  the number of the curves in  $\Sigma$  that arrive at this point is the same as the number of the curves in  $\Sigma$  that leave from this point. Then we can partition  $\Sigma$  into subsets  $\Sigma_1, \dots, \Sigma_k$  so that each  $\Sigma_j$  consists of successive curves and the sum  $\gamma_j$  of the curves in  $\Sigma_j$  is a closed curve.*

*Proof.* We describe the following algorithm for the partitioning of  $\Sigma$ .

We start with  $\sigma_1$ . The final endpoint of  $\sigma_1$  is the initial endpoint of at least one curve in  $\Sigma$ . If the final endpoint of  $\sigma_1$  coincides with its initial endpoint, then  $\sigma_1$  is closed and we stop the process. If this is not the case, then, renumbering if necessary the curves  $\sigma_2, \dots, \sigma_n$ , we may assume that the final endpoint of  $\sigma_1$  coincides with the initial endpoint of  $\sigma_2$ . If the final endpoint of  $\sigma_2$  coincides with the initial endpoint of  $\sigma_1$ , then the sum of  $\sigma_1, \sigma_2$  is a closed curve and we stop the process. If the final endpoint of  $\sigma_2$  coincides with its initial endpoint, then  $\sigma_2$  is a closed curve and we stop the process. If the final endpoint of  $\sigma_2$  is not the initial point of either  $\sigma_1$  or  $\sigma_2$ , then renumbering if necessary the curves  $\sigma_3, \dots, \sigma_n$ , we may assume that the final endpoint of  $\sigma_2$  coincides with the initial endpoint of  $\sigma_3$ . Then, exactly as before, we examine whether the final endpoint of  $\sigma_3$  coincides with the initial endpoint of  $\sigma_1$  or of  $\sigma_2$  or of  $\sigma_3$ . Then, respectively, the sum of  $\sigma_1, \sigma_2, \sigma_3$  or the sum of  $\sigma_2, \sigma_3$  or  $\sigma_3$  by itself is a closed curve and we stop the process. If the final endpoint of  $\sigma_3$  is not the initial endpoint of either  $\sigma_1$  or  $\sigma_2$  or  $\sigma_3$ , then renumbering if necessary the curves  $\sigma_4, \dots, \sigma_n$ , we may assume that the final endpoint of  $\sigma_3$  coincides with the initial endpoint of  $\sigma_4$ . Now, it is clear that this process will eventually stop, because we have only finitely many curves. Therefore, we shall eventually find successive curves  $\sigma_1, \sigma_2, \dots, \sigma_{k-1}, \sigma_k$  ( $1 \leq k \leq n$ ) so that the final endpoint of  $\sigma_k$  coincides with the initial endpoint of one of the same curves  $\sigma_1, \sigma_2, \dots, \sigma_{k-1}, \sigma_k$ . Let the final endpoint of  $\sigma_k$  coincide with the initial endpoint of  $\sigma_l$  for some  $l$  with  $1 \leq l \leq k$ . Then the sum of  $\sigma_l, \sigma_{l+1}, \dots, \sigma_{k-1}, \sigma_k$  is a closed curve and we stop the process. Now we set  $\Sigma_1 = \{\sigma_l, \sigma_{l+1}, \dots, \sigma_{k-1}, \sigma_k\}$  and call  $\gamma_1$  the closed curve which is the sum of  $\sigma_l, \sigma_{l+1}, \dots, \sigma_{k-1}, \sigma_k$ . Then we drop the curves of  $\Sigma_1$  from  $\Sigma$ , i.e. we consider the set  $\Sigma' = \Sigma \setminus \Sigma_1 = \{\sigma_1, \dots, \sigma_{l-1}, \sigma_{k+1}, \dots, \sigma_n\}$ .

Each endpoint of the curves in  $\Sigma'$  is one of the points of  $A = \{a_1, \dots, a_m\}$ , say it is  $a_j$ . Then the number of the curves in  $\Sigma$  that arrive at  $a_j$  is the same as the number of the curves in  $\Sigma$  that leave from  $a_j$ . But the curves  $\sigma_l, \sigma_{l+1}, \dots, \sigma_{k-1}, \sigma_k$  are successive and hence if one of them arrives at  $a_j$  then the next one leaves from  $a_j$ . Therefore, the remaining curves of  $\Sigma'$  have the same property: the number of the curves in  $\Sigma'$  that arrive at  $a_j$  is the same as the number of the curves in  $\Sigma'$  that leave from  $a_j$ . Thus  $\Sigma'$  has the same property as the original  $\Sigma$ .

Now we continue our algorithm with  $\Sigma'$ . We find a subset  $\Sigma_2$  of  $\Sigma'$  which consists of successive curves and we call  $\gamma_2$  the closed curve which is the sum of the curves in  $\Sigma_2$ . Then we drop the curves of  $\Sigma_2$  from  $\Sigma'$ , i.e. we consider the set  $\Sigma'' = \Sigma' \setminus \Sigma_2 = \Sigma \setminus (\Sigma_1 \cup \Sigma_2)$ . We go on until we exhaust the original  $\Sigma$ .  $\square$

**Lemma 6.2.** *We take any  $\delta > 0$  and two perpendicular lines. For each of them we consider all its parallel lines at distances equal to integer multiples of  $\delta$ . The result is a net of closed square regions of sidelength  $\delta$  which cover the plane and have disjoint interiors. We choose any of those closed*

square regions, say  $Q_1, \dots, Q_l$ . We consider the closed boundary curves  $\partial Q_1, \dots, \partial Q_l$  with their positive direction. Each of them is the sum of four corresponding linear segments, considered as curves with the same direction. We drop the linear segments (with necessarily opposite directions) which are common to any two neighboring square regions from among the  $Q_1, \dots, Q_l$  and we consider the set  $\Sigma = \{\sigma_1, \dots, \sigma_n\}$  of all the remaining linear segments, i.e. those which belong to only one of  $Q_1, \dots, Q_l$ . Then we can partition  $\Sigma$  into subsets  $\Sigma_1, \dots, \Sigma_k$  so that each  $\Sigma_j$  consists of successive linear segments and the sum  $\gamma_j$  of the linear segments in  $\Sigma_j$  is a closed curve.

*Proof.* It is enough to prove that  $\Sigma$  has the property described in lemma 6.1, i.e. that for every point of intersection  $a$  of our lines the number of the curves in  $\Sigma$  that arrive at  $a$  is the same as the number of the curves in  $\Sigma$  that leave from  $a$ . This can be done easily, considering cases for the number, 0 or 1 or 2 or 3 or 4, of the squares among  $Q_1, \dots, Q_l$  which have  $a$  as one of their corners.  $\square$

**Lemma 6.3.** *Let  $\Omega$  be an open set and  $K \subseteq \Omega$  be compact. Then there is  $\delta > 0$  so that  $|z-w| \geq 2\delta$  for every  $z \in K$  and every  $w \in \Omega^c$ . For this  $\delta > 0$  we consider the net of closed square regions of lemma 6.2 and we take all closed square regions  $Q_1, \dots, Q_l$  of the net which intersect  $K$ . Then  $Q_1, \dots, Q_l$  are contained in  $\Omega$ . As in lemma 6.2, we consider the set  $\Sigma = \{\sigma_1, \dots, \sigma_n\}$  of all linear segments which belong to only one of  $Q_1, \dots, Q_l$  and we partition  $\Sigma$  into subsets  $\Sigma_1, \dots, \Sigma_k$  so that each  $\Sigma_j$  consists of successive linear segments and the sum  $\gamma_j$  of the linear segments in  $\Sigma_j$  is a closed curve. Then  $\gamma_1^* \cup \dots \cup \gamma_k^* \subseteq \Omega \setminus K$  (i.e. every  $\gamma_j$  is in  $\Omega \setminus K$ ) and the distance of every  $z \in \gamma_1^* \cup \dots \cup \gamma_k^*$  from  $K$  is  $\leq \delta\sqrt{2}$ . Moreover, for every  $f$  holomorphic in  $\Omega$  and for every  $z \in K$ , we have*

$$f(z) = \sum_{j=1}^k \frac{1}{2\pi i} \oint_{\gamma_j} \frac{f(\zeta)}{\zeta-z} d\zeta. \quad (6.8)$$

*Proof.* Each of the closed square regions  $Q_1, \dots, Q_l$ , say  $Q_m$ , intersects  $K$  and its diameter is equal to  $\sqrt{2}\delta$ . Therefore, the distance of every point of  $Q_m$  from  $K$  is  $\leq \sqrt{2}\delta$ . Since  $\sqrt{2}\delta < 2\delta$ , we see that  $Q_m$  is contained in  $\Omega$ .

Consider any of the linear segments  $\sigma_1, \dots, \sigma_n$ , say  $\sigma_j$ . Now,  $\sigma_j$  belongs to one of  $Q_1, \dots, Q_l$ , say  $Q_m$ . Since  $Q_m$  is contained in  $\Omega$ , we have that  $\sigma_j$  is also contained in  $\Omega$ . Moreover, the distance of every point of  $Q_m$  from  $K$  is  $\leq \sqrt{2}\delta$  and hence the distance of every point of  $\sigma_j$  from  $K$  is  $\leq \sqrt{2}\delta$ . If  $\sigma_j$  intersects  $K$ , then both closed square regions of our net which lie on the two sides of  $\sigma_j$  intersect  $K$  and hence both are among  $Q_1, \dots, Q_l$ . This is impossible because  $\sigma_j$  belongs to only one of  $Q_1, \dots, Q_l$ . Therefore,  $\sigma_j$  does not intersect  $K$  and hence it is contained in  $\Omega \setminus K$ . Finally, since each of  $\gamma_1, \dots, \gamma_k$  is the sum of certain of the  $\sigma_1, \dots, \sigma_n$ , we have proved that  $\gamma_1^* \cup \dots \cup \gamma_k^* \subseteq \Omega \setminus K$  and that the distance of every  $z \in \gamma_1^* \cup \dots \cup \gamma_k^*$  from  $K$  is  $\leq \delta\sqrt{2}$ .

Now we take any  $z \in K$ . Then  $z$  belongs to one  $Q_1, \dots, Q_l$ , say  $Q_m$ . Let us assume that  $z$  is an interior point of  $Q_m$ . Since the closed square region  $Q_m$  is contained in  $\Omega$ , there is a slightly larger open square region  $Q'$  which is also contained in  $\Omega$ . Now  $f$  is holomorphic in the convex region  $Q'$  and Cauchy's formula in section 6.1 says that

$$f(z) = \frac{1}{2\pi i} \oint_{\partial Q_m} \frac{f(\zeta)}{\zeta-z} d\zeta, \quad (6.9)$$

because the index of  $\partial Q_m$  with respect to  $z$  is equal to 1. Now we take any closed square region  $Q_p$  with  $p \neq m$ . Then  $z$  is not contained in  $Q_p$  and again we may find an open square region  $Q'$  slightly larger than  $Q_p$  which is contained in  $\Omega$  and which does not contain  $z$ . Then  $\frac{f(\zeta)}{\zeta-z}$  is a holomorphic function of  $\zeta$  in the convex region  $Q'$  and hence

$$0 = \frac{1}{2\pi i} \oint_{\partial Q_p} \frac{f(\zeta)}{\zeta-z} d\zeta \quad \text{for } p \neq m. \quad (6.10)$$

We add (6.9) and (6.10) and we get

$$f(z) = \sum_{p=1}^l \frac{1}{2\pi i} \oint_{\partial Q_p} \frac{f(\zeta)}{\zeta-z} d\zeta. \quad (6.11)$$

Now we split the integral over each  $\partial Q_p$  in four integrals over the boundary linear segments of  $\partial Q_p$  and we get  $4l$  integrals. We recall that if a linear segment belongs to two neighboring closed square regions, then it appears twice among the integrals with opposite directions and hence the two integrals cancel. Therefore, the remaining integrals will be only over the boundary linear segments which belong to exactly one of  $Q_1, \dots, Q_l$ , i.e. the linear segments of the set  $\Sigma = \{\sigma_1, \dots, \sigma_n\}$ . Thus (6.11) becomes

$$f(z) = \sum_{\sigma \in \Sigma} \frac{1}{2\pi i} \int_{\sigma} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

The subsets  $\Sigma_1, \dots, \Sigma_k$  form a partition of  $\Sigma$  and hence

$$f(z) = \sum_{j=1}^k \sum_{\sigma \in \Sigma_j} \frac{1}{2\pi i} \int_{\sigma} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

Finally, since  $\gamma_j$  is the sum of the successive linear segments  $\sigma \in \Sigma_j$ , we end up with (6.8). Now let  $z$  be a boundary point of  $Q_m$ . Then we may consider a variable point  $z'$  in the interior of  $Q_m$  so that  $z' \rightarrow z$ . We have proved (6.8) for  $z'$ , i.e.

$$f(z') = \sum_{j=1}^k \frac{1}{2\pi i} \oint_{\gamma_j} \frac{f(\zeta)}{\zeta - z'} d\zeta.$$

Proposition 4.12 implies the continuity of the right side as a function of  $z'$ . Therefore, taking the limit as  $z' \rightarrow z$ , we end up again with (6.8).  $\square$

## 6.4 The theorem of Cauchy in general open sets.

Let  $\sigma_1, \dots, \sigma_n$  be any curves (not necessarily closed) and  $k_1, \dots, k_n$  be any integers (not necessarily non-negative). Then we say that the curves  $\sigma_1, \dots, \sigma_n$  considered  $k_1, \dots, k_n$  times, respectively, form a **chain**  $\Sigma$ . The integer  $k_j$  is called **multiplicity** of the corresponding  $\sigma_j$  in the chain  $\Sigma$ . If every  $\sigma_j$  is closed, then  $\Sigma$  is called **closed chain** or **cycle**. If every  $\sigma_j$  is in a set  $A$ , then we say that  $\Sigma$  is in  $A$ .

If a curve  $\sigma$  is not among the curves which constitute a chain  $\Sigma$ , we may include it among those curves by assigning multiplicity 0 to  $\sigma$ . And now we may introduce the algebraic structure of a *module* in the set of all chains in the following manner. If  $\Sigma'$  and  $\Sigma''$  are two chains, we may assume that they are formed by the same collection  $\sigma_1, \dots, \sigma_n$  of curves. If  $k'_1, \dots, k'_n$  and  $k''_1, \dots, k''_n$  are the corresponding multiplicities in the chains  $\Sigma'$  and  $\Sigma''$ , then we define  $\Sigma' + \Sigma''$  to be the chain which consists of  $\sigma_1, \dots, \sigma_n$  with multiplicities  $k'_1 + k''_1, \dots, k'_n + k''_n$ . Moreover, if  $k$  is an integer and  $\Sigma$  is a chain formed by the curves  $\sigma_1, \dots, \sigma_n$  with multiplicities  $k_1, \dots, k_n$ , then we define  $k\Sigma$  to be the chain formed by  $\sigma_1, \dots, \sigma_n$  with multiplicities  $kk_1, \dots, kk_n$ . It is very easy to show that, under this addition of chains and this multiplication of chains and integers, the set of chains is a  $\mathbb{Z}$ -module. The opposite  $-\Sigma$  of a chain  $\Sigma$  is  $(-1)\Sigma$  and the neutral element of addition is the chain which contains no curve (or any curves with multiplicities 0).

If  $\Sigma$  is a chain formed by the curves  $\sigma_1, \dots, \sigma_n$  with multiplicities  $k_1, \dots, k_n$ , we immediately see that, under the above definitions of addition and multiplication, we have  $\Sigma = k_1\sigma_1 + \dots + k_n\sigma_n$ . Here we consider each  $\sigma_j$  as a chain consisting of only one curve with multiplicity 1.

We shall not go into this algebraic point of view, since it does not have much to offer in our study of complex analysis. We shall keep in mind, though, the definition and notation of  $\Sigma' + \Sigma''$  and  $k\Sigma$  and from time to time we shall feel free to make certain mild algebraic comments.

Now we consider a chain  $\Sigma$  formed by the piecewise smooth curves  $\sigma_1, \dots, \sigma_n$  with multiplicities  $k_1, \dots, k_n$  and a continuous  $\phi : \sigma_1^* \cup \dots \cup \sigma_n^* \rightarrow \mathbb{C}$ . We define the **curvilinear integral** of  $\phi$  over  $\Sigma$  by

$$\int_{\Sigma} \phi(z) dz = \sum_{j=1}^n k_j \int_{\sigma_j} \phi(z) dz.$$

If  $\Sigma$  is a cycle, we may use the notation

$$\oint_{\Sigma} \phi(z) dz.$$

It is easy to show that

$$\int_{k_1 \Sigma_1 + k_2 \Sigma_2} \phi(z) dz = k_1 \int_{\Sigma_1} \phi(z) dz + k_2 \int_{\Sigma_2} \phi(z) dz.$$

This says that integration “respects” the linear structure of the  $\mathbb{Z}$ -module of chains.

Moreover, for every  $z$  which does not belong to  $\sigma_1^* \cup \dots \cup \sigma_n^*$  we define the **rotation number** or **index** of the chain  $\Sigma$  with respect to  $z$  by

$$n(\Sigma; z) = \sum_{j=1}^n k_j n(\sigma_j; z).$$

We may say that  $n(\Sigma; z)$  is the total number of rotations around  $z$  of the curves forming  $\Sigma$ , taking into account their multiplicities.

Combining the last two definitions, we easily see that the index of a cycle consisting of piecewise smooth closed curves is given by the same integral form which gives the index of a piecewise smooth closed curve:

$$n(\Sigma; z) = \frac{1}{2\pi i} \oint_{\Sigma} \frac{1}{\zeta - z} d\zeta.$$

$$\text{Indeed, } n(\Sigma; z) = \sum_{j=1}^n k_j n(\sigma_j; z) = \sum_{j=1}^n k_j \frac{1}{2\pi i} \oint_{\sigma_j} \frac{1}{\zeta - z} d\zeta = \frac{1}{2\pi i} \oint_{\Sigma} \frac{1}{\zeta - z} d\zeta.$$

Now we state a basic definition.

Let  $\Sigma$  be a cycle in the open set  $\Omega$ . We say that  $\Sigma$  is **null-homologous** in  $\Omega$  if  $n(\Sigma; z) = 0$  for every  $z \in \Omega^c$ .

In other words, a cycle  $\Sigma$  in  $\Omega$  is null-homologous in  $\Omega$  if the total number of rotations of the curves forming  $\Sigma$ , taking into account their multiplicities, around every point of the complement of  $\Omega$  is zero.

**Lemma 6.4.** *Let  $\Omega \subseteq \mathbb{C}$  be open,  $\delta > 0$  and  $K = \{z \in \Omega \mid |z| \leq R, |z - w| \geq \delta \text{ for all } w \in \Omega^c\}$ . Then  $K$  is a compact subset of  $\Omega$ .*

*Proof.* Since  $K \subseteq \overline{D}_0(R)$ , the set  $K$  is bounded.

Now, let  $z_n \in K$  for every  $n$  and  $z_n \rightarrow z$ . If we prove that  $z \in K$ , then  $K$  is closed and hence compact. We have  $|z_n| \leq R$  for every  $n$  and thus  $|z| \leq R$ . For every  $w \in \Omega^c$  we have  $|z_n - w| \geq \delta$  for every  $n$  and hence  $|z - w| \geq \delta$ . Therefore  $z \in K$ .  $\square$

**The theorem of Cauchy in general open sets.** *If  $f$  is holomorphic in the open set  $\Omega$  and if the cycle  $\Sigma$ , consisting of piecewise smooth closed curves, is null-homologous in  $\Omega$ , then*

$$\oint_{\Sigma} f(z) dz = 0.$$

*Proof.* Let the cycle  $\Sigma$  consist of the piecewise smooth closed curves  $\sigma_1, \dots, \sigma_n$  with multiplicities  $k_1, \dots, k_n$ . Since  $\sigma_1^* \cup \dots \cup \sigma_n^*$  is a compact subset of  $\Omega$ , there is  $\delta > 0$  so that every point of  $\sigma_1^* \cup \dots \cup \sigma_n^*$  has a distance  $\geq 2\delta$  from  $\Omega^c$  and there is  $R > 0$  so that  $\sigma_1^* \cup \dots \cup \sigma_n^*$  is contained in the closed disc  $\overline{D}_0(R)$ . We consider the set

$$K = \{z \in \Omega \mid |z| \leq R, |z - w| \geq 2\delta \text{ for every } w \in \Omega^c\}$$

of lemma 6.4 (with  $2\delta$  instead of  $\delta$ ). Then  $K$  is a compact subset of  $\Omega$  which contains  $\sigma_1^* \cup \dots \cup \sigma_n^*$ . With the same  $\delta$  and with this set  $K$  we form the closed curves  $\gamma_1, \dots, \gamma_k$  in  $\Omega \setminus K$ , which are described in lemma 6.3. According to lemma 6.3 we have

$$f(z) = \sum_{l=1}^k \frac{1}{2\pi i} \oint_{\gamma_l} \frac{f(\zeta)}{\zeta - z} d\zeta \quad \text{for every } z \in \sigma_1^* \cup \dots \cup \sigma_n^*.$$

Hence

$$\begin{aligned} \oint_{\Sigma} f(z) dz &= \sum_{j=1}^n k_j \oint_{\sigma_j} f(z) dz = \sum_{j=1}^n k_j \oint_{\sigma_j} \left( \sum_{l=1}^k \frac{1}{2\pi i} \oint_{\gamma_l} \frac{f(\zeta)}{\zeta - z} d\zeta \right) dz \\ &= - \sum_{l=1}^k \oint_{\gamma_l} \left( \sum_{j=1}^n k_j \frac{1}{2\pi i} \oint_{\sigma_j} \frac{1}{z - \zeta} dz \right) f(\zeta) d\zeta \\ &= - \sum_{l=1}^k \oint_{\gamma_l} \left( \sum_{j=1}^n k_j n(\sigma_j; \zeta) \right) f(\zeta) d\zeta \\ &= - \sum_{l=1}^k \oint_{\gamma_l} n(\Sigma; \zeta) f(\zeta) d\zeta. \end{aligned} \tag{6.12}$$

Now we consider the index  $n(\Sigma; \zeta) = \sum_{j=1}^n k_j n(\sigma_j; \zeta)$  when, as in the integrals in (6.12),  $\zeta$  belongs to any of  $\gamma_1^*, \dots, \gamma_k^*$ . Since every such  $\zeta$  is in  $\Omega \setminus K$ , either  $\zeta \notin \overline{D_0}(R)$  or the distance of  $\zeta$  from  $\Omega^c$  is  $< 2\delta$ . If  $\zeta \notin \overline{D_0}(R)$ , then, since  $\Sigma$  is in  $\overline{D_0}(R)$ , we have that  $n(\Sigma; \zeta) = 0$ . If the distance of  $\zeta$  from  $\Omega^c$  is  $< 2\delta$ , then there is  $w \in \Omega^c$  so that  $|\zeta - w| < 2\delta$ . Then every point of the linear segment  $[\zeta, w]$  has distance  $< 2\delta$  from  $w$  and hence from  $\Omega^c$ . Thus  $[\zeta, w]$  is not contained in  $K$  which implies that  $[\zeta, w]$  is in the complement of  $\sigma_1^*, \dots, \sigma_n^*$ . Since  $[\zeta, w]$  is connected and it is contained in the complement of every  $\sigma_j^*$  we have that  $n(\sigma_j; \zeta) = n(\sigma_j; w)$  for every  $j = 1, \dots, n$ . Therefore,

$$n(\Sigma; \zeta) = \sum_{j=1}^n k_j n(\sigma_j; \zeta) = \sum_{j=1}^n k_j n(\sigma_j; w) = n(\Sigma; w) = 0$$

because  $w \in \Omega^c$  and  $\Sigma$  is null-homologous in  $\Omega$ . Now (6.12) implies  $\oint_{\Sigma} f(z) dz = 0$ .  $\square$

It is interesting to see that the assumption of our last result is at the same time a special case of it. Indeed, if we take any  $w \in \Omega^c$ , then the function  $f(z) = \frac{1}{z-w}$  is holomorphic in  $\Omega$  and the theorem of Cauchy implies that  $\oint_{\Sigma} \frac{1}{z-w} dz = 0$ . But this says that  $n(\Sigma; w) = 0$ . In other words, we have the following situation. The assumption that  $\Sigma$  is null-homologous in  $\Omega$  is equivalent to the validity of the theorem of Cauchy for the very particular holomorphic functions of the form  $f(z) = \frac{1}{z-w}$  for every  $w \in \Omega^c$ . Therefore the real content of the theorem of Cauchy is that *the validity of  $\oint_{\Sigma} f(z) dz = 0$  for the special holomorphic functions in  $\Omega$  of the form  $f(z) = \frac{1}{z-w}$  for every  $w \in \Omega^c$  implies its validity for every function  $f$  which is holomorphic in  $\Omega$ .*

**Example 6.4.1.** Let  $\gamma$  be any piecewise smooth closed curve in the convex region  $\Omega$  and let  $w \in \Omega^c$ . Then  $w$  is contained in the unbounded connected component of  $\mathbb{C} \setminus \gamma^*$  and proposition 6.10 implies that  $n(\gamma; w) = 0$ . Hence  $\gamma$  is null-homologous in  $\Omega$ . Now the theorem of Cauchy for general open sets says that  $\oint_{\gamma} f(z) dz = 0$  for every  $f$  holomorphic in  $\Omega$ . We conclude that *the theorem of Cauchy for convex regions is a corollary of the theorem of Cauchy for general open sets.*

**Example 6.4.2.** We consider the open set  $D_{z_0}(R_1, R_2)$  with  $0 \leq R_1 < R_2 \leq +\infty$ . We consider the closed curve  $\gamma$  which describes the circle  $C_{z_0}(r)$ , with  $R_1 < r < R_2$ , once and in the positive direction. This curve is *not* null-homologous in  $D_{z_0}(R_1, R_2)$ . Indeed,  $z_0$  is in the complement of  $D_{z_0}(R_1, R_2)$  and  $n(\gamma; z_0) = \frac{1}{2\pi i} \oint_{C_{z_0}(r)} \frac{1}{z-z_0} dz = 1$ . Therefore, we do not expect that  $\oint_{\gamma} f(z) dz = 0$  is true for *every*  $f$  which is holomorphic in  $D_{z_0}(R_1, R_2)$ . In fact, this is certainly not true for  $f(z) = \frac{1}{z-z_0}$  which is holomorphic in  $D_{z_0}(R_1, R_2)$ .

**Example 6.4.3.** We consider the same open set  $D_{z_0}(R_1, R_2)$  as in the previous example and an arbitrary piecewise smooth closed curve  $\gamma$  in  $D_{z_0}(R_1, R_2)$ . We shall see how we can evaluate  $\oint_{\gamma} f(z) dz$  with a minimum of effort for any  $f$  holomorphic in  $D_{z_0}(R_1, R_2)$ . It is clear that, depending on the specific curve  $\gamma$ , it may be difficult to evaluate the integral using a parametric equation of  $\gamma$ .

Let us assume that the shape of the trajectory and the direction of  $\gamma$  allow us to count the number of rotations of  $\gamma$  around  $z_0$ , i.e. we assume that we know the integer  $k = n(\gamma; z_0)$ . Since the disc  $\overline{D_{z_0}}(R_1)$  is one of the two connected components of the complement of  $D_{z_0}(R_1, R_2)$ , we have that  $n(\gamma; z) = k$  for every  $z \in \overline{D_{z_0}}(R_1)$ . On the other hand, we have that  $n(\gamma; z) = 0$  for every  $z$  in the unbounded connected component of the complement of  $D_{z_0}(R_1, R_2)$ , which is  $\overline{D_{z_0}}(R_2, +\infty)$ . Now we take a piecewise smooth closed curve  $\gamma_1$  in  $D_{z_0}(R_1, R_2)$  such that the  $\oint_{\gamma_1} f(z) dz$  may be much easier to evaluate than the original  $\oint_{\gamma} f(z) dz$ . For instance, we may consider  $\gamma_1$  to describe the circle  $C_{z_0}(r)$  with  $R_1 < r < R_2$  once and in the positive direction. In this case we have that  $n(\gamma_1; z) = 1$  for every  $z \in \overline{D_{z_0}}(R_1)$  and  $n(\gamma_1; z) = 0$  for every  $z \in \overline{D_{z_0}}(R_2, +\infty)$ . Now we form the cycle  $\Sigma = 1\gamma + (-k)\gamma_1$  and we have  $n(\Sigma; z) = 1n(\gamma; z) + (-k)n(\gamma_1; z) = k + (-k) = 0$  for every  $z \in \overline{D_{z_0}}(R_1)$  and also  $n(\Sigma; z) = 1n(\gamma; z) + (-k)n(\gamma_1; z) = 0 + 0 = 0$  for every  $z \in \overline{D_{z_0}}(R_2, +\infty)$ . Therefore,  $\Sigma$  is null-homologous in  $D_{z_0}(R_1, R_2)$  and the theorem of Cauchy implies  $0 = \oint_{\Sigma} f(z) dz = 1 \oint_{\gamma} f(z) dz + (-k) \oint_{\gamma_1} f(z) dz$  and hence

$$\oint_{\gamma} f(z) dz = k \oint_{\gamma_1} f(z) dz = k \oint_{C_{z_0}(r)} f(z) dz.$$

We see that the evaluation of  $\oint_{\gamma} f(z) dz$  has been reduced to the evaluation of the possibly much simpler integral  $\oint_{C_{z_0}(r)} f(z) dz$  and the evaluation of the index  $n(\gamma; z_0)$ . We shall generalize this technique in the following sections and chapters.

Now we generalize Cauchy's formulas for derivatives.

**Cauchy's formula for derivatives and closed curves in general open sets.** *If  $f$  is holomorphic in the open set  $\Omega$  and if the cycle  $\Sigma$ , consisting of piecewise smooth closed curves, is null-homologous in  $\Omega$ , then for all  $n \in \mathbb{N}_0$  we have*

$$n(\Sigma; z)f^{(n)}(z) = \frac{n!}{2\pi i} \oint_{\Sigma} \frac{f(\zeta)}{(\zeta-z)^{n+1}} d\zeta$$

for every  $z \in \Omega$  which does not belong to the trajectory of any closed curve forming  $\Sigma$ .

*Proof.* The function  $F(\zeta) = \frac{f(\zeta)-f(z)}{\zeta-z}$  is holomorphic in  $\Omega \setminus \{z\}$ . Since  $z$  is a root of  $f(\zeta) - f(z)$ , the singularity  $z$  is removable. Therefore, we may define  $F$  at  $z$  with  $F(z) = \lim_{\zeta \rightarrow z} \frac{f(\zeta)-f(z)}{\zeta-z} = f'(z)$  and then  $F$  becomes holomorphic in  $\Omega$ . Now we apply the theorem of Cauchy in general open sets and get  $\oint_{\Sigma} \frac{f(\zeta)-f(z)}{\zeta-z} d\zeta = \oint_{\Sigma} F(\zeta) d\zeta = 0$ , which implies

$$\oint_{\Sigma} \frac{f(\zeta)}{\zeta-z} d\zeta = f(z) \oint_{\Sigma} \frac{1}{\zeta-z} d\zeta = f(z)2\pi i n(\Sigma; z)$$

for every  $z \in \Omega$  which does not belong to the trajectory of any closed curve forming  $\Sigma$ . This is the result of the statement in the case  $n = 0$ . For derivatives of order  $n \geq 1$  we differentiate both sides of the last formula, just as in the proof of the same theorem in convex sets, using the fact that the index of  $\Sigma$  is constant in a neighborhood of  $z$ .  $\square$

### Exercises.

**6.4.1.** Let  $f$  be holomorphic in  $\mathbb{D} \setminus \{0\}$ . If the piecewise smooth closed curve  $\gamma$  is in  $\mathbb{D} \setminus \{0\}$  and  $n(\gamma; 0) = 0$ , evaluate  $\oint_{\gamma} f(z) dz$ .

**6.4.2.** Let  $f$  be holomorphic in  $\mathbb{C}$  and  $f(1) = 6$ ,  $f(-1) = 10$ . Prove that, if  $\gamma$  is any piecewise smooth closed curve in  $\mathbb{C} \setminus \{-1, 1\}$ , then  $\frac{1}{2\pi i} \oint_{\gamma} \frac{f(z)}{z^2-1} dz$  can take every integral value.

**6.4.3.** Let  $f(z) = (\frac{1}{z} + \frac{a}{z^3})e^z$  for  $z \neq 0$ . Find all the values of  $a$  so that  $\oint_{\gamma} f(z) dz = 0$  for every piecewise smooth closed curve  $\gamma$  in  $\mathbb{C} \setminus \{0\}$ .

**6.4.4.** (i) Find all possible values of  $\oint_{\gamma} \frac{2z-1}{z^2-z} dz$ , where  $\gamma$  is an arbitrary piecewise smooth closed curve in  $\mathbb{C} \setminus \{0, 1\}$ .

(ii) Find all possible values of  $\int_{\gamma} \frac{2z-1}{z^2-z} dz$ , where  $\gamma$  is an arbitrary piecewise smooth curve in  $\mathbb{C} \setminus \{0, 1\}$  with initial endpoint  $-i$  and final endpoint  $i$ .

**6.4.5.** Find all possible values of  $\oint_{\gamma} \frac{\cos z}{z^2-\pi z} dz$ , where  $\gamma$  is an arbitrary piecewise smooth closed curve in  $\mathbb{C} \setminus \{0, \pi\}$ .

**6.4.6.** Let  $f$  be holomorphic in the open set  $\Omega$  and  $\gamma$  be a piecewise smooth closed curve null-homologous in  $\Omega$ . If  $|f(\zeta)| \leq 1$  for every  $\zeta \in \gamma^*$ ,  $z_0 \in \Omega$  and  $n(\gamma; z_0) \neq 0$ , prove that  $|f(z_0)| \leq 1$ .

## 6.5 The residue theorem.

Let  $z_0$  be an isolated singularity of  $f$  and let  $\sum_{-\infty}^{+\infty} a_n(z-z_0)^n$  be the Laurent series of  $f$  in the ring  $D_{z_0}(R) \setminus \{z_0\}$ . Then the coefficient  $a_{-1}$  is called **residue** of  $f$  at  $z_0$  and we denote

$$\text{Res}(f; z_0) = a_{-1}.$$

We know that  $\text{Res}(f; z_0) = a_{-1} = \frac{1}{2\pi i} \oint_{C_{z_0}(r)} f(\zeta) d\zeta$  for  $0 < r < R$ .

**Example 6.5.1.** If  $z_0$  is a removable singularity of  $f$ , then  $a_n = 0$  for every  $n < 0$  and in particular  $\text{Res}(f; z_0) = 0$ .

**Example 6.5.2.** Every function of the form  $f(z) = \frac{1}{(z-z_0)^N}$  with  $N \geq 2$  has residue 0 at  $z_0$ .

**Example 6.5.3.** If  $z_0$  is a pole of  $f$  of order  $N \geq 1$ , then we can find “easily” the residue of  $f$  at  $z_0$ . Indeed, there is a function  $g$  holomorphic in a disc  $D_{z_0}(R)$  so that  $g(z_0) \neq 0$  and  $f(z) = \frac{g(z)}{(z-z_0)^N}$  for every  $z \in D_{z_0}(R) \setminus \{z_0\}$ . From the Taylor series  $\sum_{n=0}^{+\infty} b_n(z-z_0)^n$  of  $g$  we see that  $\text{Res}(f; z_0) = b_{N-1} = \frac{g^{(N-1)}(z_0)}{(N-1)!}$ . For instance, if  $N = 1$ , then  $\text{Res}(f; z_0) = g(z_0)$  and, if  $N = 2$ , then  $\text{Res}(f; z_0) = g'(z_0)$ .

**Example 6.5.4.** We consider a power series of the form  $\sum_{n=-\infty}^{n=-1} a_n(z-z_0)^n$  and we assume that its radius of convergence is 0, i.e. that it converges in the ring  $D_{z_0}(0, +\infty)$ . If  $f$  is the holomorphic function defined by the power series in  $D_{z_0}(0, +\infty)$ , then

$$\frac{1}{2\pi i} \oint_{\gamma} f(\zeta) d\zeta = n(\gamma; z_0) a_{-1} = n(\gamma; z_0) \text{Res}(f; z_0)$$

for every piecewise smooth closed curve  $\gamma$  in  $\mathbb{C} \setminus \{z_0\}$ . Indeed, since the power series converges uniformly in the compact set  $\gamma^*$  which is contained in its ring of convergence, we have

$$\frac{1}{2\pi i} \oint_{\gamma} f(\zeta) d\zeta = \sum_{n=-\infty}^{n=-1} \frac{a_n}{2\pi i} \oint_{\gamma} (\zeta - z_0)^n d\zeta = \frac{a_{-1}}{2\pi i} \oint_{\gamma} \frac{1}{\zeta - z_0} d\zeta = n(\gamma; z_0) \text{Res}(f; z_0),$$

where, for  $n \leq -2$  we used the result of example 4.5.3. Of course, this result holds for a general cycle  $\Sigma$  which consists of piecewise smooth closed curves  $\gamma$  in  $\mathbb{C} \setminus \{z_0\}$ .

The residue theorem is a generalization of the last example.

**The residue theorem.** *Let  $f$  be holomorphic, except for isolated singularities, in the open set  $\Omega$  and  $\Sigma$  be a cycle which is null-homologous in  $\Omega$  and so that no isolated singularity of  $f$  is in the trajectory of any of the closed curves forming  $\Sigma$ . Then  $n(\Sigma; z) \neq 0$  for at most finitely many isolated singularities  $z$  of  $f$ . Moreover, if  $\Sigma$  consists of piecewise smooth closed curves, then*

$$\frac{1}{2\pi i} \oint_{\Sigma} f(\zeta) d\zeta = \sum_{z \text{ sing. of } f} n(\Sigma; z) \text{Res}(f; z),$$

where the sum, extended over all isolated singularities of  $f$  in  $\Omega$ , is finite.

*First proof.* Let us assume that  $n(\Sigma; z) \neq 0$  for infinitely many isolated singularities  $z$  of  $f$  in  $\Omega$ . Then there is a sequence  $(z_n)$  of distinct isolated singularities of  $f$  in  $\Omega$  so that  $n(\Sigma; z_n) \neq 0$  for every  $n$ . Since the trajectories of the curves which form  $\Sigma$  are bounded sets,  $\Sigma$  is in some disc  $D_0(R)$ . Hence  $n(\Sigma; z) = 0$  for every  $z$  outside  $D_0(R)$ . Thus, the sequence  $(z_n)$  is in  $D_0(R)$ . The Bolzano-Weierstrass theorem implies that there is a subsequence  $(z_{n_k})$  so that  $z_{n_k} \rightarrow z$  for some  $z$ . Then  $z$  is a limit point of  $\Omega$ .

If  $z \in \Omega$ , then either  $f$  is holomorphic at  $z$  or  $z$  is an isolated singularity of  $f$ . In any case, there are no isolated singularities of  $f$  in a neighborhood of  $z$ , except perhaps  $z$  itself. This contradicts  $z_{n_k}$  being distinct and  $z_{n_k} \rightarrow z$ .

If  $z \in \partial\Omega$  and hence  $z \in \Omega^c$ , then  $n(\Sigma; z) = 0$ . Now there is a disc  $D_z(r)$  which does not intersect any of the trajectories of the curves which form  $\Sigma$ . Since  $D_z(r)$  is connected, we have that  $n(\Sigma; w) = 0$  for every  $w \in D_z(r)$ . But this contradicts  $z_{n_k} \rightarrow z$  and  $n(\Sigma; z_{n_k}) \neq 0$  for all  $k$ . In any case we arrive at a contradiction and thus  $n(\Sigma; z) \neq 0$  for at most finitely many isolated singularities  $z$  of  $f$  in  $\Omega$ . Therefore, the sum  $\sum_{z \text{ sing. of } f} n(\Sigma; z) \text{Res}(f; z)$  is finite.

Let  $z_1, \dots, z_n$  be the isolated singularities of  $f$  in  $\Omega$  with  $n(\Sigma; z_k) \neq 0$  for  $k = 1, \dots, n$ . I.e.  $n(\Sigma; z) = 0$  for every other isolated singularity  $z$  of  $f$  in  $\Omega$ .

We define the integers

$$p_1 = n(\Sigma; z_1), \dots, p_n = n(\Sigma; z_n)$$



and then  $\sum_z \text{sing. of } f \ n(\Sigma; z) \text{ Res}(f; z) = \sum_{k=1}^n p_k \text{ Res}(f; z_k)$ . Therefore, it is enough to prove

$$\frac{1}{2\pi i} \oint_{\Sigma} f(\zeta) d\zeta = \sum_{k=1}^n p_k \text{ Res}(f; z_k). \quad (6.13)$$

Since every  $z_1, \dots, z_n$  is an isolated singularity, there are disjoint closed discs  $\overline{D}_{z_k}(r_k)$  for  $k = 1, \dots, n$  so that each of them contains no singularity of  $f$  except its center. We denote  $\gamma_k$  the closed curve which describes the circle  $C_{z_k}(r_k)$  once and in the positive direction. We consider the cycle

$$\Sigma' = \Sigma + (-p_1) \gamma_1 + \dots + (-p_n) \gamma_n$$

and the open set

$$\Omega' = \Omega \setminus \{z \in \Omega \mid z \text{ singularity of } f\}.$$

Clearly,  $f$  is holomorphic in  $\Omega'$  and we shall prove that the cycle  $\Sigma'$  is null-homologous in  $\Omega'$ , i.e.  $n(\Sigma'; z) = 0$  for every  $z \notin \Omega'$ . If  $z \notin \Omega'$ , then either  $z \notin \Omega$  or  $z = z_1, \dots, z_n$  or  $z$  is any other isolated singularity of  $f$  in  $\Omega$ .

If  $z \notin \Omega$  or if  $z$  is any isolated singularity of  $f$  in  $\Omega$  different from  $z_1, \dots, z_n$ , then  $n(\Sigma; z) = 0$  and  $n(\gamma_k; z) = 0$  for every  $k$ . Therefore

$$n(\Sigma'; z) = n(\Sigma; z) - p_1 n(\gamma_1; z) - \dots - p_n n(\gamma_n; z) = 0.$$

If  $z = z_{k_0}$  for some  $k_0$ , then  $n(\Sigma; z) = n(\Sigma; z_{k_0}) = p_{k_0}$  and  $n(\gamma_{k_0}; z) = n(\gamma_{k_0}; z_{k_0}) = 1$  and  $n(\gamma_k; z) = n(\gamma_k; z_{k_0}) = 0$  for every  $k \neq k_0$ . Therefore

$$n(\Sigma'; z) = n(\Sigma; z) - p_1 n(\gamma_1; z) - \dots - p_n n(\gamma_n; z) = p_{k_0} - p_{k_0} = 0.$$

Thus,  $\Sigma'$  is null-homologous in  $\Omega'$ . Since  $f$  is holomorphic in  $\Omega'$ , the theorem of Cauchy implies  $\oint_{\Sigma'} f(\zeta) d\zeta = 0$ . Hence

$$\oint_{\Sigma} f(\zeta) d\zeta = \sum_{k=1}^n p_k \oint_{\gamma_k} f(\zeta) d\zeta = 2\pi i \sum_{k=1}^n p_k \text{ Res}(f; z_k)$$

and we proved (6.13).

*Second proof.* We follow the first proof up to the point where we considered the isolated singularities  $z_1, \dots, z_n$  of  $f$ . I.e.  $n(\Sigma; z) = 0$  for every isolated singularity of  $f$  different from  $z_1, \dots, z_n$ . Now, we consider the corresponding singular parts  $s_1, \dots, s_n$  of  $f$  at  $z_1, \dots, z_n$ . Then we know from section 5.9 that  $f - s_k$  is holomorphic at  $z_k$  and also that  $s_k$  is holomorphic in  $\mathbb{C} \setminus \{z_k\}$ . Hence the function  $g = f - s_1 - \dots - s_n$  is holomorphic in  $\Omega$  except at the isolated singularities of  $f$  which are different from  $z_1, \dots, z_n$ . We consider the open set

$$\Omega'' = \Omega \setminus \{z \in \Omega \mid z \text{ is a singularity of } f, z \neq z_1, \dots, z_n\}.$$

and then  $g$  is holomorphic in  $\Omega''$ . Also,  $\Sigma$  is null-homologous in  $\Omega''$ . Therefore, the theorem of Cauchy implies that  $\frac{1}{2\pi i} \oint_{\Sigma} g(\zeta) d\zeta = 0$  and hence

$$\begin{aligned} \frac{1}{2\pi i} \oint_{\Sigma} f(\zeta) d\zeta &= \sum_{k=1}^n \frac{1}{2\pi i} \oint_{\Sigma} s_k(\zeta) d\zeta = \sum_{k=1}^n n(\Sigma; z_k) \text{ Res}(s_k; z_k) \\ &= \sum_{k=1}^n n(\Sigma; z_k) \text{ Res}(f; z_k), \end{aligned}$$

where for the second equality we used the result of example 6.5.4. □

The residue theorem is a powerful tool for the evaluation of integrals, because it reduces this evaluation to the location of the isolated singularities of the function to be integrated and to the evaluation of the corresponding residues. Let us see some characteristic examples.

**Example 6.5.5.** Evaluation of  $\int_{-\infty}^{+\infty} r(x) dx$ , where  $r = \frac{p}{q}$  is a rational function,  $\deg q \geq \deg p + 2$  and  $q$  has no real roots.

Let  $p(x) = a_n x^n + \cdots + a_1 x + a_0$ , with  $a_n \neq 0$ , and  $q(x) = b_m x^m + \cdots + b_1 x + b_0$ , with  $b_m \neq 0$ , and  $m \geq n + 2$ . Then  $r$  is continuous in  $\mathbb{R}$  and the generalized integral  $\int_{-\infty}^{+\infty} r(x) dx$  converges.

To see this, we observe that  $\lim_{z \rightarrow \infty} z^{m-n} r(z) = \frac{a_n}{b_m}$ . Hence, if  $c = \frac{|a_n|}{|b_m|} > 0$ , there is  $R_0 > 0$  so that

$$\frac{c}{2} \leq |z|^{m-n} |r(z)| \leq 2c \quad \text{when } |z| \geq R_0. \quad (6.14)$$

Now, since  $m - n \geq 2$ , we get  $\int_{-\infty}^{-R_0} |r(x)| dx \leq 2c \int_{-\infty}^{-R_0} \frac{1}{|x|^{m-n}} dx < +\infty$  and  $\int_{R_0}^{+\infty} |r(x)| dx \leq 2c \int_{R_0}^{+\infty} \frac{1}{x^{m-n}} dx < +\infty$ . Thus, the integrals  $\int_{-\infty}^{-R_0} r(x) dx$ ,  $\int_{R_0}^{+\infty} r(x) dx$  converge absolutely and so they converge. Moreover,  $r$  is continuous in  $[-R_0, R_0]$  and so the integral  $\int_{-\infty}^{+\infty} r(x) dx$  converges.

We consider the roots of  $q$  in the upper halfplane and let them be  $z_1, \dots, z_M$ , where  $M \leq m$ . We take any  $R > R_0$  so that  $z_1, \dots, z_M$  are contained in the disc  $D_0(R)$ . We apply the residue theorem with  $r = \frac{p}{q}$  which is holomorphic in  $\mathbb{C}$  except for the roots of  $q$  and with the closed curve  $\gamma_R$  which is the sum of the linear segment  $[-R, R]$ , with parametric equation  $z = x$ ,  $x \in [-R, R]$ , and of the curve  $\sigma_R$ , with parametric equation  $z = Re^{it}$ ,  $t \in [0, \pi]$ , which describes the upper semicircle of  $C_0(R)$  from  $R$  to  $-R$ . The trajectory of  $\gamma_R$  contains no isolated singularity of  $r$ . Since  $\gamma_R$  rotates around each of  $z_1, \dots, z_M$  once and in the positive direction, the residue theorem implies

$$\frac{1}{2\pi i} \oint_{\gamma_R} r(z) dz = \text{Res}(r; z_1) + \cdots + \text{Res}(r; z_M).$$

We have that  $\oint_{\gamma_R} r(z) dz = \int_{[-R, R]} r(z) dz + \int_{\sigma_R} r(z) dz$  and hence

$$\int_{-R}^R r(x) dx = \int_{[-R, R]} r(z) dz = 2\pi i (\text{Res}(r; z_1) + \cdots + \text{Res}(r; z_M)) - \int_{\sigma_R} r(z) dz.$$

Since  $R > R_0$ , (6.14) and  $m \geq n + 2$  imply  $|\int_{\sigma_R} r(z) dz| \leq \frac{2c}{R^{m-n}} \pi R \rightarrow 0$  when  $R \rightarrow +\infty$  and we conclude that

$$\int_{-\infty}^{+\infty} r(x) dx = 2\pi i (\text{Res}(r; z_1) + \cdots + \text{Res}(r; z_M)).$$

Thus, to evaluate  $\int_{-\infty}^{+\infty} r(x) dx$  we need only to find the residues of  $r$  at the poles  $z_1, \dots, z_M$  of  $r$  in the upper halfplane.

**Example 6.5.6.** Evaluation of  $\text{pv} \int_{-\infty}^{+\infty} r(x) dx$ , where  $r = \frac{p}{q}$  is a rational function,  $\deg q = \deg p + 1$  and  $q$  has no real root.

Let  $p(x) = a_n x^n + \cdots + a_1 x + a_0$ , with  $a_n \neq 0$ , and  $q(x) = b_{n+1} x^{n+1} + \cdots + b_1 x + b_0$ , with  $b_{n+1} \neq 0$ . It is easy to see that the generalized integral  $\int_{-\infty}^{+\infty} r(x) dx$  does not converge. Indeed, we recall the estimate (6.14), i.e.  $|r(z)| \geq \frac{c}{2|z|}$  when  $|z| \geq R_0$ . Therefore, for real  $z = x$  we have that  $|r(x)| \geq \frac{c}{2x}$  when  $x \geq R_0$ . Now,  $r$  has constant sign in  $[R_0, +\infty)$  and hence  $|\int_{R_0}^{+\infty} r(x) dx| = \int_{R_0}^{+\infty} |r(x)| dx \geq \frac{c}{2} \int_{R_0}^{+\infty} \frac{1}{x} dx = +\infty$ . Thus,  $\int_{R_0}^{+\infty} r(x) dx = +\infty$  or  $-\infty$  and, similarly,  $\int_{-\infty}^{-R_0} r(x) dx = +\infty$  or  $-\infty$ .

Since the generalized integral diverges, we examine its **principal value**, i.e.

$$\text{pv} \int_{-\infty}^{+\infty} r(x) dx = \lim_{R \rightarrow +\infty} \int_{-R}^R r(x) dx.$$

It is easy to see that  $r(z) - \frac{a_n}{b_{n+1}} \frac{1}{z}$  is a rational function whose denominator has degree two units larger than the degree of its numerator. According to the previous example, there is  $R_0 > 0$  so that

$$\left| r(z) - \frac{a_n}{b_{n+1}} \frac{1}{z} \right| \leq \frac{C}{|z|^2} \quad \text{when } |z| \geq R_0. \quad (6.15)$$

As in the previous example, we consider the roots  $z_1, \dots, z_M$  of  $q$  in the upper halfplane and we take  $R > R_0$  so that  $z_1, \dots, z_M$  are contained in  $D_0(R)$ . We apply the residue theorem with  $r = \frac{p}{q}$  and the same closed curve  $\gamma_R$  and we get

$$\frac{1}{2\pi i} \oint_{\gamma_R} r(z) dz = \text{Res}(r; z_1) + \cdots + \text{Res}(r; z_M).$$

Now,  $\oint_{\gamma_R} r(z) dz = \int_{[-R,R]} r(z) dz + \int_{\sigma_R} r(z) dz$  and hence

$$\int_{-R}^R r(x) dx = 2\pi i(\text{Res}(r; z_1) + \cdots + \text{Res}(r; z_M)) - \int_{\sigma_R} \left(r(z) - \frac{a_n}{b_{n+1}} \frac{1}{z}\right) dz - \frac{a_n}{b_{n+1}} \int_{\sigma_R} \frac{1}{z} dz.$$

The last term is

$$\frac{a_n}{b_{n+1}} \int_{\sigma_R} \frac{1}{z} dz = \frac{a_n}{b_{n+1}} \int_0^\pi \frac{1}{Re^{it}} iRe^{it} dt = i\pi \frac{a_n}{b_{n+1}}.$$

Since  $R > R_0$ , we have from (6.15) that  $\left| \int_{\sigma_R} \left(r(z) - \frac{a_n}{b_{n+1}} \frac{1}{z}\right) dz \right| \leq \frac{C}{R^2} \pi R \rightarrow 0$  when  $R \rightarrow +\infty$  and we finally get

$$\text{pv} \int_{-\infty}^{+\infty} r(x) dx = 2\pi i(\text{Res}(r; z_1) + \cdots + \text{Res}(r; z_M)) - i\pi \frac{a_n}{b_{n+1}}.$$

**Example 6.5.7.** Evaluation of  $\text{pv} \int_{-\infty}^{+\infty} r(x) dx$ , where  $r = \frac{p}{q}$  is a rational function,  $\deg q \geq \deg p + 1$  and  $q$  has real roots, all with multiplicity 1.

Let  $p(x) = a_n x^n + \cdots + a_1 x + a_0$ , with  $a_n \neq 0$ , and  $q(x) = b_m x^m + \cdots + b_1 x + b_0$ , with  $b_m \neq 0$ , and  $m \geq n + 1$ . We assume that the real roots of  $q$  are  $x_1, \dots, x_n$  with  $x_1 < \dots < x_n$  and that these are not roots of  $p$ . We take  $\epsilon_0 > 0$  so that the intervals  $[x_1 - \epsilon_0, x_1 + \epsilon_0], \dots, [x_n - \epsilon_0, x_n + \epsilon_0]$  around the real roots of  $q$  are disjoint. In order for  $\int_{-\infty}^{+\infty} r(x) dx$  to converge, the generalized integrals  $\int_{x_k - \epsilon_0}^{x_k} r(x) dx$  and  $\int_{x_k}^{x_k + \epsilon_0} r(x) dx$  must converge for every  $x_k$ . This is not correct. Indeed, we write  $r(z) = \frac{p(z)}{(z-x_k)q_k(z)} = \frac{g_k(z)}{z-x_k}$ , where  $q_k$  is a polynomial with  $q_k(x_k) \neq 0$  and where  $g_k = \frac{p}{q_k}$  is a rational function holomorphic at  $x_k$ . Since  $\lim_{z \rightarrow x_k} g_k(z) = g_k(x_k) \neq 0$ , there is  $\epsilon_k$  with  $0 < \epsilon_k \leq \epsilon_0$  so that  $|g_k(z)| \geq \frac{1}{2} |g_k(x_k)|$  for every  $z$  with  $|z - x_k| \leq \epsilon_k$ . Hence,  $|r(z)| \geq \frac{1}{2} \frac{|g_k(x_k)|}{|z - x_k|}$  for every  $z$  with  $0 < |z - x_k| \leq \epsilon_k$ . The function  $r$  has constant sign in  $(x_k, x_k + \epsilon_k]$ . Therefore,  $\left| \int_{x_k}^{x_k + \epsilon_k} r(x) dx \right| = \int_{x_k}^{x_k + \epsilon_k} |r(x)| dx \geq \frac{|g_k(x_k)|}{2} \int_{x_k}^{x_k + \epsilon_k} \frac{1}{x - x_k} dx = +\infty$  and the generalized integral  $\int_{x_k}^{x_k + \epsilon_k} r(x) dx$  does not converge. Similarly,  $\int_{x_k - \epsilon_k}^{x_k} r(x) dx$  does not converge either. This is why we examine the **principal value** of  $\int_{-\infty}^{+\infty} r(x) dx$ , i.e.

$$\begin{aligned} \text{pv} \int_{-\infty}^{+\infty} r(x) dx &= \lim_{R \rightarrow +\infty, \epsilon \rightarrow 0+} \left( \int_{-R}^{x_1 - \epsilon} r(x) dx + \int_{x_1 + \epsilon}^{x_2 - \epsilon} r(x) dx + \cdots \right. \\ &\quad \left. \cdots + \int_{x_{n-1} - \epsilon}^{x_n - \epsilon} r(x) dx + \int_{x_n + \epsilon}^R r(x) dx \right) = \lim_{R \rightarrow +\infty, \epsilon \rightarrow 0+} I(R, \epsilon). \end{aligned} \quad (6.16)$$

We evaluate  $I(R, \epsilon)$  using a variant of the curve  $\gamma_R$  of the previous examples: the curve  $\gamma_{R, \epsilon}$ , which is the sum of the linear segments  $[-R, x_1 - \epsilon], [x_1 + \epsilon, x_2 - \epsilon], \dots, [x_{n-1} + \epsilon, x_n - \epsilon], [x_n + \epsilon, R]$ , of the curve  $\sigma_R$ , which describes the upper semicircle of  $C_0(R)$  from  $R$  to  $-R$ , and of the curves  $\sigma_{1, \epsilon}, \dots, \sigma_{n, \epsilon}$ , where each  $\sigma_{k, \epsilon}$  describes the upper semicircle of the corresponding  $C_{x_k}(\epsilon)$  from  $x_k - \epsilon$  to  $x_k + \epsilon$ . We just take  $R$  large enough and  $\epsilon$  small enough so that the curve  $\gamma_{R, \epsilon}$  rotates once and in the positive direction around each of the roots  $z_1, \dots, z_M$  of  $q$  in the upper halfplane. Then  $\gamma_{R, \epsilon}$  rotates no times around each of the remaining roots of  $q$ . The residue theorem implies that

$$\oint_{\gamma_{R, \epsilon}} r(z) dz = 2\pi i(\text{Res}(r; z_1) + \cdots + \text{Res}(r; z_M))$$

and hence

$$\begin{aligned} I(R, \epsilon) &= 2\pi i(\text{Res}(r; z_1) + \cdots + \text{Res}(r; z_M)) - \int_{\sigma_R} r(z) dz \\ &\quad - \int_{\sigma_{1, \epsilon}} r(z) dz - \cdots - \int_{\sigma_{n, \epsilon}} r(z) dz. \end{aligned} \quad (6.17)$$

Now,  $x_k$  is a pole of  $r$  of order 1 and  $r$  can be written  $r(z) = \frac{c_k}{z - x_k} + f_k(z)$  for  $z \neq x_k$  in a disc with center  $x_k$ , where  $f_k$  is holomorphic at  $x_k$  and  $c_k = \text{Res}(r; x_k)$ . Since  $f_k$  is bounded in a disc with center  $x_k$ , there is  $M_k \geq 0$  and  $\epsilon'_k > 0$  so that  $|f_k(z)| \leq M_k$  for  $|z - x_k| \leq \epsilon'_k$ . Thus,  $0 < \epsilon \leq \epsilon'_k$  implies  $\left| \int_{\sigma_{k, \epsilon}} f_k(z) dz \right| \leq M_k \pi \epsilon$  and hence  $\lim_{\epsilon \rightarrow 0+} \int_{\sigma_{k, \epsilon}} f_k(z) dz = 0$ . Therefore,

$$\begin{aligned} \int_{\sigma_{k, \epsilon}} r(z) dz &= c_k \int_{\sigma_{k, \epsilon}} \frac{1}{z - x_k} dz + \int_{\sigma_{k, \epsilon}} f_k(z) dz \\ &= -\pi i c_k + \int_{\sigma_{k, \epsilon}} f_k(z) dz \rightarrow -\pi i c_k \quad \text{when } \epsilon \rightarrow 0+. \end{aligned} \quad (6.18)$$

The limit of  $\int_{\sigma_R} r(z) dz$  when  $R \rightarrow +\infty$  has been evaluated in the previous two examples:

$$\lim_{R \rightarrow +\infty} \int_{\sigma_R} r(z) dz = \begin{cases} 0, & \text{if } m \geq n + 2 \\ i\pi \frac{a_n}{b_{n+1}}, & \text{if } m = n + 1 \end{cases} \quad (6.19)$$

Now, (6.16), (6.17), (6.18) and (6.19) imply

$$\begin{aligned} \text{pv} \int_{-\infty}^{+\infty} r(x) dx &= 2\pi i (\text{Res}(r; z_1) + \cdots + \text{Res}(r; z_M)) \\ &+ \pi i (\text{Res}(r; x_1) + \cdots + \text{Res}(r; x_n)) - \begin{cases} 0, & \text{if } m \geq n + 2 \\ i\pi \frac{a_n}{b_{n+1}}, & \text{if } m = n + 1 \end{cases} \end{aligned}$$

**Example 6.5.8.** Evaluation of  $\int_{-\infty}^{+\infty} r(x) \cos x dx$ ,  $\int_{-\infty}^{+\infty} r(x) \sin x dx$  (or of their principal values), where  $r = \frac{p}{q}$  is a rational function,  $\deg q \geq \deg p + 1$ , the real roots of  $q$  (if they exist) have multiplicity 1 and, also, the coefficients of  $p, q$  are real numbers.

Since the coefficients of  $p, q$  are real, we have that  $r(x) \in \mathbb{R}$  for every  $x \in \mathbb{R}$  which is not a root of  $q$ . Hence,

$$\int_{-\infty}^{+\infty} r(x) \cos x dx = \text{Re} \int_{-\infty}^{+\infty} r(x) e^{ix} dx, \quad \int_{-\infty}^{+\infty} r(x) \sin x dx = \text{Im} \int_{-\infty}^{+\infty} r(x) e^{ix} dx$$

and we evaluate  $\int_{-\infty}^{+\infty} r(x) e^{ix} dx$  (or its principal value).

The method of evaluation has been described already in the previous three examples. We use either the curve  $\gamma_R$  or the curve  $\gamma_{R,\epsilon}$  and we evaluate the residues of  $r(z)e^{iz}$  at the roots of  $q$ .

We shall concentrate on the important specific generalized integral

$$\int_0^{+\infty} \frac{\sin x}{x} dx = \frac{1}{2} \int_{-\infty}^{+\infty} \frac{\sin x}{x} dx.$$

(Equality holds because  $\frac{\sin x}{x}$  is even.) We shall evaluate  $\text{pv} \int_{-\infty}^{+\infty} \frac{e^{ix}}{x} dx$  instead of  $\int_{-\infty}^{+\infty} \frac{\sin x}{x} dx$ . Observe that  $\frac{e^{ix}}{x} = \frac{\cos x}{x} + i \frac{\sin x}{x}$  diverges at 0 because its real part  $\frac{\cos x}{x}$  diverges at 0. The imaginary part  $\frac{\sin x}{x}$  converges at 0 and, in fact, if we define  $\frac{\sin x}{x}$  at 0 to have value  $\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$ , then it becomes continuous at 0.

The function  $\frac{e^{iz}}{z}$  is holomorphic in  $\mathbb{C}$  except for a pole at 0 of order 1. We consider the closed curve  $\gamma_{R,\epsilon}$  which is the sum of the linear segments  $[-R, -\epsilon]$  and  $[\epsilon, R]$ , of the curve  $\sigma_R$ , which describes the upper semicircle of  $C_0(R)$  from  $R$  to  $-R$ , and of the curve  $\sigma_\epsilon$ , which describes the upper semicircle of  $C_0(\epsilon)$  from  $-\epsilon$  to  $\epsilon$ . Then  $\gamma_{R,\epsilon}$  does not rotate around the pole 0 of  $\frac{e^{iz}}{z}$ . The residue theorem implies  $\oint_{\gamma_{R,\epsilon}} \frac{e^{iz}}{z} dz = 0$  and hence

$$\int_{-R}^{-\epsilon} \frac{e^{ix}}{x} dx + \int_{\epsilon}^R \frac{e^{ix}}{x} dx = - \int_{\sigma_R} \frac{e^{iz}}{z} dz - \int_{\sigma_\epsilon} \frac{e^{iz}}{z} dz. \quad (6.20)$$

Now,

$$\int_{\sigma_R} \frac{e^{iz}}{z} dz = \int_0^\pi \frac{e^{iRe^{it}}}{Re^{it}} iRe^{it} dt = i \int_0^\pi e^{-R \sin t + iR \cos t} dt$$

and

$$\begin{aligned} \left| \int_{\sigma_R} \frac{e^{iz}}{z} dz \right| &\leq \int_0^\pi e^{-R \sin t} dt = 2 \int_0^{\pi/2} e^{-R \sin t} dt \leq 2 \int_0^{\pi/2} e^{-\frac{2R}{\pi} t} dt \\ &= \frac{\pi}{R} (1 - e^{-R}) \rightarrow 0 \quad \text{when } R \rightarrow +\infty. \end{aligned} \quad (6.21)$$

For the second inequality we used the well known inequality  $\sin t \geq \frac{2t}{\pi}$  for  $0 \leq t \leq \frac{\pi}{2}$ . From the Laurent series of  $\frac{e^{iz}}{z}$  at 0 we see that  $\frac{e^{iz}}{z} = \frac{1}{z} + h(z)$  for  $z \neq 0$ , where  $h$  is holomorphic in  $\mathbb{C}$ . Now,  $h$  is bounded in a neighborhood of 0, i.e. there is  $M \geq 0$  so that  $|h(z)| \leq 1$  when  $|z| \leq 1$ . Hence, for  $\epsilon \leq 1$  we have  $|\int_{\sigma_\epsilon} h(z) dz| \leq M\pi\epsilon \rightarrow 0$  when  $\epsilon \rightarrow 0+$ . Therefore

$$\begin{aligned} \int_{\sigma_\epsilon} \frac{e^{iz}}{z} dz &= \int_{\sigma_\epsilon} \frac{1}{z} dz + \int_{\sigma_\epsilon} h(z) dz \\ &= -\pi i + \int_{\sigma_\epsilon} h(z) dz \rightarrow -\pi i \quad \text{when } \epsilon \rightarrow 0+. \end{aligned} \quad (6.22)$$

From (6.20), (6.21) and (6.22):

$$\text{pv} \int_{-\infty}^{+\infty} \frac{e^{ix}}{x} dx = \lim_{\epsilon \rightarrow 0+, R \rightarrow +\infty} \left( \int_{-R}^{-\epsilon} \frac{e^{ix}}{x} dx + \int_{\epsilon}^R \frac{e^{ix}}{x} dx \right) = \pi i.$$

Since  $\frac{\cos x}{x}$  is odd and  $\frac{\sin x}{x}$  is even, we get  $\int_{-R}^{-\epsilon} \frac{e^{ix}}{x} dx + \int_{\epsilon}^R \frac{e^{ix}}{x} dx = 2i \int_{\epsilon}^R \frac{\sin x}{x} dx$  and hence

$$\int_0^{+\infty} \frac{\sin x}{x} dx = \lim_{\epsilon \rightarrow 0+, R \rightarrow +\infty} \int_{\epsilon}^R \frac{\sin x}{x} dx = \frac{\pi}{2}.$$

**Example 6.5.9.** We shall evaluate  $\int_0^{+\infty} \frac{\ln x}{x^2+4} dx$ .

We consider the holomorphic branch of the logarithm, which we shall denote  $\log z$ , in the open region  $A = \mathbb{C} \setminus \{iy \mid y \leq 0\}$  and which takes the value 0 at 1. This branch is given by

$$\log z = \ln r + i\theta \quad \text{for } z = re^{i\theta} \text{ with } r > 0 \text{ and } -\frac{\pi}{2} < \theta < \frac{3\pi}{2}.$$

The function  $\frac{\log z}{z^2+4}$  is holomorphic in  $A$  except for the point  $2i$  which is a pole of order 1. Indeed, we write  $\frac{\log z}{z^2+4} = \frac{(\log z)/(z+2i)}{z-2i} = \frac{g(z)}{z-2i}$  and we have that  $g(z) = \frac{\log z}{z+2i}$  is holomorphic in  $A$  with  $g(2i) = \frac{\pi}{8} - \frac{\ln 2}{4}i$ . Moreover,  $\text{Res}\left(\frac{\log z}{z^2+4}; 2i\right) = g(2i) = \frac{\pi}{8} - \frac{\ln 2}{4}i$ . Now we consider the closed curve  $\gamma_{R,\epsilon}$  of the previous example. We take  $R$  large enough and  $\epsilon$  small enough so that  $\gamma_{R,\epsilon}$  rotates once and in the positive direction around the pole  $2i$ . From the residue theorem we have that

$$\oint_{\gamma_{R,\epsilon}} \frac{\log z}{z^2+4} dz = 2\pi i \text{Res}\left(\frac{\log z}{z^2+4}; 2i\right) = \frac{\pi \ln 2}{2} + \frac{\pi^2}{4}i.$$

Taking real parts of both sides, we find

$$2 \int_{\epsilon}^R \frac{\ln x}{x^2+4} dx = \frac{\pi \ln 2}{2} - \text{Re} \int_{\sigma_R} \frac{\log z}{z^2+4} dz - \text{Re} \int_{\sigma_{\epsilon}} \frac{\log z}{z^2+4} dz.$$

Now,  $\left| \int_{\sigma_R} \frac{\log z}{z^2+4} dz \right| \leq \frac{\ln R + \pi}{R^2-4} \pi R \rightarrow 0$  when  $R \rightarrow +\infty$  and  $\left| \int_{\sigma_{\epsilon}} \frac{\log z}{z^2+4} dz \right| \leq \frac{\ln \epsilon + \pi}{4-\epsilon^2} \pi \epsilon \rightarrow 0$  when  $\epsilon \rightarrow 0+$ . Hence

$$\int_0^{+\infty} \frac{\ln x}{x^2+4} dx = \lim_{\epsilon \rightarrow 0+, R \rightarrow +\infty} \int_{\epsilon}^R \frac{\ln x}{x^2+4} dx = \frac{\pi \ln 2}{4}.$$

**Example 6.5.10.** We shall evaluate  $\int_0^{+\infty} \frac{x^{a-1}}{x+1} dx$  when  $0 < a < 1$ .

We write  $x^2$  instead of  $x$ :

$$\int_0^{+\infty} \frac{x^{a-1}}{x+1} dx = 2 \int_0^{+\infty} \frac{x^{2a-1}}{x^2+1} dx = 2 \int_0^{+\infty} \frac{x^b}{x^2+1} dx$$

with  $b = 2a - 1$  and  $-1 < b < 1$ .

We consider the holomorphic branch  $\log z$  of the previous example in the same region  $A$ . The function  $h(z) = e^{b \log z}$  is holomorphic in  $A$  and, if  $z = x$  is real, we have  $h(x) = e^{b \ln x} = x^b$ . The function  $\frac{h(z)}{z^2+1}$  is holomorphic in  $A$  except for a pole at  $i$  of order 1. Indeed, we write  $\frac{h(z)}{z^2+1} = \frac{h(z)/(z+i)}{z-i} = \frac{g(z)}{z-i}$  and we have that  $g(z) = \frac{h(z)}{z+i}$  is holomorphic in  $A$  with  $g(i) = \frac{h(i)}{2i} = \frac{e^{\frac{b\pi}{2}i}}{2i}$ . Moreover,  $\text{Res}\left(\frac{h(z)}{z^2+1}; i\right) = g(i) = \frac{e^{\frac{b\pi}{2}i}}{2i}$ . Now we consider the same closed curve  $\gamma_{R,\epsilon}$  of the previous example. The residue theorem implies

$$\oint_{\gamma_{R,\epsilon}} \frac{h(z)}{z^2+1} dz = 2\pi i \text{Res}\left(\frac{h(z)}{z^2+1}; i\right) = \pi e^{\frac{b\pi}{2}i},$$

and hence

$$(e^{b\pi i} + 1) \int_{\epsilon}^R \frac{x^b}{x^2+1} dx = \pi e^{\frac{b\pi}{2}i} - \int_{\sigma_R} \frac{h(z)}{z^2+1} dz - \int_{\sigma_{\epsilon}} \frac{h(z)}{z^2+1} dz.$$

Now  $\left| \int_{\sigma_R} \frac{h(z)}{z^2+1} dz \right| \leq \frac{R^b}{R^2-1} \pi R \rightarrow 0$  when  $R \rightarrow +\infty$  and  $\left| \int_{\sigma_{\epsilon}} \frac{h(z)}{z^2+1} dz \right| \leq \frac{\epsilon^b}{1-\epsilon^2} \pi \epsilon \rightarrow 0$  when  $\epsilon \rightarrow 0+$ . Hence

$$\int_0^{+\infty} \frac{x^{a-1}}{x+1} dx = 2 \int_0^{+\infty} \frac{x^b}{x^2+1} dx = \lim_{\epsilon \rightarrow 0+, R \rightarrow +\infty} 2 \int_{\epsilon}^R \frac{x^b}{x^2+1} dx = \frac{2\pi e^{\frac{b\pi}{2}i}}{e^{b\pi i} + 1} = \frac{\pi}{\sin a\pi}.$$

We shall evaluate  $\int_0^{+\infty} \frac{x^{a-1}}{x+1} dx$  in a different way.

We consider the holomorphic branch of the logarithm, which we shall denote  $\log z$  again, in the region  $B = \mathbb{C} \setminus \{x \mid x \geq 0\}$  and which takes the value  $i\pi$  at  $-1$ . This branch is given by

$$\log z = \ln r + i\theta \quad \text{for } z = re^{i\theta} \text{ with } r > 0 \text{ and } 0 < \theta < 2\pi.$$

The function  $h(z) = e^{(a-1)\log z}$  is holomorphic in  $B$ , and hence  $\frac{h(z)}{z+1}$  is holomorphic in  $B$  except at the point  $-1$  which is a pole of order 1. Indeed, we have  $\text{Res}\left(\frac{h(z)}{z+1}; -1\right) = h(-1) = e^{(a-1)\pi i}$ . We also consider the closed curve  $\gamma_{R,\epsilon,\delta}$  which is the sum of the curve  $\sigma_{R,\delta}$ , which describes the arc of  $C_0(R)$  from  $Re^{i\delta}$  to  $Re^{i(2\pi-\delta)}$  in the positive direction, of the curve  $\sigma_{\epsilon,\delta}$ , which describes the arc of  $C_0(\epsilon)$  from  $\epsilon e^{i(2\pi-\delta)}$  to  $\epsilon e^{i\delta}$  in the negative direction, of the linear segment  $[\epsilon e^{i\delta}, Re^{i\delta}]$  and of the linear segment  $[Re^{i(2\pi-\delta)}, \epsilon e^{i(2\pi-\delta)}]$ . The residue theorem implies that

$$\oint_{\gamma_{R,\epsilon,\delta}} \frac{h(z)}{z+1} dz = 2\pi i \text{Res}\left(\frac{h(z)}{z+1}; -1\right) = 2\pi i e^{(a-1)\pi i}$$

and hence

$$\begin{aligned} \int_{[\epsilon e^{i\delta}, Re^{i\delta}]} \frac{h(z)}{z+1} dz + \int_{[Re^{i(2\pi-\delta)}, \epsilon e^{i(2\pi-\delta)}]} \frac{h(z)}{z+1} dz \\ = 2\pi i e^{(a-1)\pi i} - \int_{\sigma_{R,\delta}} \frac{h(z)}{z+1} dz - \int_{\sigma_{\epsilon,\delta}} \frac{h(z)}{z+1} dz. \end{aligned}$$

Now,  $\left| \int_{\sigma_{R,\delta}} \frac{h(z)}{z+1} dz \right| \leq \frac{2\pi R^a}{R-1}$  and  $\left| \int_{\sigma_{\epsilon,\delta}} \frac{h(z)}{z+1} dz \right| \leq \frac{2\pi \epsilon^a}{1-\epsilon}$ . Therefore

$$\left| \int_{[\epsilon e^{i\delta}, Re^{i\delta}]} \frac{h(z)}{z+1} dz + \int_{[Re^{i(2\pi-\delta)}, \epsilon e^{i(2\pi-\delta)}]} \frac{h(z)}{z+1} dz - 2\pi i e^{(a-1)\pi i} \right| \leq \frac{2\pi R^a}{R-1} + \frac{2\pi \epsilon^a}{1-\epsilon}. \quad (6.23)$$

We have

$$\int_{[\epsilon e^{i\delta}, Re^{i\delta}]} \frac{h(z)}{z+1} dz = e^{ia\delta} \int_{\epsilon}^R \frac{r^{a-1}}{re^{i\delta}+1} dr.$$

Keeping  $\epsilon$  and  $R$  fixed, we take the limit when  $\delta \rightarrow 0+$ . Clearly,  $e^{ia\delta} \rightarrow 1$ . Also,  $\frac{1}{re^{i\delta}+1} \rightarrow \frac{1}{r+1}$  uniformly in  $[\epsilon, R]$  and hence

$$\int_{[\epsilon e^{i\delta}, Re^{i\delta}]} \frac{h(z)}{z+1} dz \rightarrow \int_{\epsilon}^R \frac{r^{a-1}}{r+1} dr \quad \text{when } \delta \rightarrow 0+. \quad (6.24)$$

We also have

$$\int_{[Re^{i(2\pi-\delta)}, \epsilon e^{i(2\pi-\delta)}]} \frac{h(z)}{z+1} dz = -e^{ia(2\pi-\delta)} \int_{\epsilon}^R \frac{r^{a-1}}{re^{-i\delta}+1} dr.$$

Keeping  $\epsilon$  and  $R$  fixed, we take the limit when  $\delta \rightarrow 0+$ . Exactly as with (6.24), we get

$$\int_{[Re^{i(2\pi-\delta)}, \epsilon e^{i(2\pi-\delta)}]} \frac{h(z)}{z+1} dz \rightarrow -e^{i2a\pi} \int_{\epsilon}^R \frac{r^{a-1}}{r+1} dr \quad \text{when } \delta \rightarrow 0+. \quad (6.25)$$

From (6.23), (6.24) and (6.25) we get

$$\left| (1 - e^{i2a\pi}) \int_{\epsilon}^R \frac{r^{a-1}}{r+1} dr - 2\pi i e^{(a-1)\pi i} \right| \leq \frac{2\pi R^a}{R-1} + \frac{2\pi \epsilon^a}{1-\epsilon}.$$

Finally, we let  $\epsilon \rightarrow 0+$  and  $R \rightarrow +\infty$  and we conclude that

$$\int_0^{+\infty} \frac{x^{a-1}}{x+1} dx = \lim_{\epsilon \rightarrow 0+, R \rightarrow +\infty} \int_{\epsilon}^R \frac{r^{a-1}}{r+1} dr = \frac{2\pi i e^{(a-1)\pi i}}{1 - e^{i2a\pi}} = \frac{\pi}{\sin a\pi}.$$

**Example 6.5.11.** Evaluation of  $\int_0^{2\pi} r(\sin \theta, \cos \theta) d\theta$ , where  $r(s, t)$  is a rational function of two variables.

We parametrize  $C_0(1)$  with  $z = e^{i\theta}$ ,  $\theta \in [0, 2\pi]$ , and we have  $\cos \theta = \frac{1}{2}(z + \frac{1}{z})$ ,  $\sin \theta = \frac{1}{2i}(z - \frac{1}{z})$  and  $\frac{dz}{d\theta} = ie^{i\theta} = iz$ . Hence

$$\int_0^{2\pi} r(\sin \theta, \cos \theta) d\theta = \frac{1}{i} \oint_{C_0(1)} r\left(\frac{z^2+1}{2z}, \frac{z^2-1}{2iz}\right) \frac{1}{z} dz.$$

The function  $s(z) = r\left(\frac{z^2+1}{2z}, \frac{z^2-1}{2iz}\right) \frac{1}{z}$  is a rational function of  $z$ . We apply the residue theorem after we evaluate the residues of  $s$  at its poles in the disc  $D_0(1)$ .

### Exercises.

**6.5.1.** Find the singular parts as well as the residues of  $\frac{1}{z^2+5z+6}$ ,  $\frac{1}{(z^2-1)^2}$ ,  $e^z + e^{1/z}$ ,  $\frac{\cos z-1}{z^4}$  at their isolated singularities.

**6.5.2.** Find the residues of  $\frac{1}{\sin z}$ ,  $\tan z$ ,  $\frac{1}{\sin^2 z}$ ,  $\frac{1}{e^z-1}$  at their isolated singularities.

**6.5.3.** If  $f = gh$ , where  $g$  is holomorphic at  $z_0$  and  $h$  has a pole of order 1 at  $z_0$ , prove that  $\text{Res}(f; z_0) = g(z_0) \text{Res}(h; z_0)$ .

**6.5.4.** Let  $f = \frac{g}{h}$ , where  $g, h$  are holomorphic in a neighborhood of  $z_0$ . Assume that  $z_0$  is a root of  $h$  of multiplicity  $N$  and not a root of  $g$ . Then  $z_0$  is a pole of  $f$  of order  $N$ .

(i) If  $N = 1$ , prove that  $\text{Res}(f; z_0) = \frac{g(z_0)}{h'(z_0)}$ .

(ii) If  $N = 2$ , prove that  $\text{Res}(f; z_0) = \frac{6g'(z_0)h''(z_0)-2g(z_0)h'''(z_0)}{3h''(z_0)^2}$ .

**6.5.5.** Evaluate  $\int_{-\infty}^{+\infty} \frac{1}{x^2+1} dx$ ,  $\int_{-\infty}^{+\infty} \frac{1}{(x^2+1)(x^2+4)} dx$ ,  $\int_{-\infty}^{+\infty} \frac{1}{(x^2+1)^2} dx$ ,  $\int_{-\infty}^{+\infty} \frac{x^4}{1+x^8} dx$ .

**6.5.6.** Evaluate  $\text{pv} \int_{-\infty}^{+\infty} \frac{x+1}{x^2+1} dx$ ,  $\text{pv} \int_{-\infty}^{+\infty} \frac{x^3}{x^4-4x^2+5} dx$ ,  $\text{pv} \int_{-\infty}^{+\infty} \frac{x^2+3}{x(x^2+1)} dx$ .

**6.5.7.** Evaluate  $\int_{-\infty}^{+\infty} \frac{\cos x}{(x^2+1)(x^2+4)} dx$ ,  $\int_{-\infty}^{+\infty} \frac{x^3 \sin x}{x^4+1} dx$ ,  $\text{pv} \int_{-\infty}^{+\infty} \frac{\cos x}{x(x^2+1)} dx$ .

**6.5.8.** Evaluate  $\int_0^{2\pi} \frac{1}{(1-a \cos \theta)^2} d\theta$ ,  $\int_0^{2\pi} \frac{\cos 2\theta}{1-2a \cos \theta+a^2} d\theta$  when  $0 < a < 1$ ,  $\int_0^{\pi/2} \frac{1}{a+\sin^2 x} dx$  when  $|a| > 1$ .

**6.5.9.** Evaluate  $\int_0^{+\infty} \frac{x^a}{x^2+3x+2} dx$  when  $-1 < a < 1$ .

**6.5.10.** Evaluate  $\int_0^{+\infty} \frac{\ln x}{(x^2+1)(x^2+4)} dx$ ,  $\int_0^{+\infty} \frac{\ln^2 x}{x^2+1} dx$ ,  $\int_0^{+\infty} \frac{\ln(1+x^2)}{x^{1+a}} dx$  when  $0 < a < 2$ .

**6.5.11.** Evaluate  $\int_{-\infty}^{+\infty} \frac{\cos x}{e^x+e^{-x}} dx$ .

**6.5.12.** Evaluate  $\int_0^{+\infty} \frac{1}{x^3+8} dx$ ,  $\int_0^{+\infty} \frac{x}{x^4+16} dx$ , using  $\int_0^{+\infty} \frac{x^{a-1}}{x+1} dx = \frac{\pi}{\sin a\pi}$ .

**6.5.13.** Evaluate  $\int_0^{2\pi} \frac{1}{2+\cos \theta} d\theta$ .

**6.5.14.** If  $z_1, \dots, z_N \in D_0(R)$  are distinct and  $f$  is holomorphic in an open set containing  $\overline{D_0(R)}$ , prove that  $\oint_{C_0(R)} \frac{f(z)}{(z-z_1)\cdots(z-z_N)} dz = 2\pi i \left( \frac{f(z_1)}{p'(z_1)} + \cdots + \frac{f(z_N)}{p'(z_N)} \right)$ , where  $p(z) = (z-z_1)\cdots(z-z_N)$ .

**6.5.15.** If  $n \in \mathbb{N}$ , evaluate  $\oint_{C_0(n)} \tan \pi z dz$ .

**6.5.16.** Let  $r = \frac{q}{p}$  be a rational function with  $\deg q \geq \deg p + 2$ . If  $z_1, \dots, z_n$  are the distinct roots of  $q$ , prove that  $\sum_{k=1}^n \text{Res}(r; z_k) = 0$ .

What is the value of  $\sum_{k=1}^n \text{Res}(r; z_k)$  if  $\deg q = \deg p + 1$ ?

**6.5.17.** If  $f(z) = e^{z+(1/z)}$ , prove that  $\text{Res}(f; 0) = \sum_{n=0}^{+\infty} \frac{1}{n!(n+1)!}$ .

**6.5.18.** (i) If  $n \in \mathbb{N}$ , prove that there is  $M \geq 0$  independent of  $n$  so that  $|\cot z| \leq M$  for every  $z \in \partial R_n$ , where  $R_n$  is the square region with corners at the points  $\pm(n + \frac{1}{2})\pi \pm i(n + \frac{1}{2})\pi$ .

(ii) Prove that  $\oint_{\partial R_n} \frac{\cot z}{z^2} dz \rightarrow 0$  when  $n \rightarrow +\infty$ .

(iii) Prove that  $\sum_{n=1}^{+\infty} \frac{1}{n^2} = \frac{\pi^2}{6}$ .

(iv) Let  $f$  be holomorphic in  $\mathbb{C}$  except for poles  $z_1, \dots, z_N$  and let  $\lim_{z \rightarrow \infty} z f(z) = 0$ . Prove that  $\lim_{n \rightarrow +\infty} \sum_{k=-n}^n f(k) = -\sum_{j=1}^N \text{Res}(f(z) \cot z; z_j)$ .

(v) Find the sums  $\sum_{n=1}^{+\infty} \frac{1}{n^2+a^2}$ , where  $a > 0$ , and the sum  $\sum_{-\infty}^{+\infty} \frac{1}{(n+a)^2}$ , where  $a \notin \mathbb{Z}$ .

**6.5.19.** Let  $p, q$  be polynomials with  $\deg q \geq \deg p + 2$ . For each  $m \in \mathbb{N}$  let  $I_m$  be the square region with corners at  $(m + \frac{1}{2})(\pm 1 \pm i)$ .

- (i) Prove that  $\oint_{\partial I_m} \frac{1}{\sin \pi z} \frac{p(z)}{q(z)} dz \rightarrow 0$  when  $m \rightarrow +\infty$ .  
(ii) Prove the same result if  $\deg q = \deg p + 1$ .

**6.5.20.** Let  $-1 < \nu < 1$  and  $n \in \mathbb{N}$ . Prove that  $\oint_{C_0(n+\frac{1}{2})} \frac{e^{i\nu\pi z}}{\sin \pi z} \frac{1}{z-a} dz \rightarrow 0$  when  $n \rightarrow +\infty$ .

**6.5.21.** (i) Let  $p, q$  be polynomials with  $\deg q \geq \deg p + 1$  and  $q(k) \neq 0$  for every  $k \in \mathbb{Z}$ . Prove that  $\lim_{m \rightarrow +\infty} \sum_{k=-m}^m \frac{p(k)}{q(k)}$  is equal to the sum of the residues of  $-\pi \cot \pi z \frac{p(z)}{q(z)}$  at the roots of  $q$ . Also, prove that  $\lim_{m \rightarrow +\infty} \sum_{k=-m}^m (-1)^k \frac{p(k)}{q(k)}$  is equal to the sum of the residues of  $-\pi \frac{1}{\sin \pi z} \frac{p(z)}{q(z)}$  at the roots of  $q$ .

(ii) Prove  $\pi \cot \pi w = \lim_{m \rightarrow +\infty} \sum_{k=-m}^m \frac{1}{w-k} = \frac{1}{w} + \sum_{k=-\infty}^{+\infty} (\frac{1}{w-k} + \frac{1}{k})$  if  $w \notin \mathbb{Z}$ .

(iii) Prove  $\frac{\pi^2}{\sin^2 \pi w} = \sum_{k=-\infty}^{+\infty} \frac{1}{(w-k)^2}$  if  $w \notin \mathbb{Z}$ .

(iv) Prove  $\sum_{k=1}^{+\infty} \frac{1}{a+bk^2} = -\frac{1}{2a} + \frac{\pi}{2\sqrt{ab}} \frac{e^{\pi\sqrt{a/b}} + e^{-\pi\sqrt{a/b}}}{e^{\pi\sqrt{a/b}} - e^{-\pi\sqrt{a/b}}}$  if  $ab > 0$ .

(v) Prove  $\sum_{k=1}^{+\infty} \frac{(-1)^k}{a+bk^2} = -\frac{1}{2a} - \frac{\pi}{\sqrt{ab}} \frac{1}{e^{\pi\sqrt{a/b}} - e^{-\pi\sqrt{a/b}}}$  if  $ab > 0$ .

## 6.6 The argument principle. The theorem of Rouché.

A function  $f$  is called **meromorphic** in the open set  $\Omega$  if it is holomorphic in  $\Omega$  except at certain points in  $\Omega$  which are poles of  $f$ .

Let  $f$  be meromorphic in the open set  $\Omega$ . If  $w \in \mathbb{C}$ , we shall denote  $A_w$  the set of solutions of  $f(z) = w$ , i.e.

$$A_w = \{z \in \Omega \mid f(z) = w\}.$$

If  $f$  is not constant in any connected component of  $\Omega$ , then the solutions of  $f(z) = w$  are isolated points.

Also, letting  $f$  have the value  $\infty$  at each of its poles in  $\Omega$ , so that  $f$  becomes continuous at its poles considered as a function from  $\Omega$  to  $\hat{\mathbb{C}}$ , we denote  $A_\infty$  the set of solutions of  $f(z) = \infty$ , i.e.

$$A_\infty = \{z \in \Omega \mid f(z) = \infty\} = \{z \in \Omega \mid z \text{ is a pole of } f\}.$$

**The argument principle.** Let  $w \in \mathbb{C}$ . We assume that  $f$  is meromorphic in the open set  $\Omega$  and that it is not constant in any connected component of  $\Omega$ . We also consider  $\Sigma$  to be a cycle, which consists of piecewise smooth closed curves and which is null-homologous in  $\Omega$ , so that no element of  $A_w \cup A_\infty$  is in the trajectory of any of the closed curves forming  $\Sigma$ . Then  $n(\Sigma; z) \neq 0$  for at most finitely many elements of  $A_w \cup A_\infty$  and hence the sums  $\sum_{z \in A_w} n(\Sigma; z) m(z)$  and  $\sum_{z \in A_\infty} n(\Sigma; z) m(z)$ , where  $m(z)$  is the corresponding multiplicity of  $z \in A_w \cup A_\infty$ , are finite. Moreover,

$$n(f(\Sigma); w) = \frac{1}{2\pi i} \oint_\Sigma \frac{f'(\zeta)}{f(\zeta)-w} d\zeta = \sum_{z \in A_w} n(\Sigma; z) m(z) - \sum_{z \in A_\infty} n(\Sigma; z) m(z). \quad (6.26)$$

Furthermore, even if the closed curves which form  $\Sigma$  are not necessarily piecewise smooth, then the left and the right side of (6.26) are still equal.

*Proof.* At first we assume that the closed curves forming  $\Sigma$  are all piecewise continuous.

We apply the residue theorem to the function  $\frac{f'}{f-w}$ . The isolated singularities of this function are the elements of  $A_w \cup A_\infty$ .

If  $m(z)$  is the multiplicity of  $z \in A_w$ , then there is a  $g$  holomorphic in some neighborhood  $D_z(r)$  of  $z$  so that  $f(\zeta) - w = (\zeta - z)^{m(z)} g(\zeta)$  when  $\zeta \in D_z(r)$  and also  $g(z) \neq 0$ . Since  $g(z) \neq 0$ , we may assume that  $r$  is small enough so that  $g(\zeta) \neq 0$  when  $\zeta \in D_z(r)$ . Therefore

$$\frac{f'(\zeta)}{f(\zeta)-w} = \frac{m(z)}{\zeta-z} + \frac{g'(\zeta)}{g(\zeta)} \quad \text{when } \zeta \in D_z(r) \setminus \{z\}.$$



Since  $\frac{g'}{g}$  is holomorphic in  $D_z(r)$ , we have that  $z$  is a pole of  $\frac{f'}{f-w}$  of order 1 with residue  $m(z)$ . If  $m(z)$  is the order of  $z \in A_\infty$ , there is a  $g$  holomorphic in some neighborhood  $D_z(r)$  of  $z$  so that  $f(\zeta) - w = \frac{g(\zeta)}{(\zeta-z)^{m(z)}}$  when  $\zeta \in D_z(r)$  and also  $g(z) \neq 0$ . Since  $g(z) \neq 0$ , we may assume that  $r$  is small enough so that  $g(\zeta) \neq 0$  when  $\zeta \in D_z(r)$ . Hence

$$\frac{f'(\zeta)}{f(\zeta)-w} = \frac{-m(z)}{\zeta-z} + \frac{g'(\zeta)}{g(\zeta)} \quad \text{when } \zeta \in D_z(r) \setminus \{z\}.$$

Since  $\frac{g'}{g}$  is holomorphic in  $D_z(r)$ , we have that  $z$  is a pole of  $\frac{f'}{f-w}$  of order 1 with residue  $-m(z)$ . Now, the residue theorem implies the second equality in (6.26). The first equality is a matter of a simple change of variable. If  $\zeta = \gamma(t)$ ,  $t \in [a, b]$ , is the parametric equation of any curve  $\gamma$  forming  $\Sigma$ , then the parametric equation of  $f(\gamma)$  is  $\eta = f(\gamma(t))$ ,  $t \in [a, b]$ , and hence:

$$n(f(\gamma); w) = \frac{1}{2\pi i} \oint_{f(\gamma)} \frac{1}{\eta-w} d\eta = \frac{1}{2\pi i} \int_a^b \frac{f'(\gamma(t))\gamma'(t)}{f(\gamma(t))-w} dt = \frac{1}{2\pi i} \oint_\gamma \frac{f'(\zeta)}{f(\zeta)-w} d\zeta.$$

The rest is simple if we recall that  $\Sigma = n_1\gamma_1 + \dots + n_k\gamma_k$  and  $f(\Sigma) = n_1f(\gamma_1) + \dots + n_kf(\gamma_k)$ . Now we assume that the curves  $\gamma$  which form  $\Sigma$  are not necessarily piecewise smooth.

We consider any of the closed curves which form  $\Sigma$  with parametric equation  $\zeta = \gamma(t)$ ,  $t \in [a, b]$ , and the corresponding  $f(\gamma)$  with parametric equation  $\eta = f(\gamma(t))$ ,  $t \in [a, b]$ . The set  $A_w \cup A_\infty$  has no accumulation point in  $\Omega$ . Thus, the set  $A_w \cup A_\infty \cup \Omega^c$  is closed and we also have that it is disjoint from  $\gamma^*$ . Therefore, there is  $\epsilon_1 > 0$  so that

$$|\gamma(t) - z| \geq 2\epsilon_1 \quad \text{for every } t \in [a, b] \text{ and every } z \in A_w \cup A_\infty \cup \Omega^c. \quad (6.27)$$

We consider the set

$$K = \{z \mid |z - \gamma(t)| \leq \epsilon_1 \text{ for at least one } t \in [a, b]\}$$

and we easily see that  $K$  is a compact subset of  $\Omega \setminus (A_w \cup A_\infty)$  and hence  $f$  is continuous in  $K$ . Also, we have  $f(z) \neq w$  for every  $z \in K$  and  $\gamma^*$  is a subset of  $K$  and hence there is  $\epsilon_2 > 0$  so that

$$|f(\gamma(t)) - w| \geq \epsilon_2 \quad \text{for every } t \in [a, b]. \quad (6.28)$$

Since  $f$  is continuous in  $K$ , there is  $\delta_1$  with  $0 < \delta_1 \leq \epsilon_1$  so that

$$|f(z') - f(z'')| < \epsilon_2 \quad \text{for every } z', z'' \in K \text{ with } |z' - z''| < \delta_1. \quad (6.29)$$

Finally, there is  $\delta > 0$  so that

$$|\gamma(t') - \gamma(t'')| < \delta_1 \quad \text{for every } t', t'' \in [a, b] \text{ with } |t' - t''| < \delta. \quad (6.30)$$

Now we take successive points  $a = t_0 < t_1 < \dots < t_{n-1} < t_n = b$  so that  $t_k - t_{k-1} < \delta$  for every  $k$  and we consider the polygonal curve  $\sigma : [a, b] \rightarrow \mathbb{C}$  consisting of the successive linear segments  $[\gamma(t_{k-1}), \gamma(t_k)]$ . It is easy to see that we have

$$|\sigma(t) - \gamma(t)| < \delta_1 \leq \epsilon_1 \quad \text{for every } t \in [a, b]. \quad (6.31)$$

Indeed, if  $t \in [t_{k-1}, t_k]$ , then, because of (6.30), we have

$$\begin{aligned} |\sigma(t) - \gamma(t)| &= \left| \left( \frac{t_k-t}{t_k-t_{k-1}} \gamma(t_{k-1}) + \frac{t-t_{k-1}}{t_k-t_{k-1}} \gamma(t_k) \right) - \gamma(t) \right| \\ &\leq \frac{t_k-t}{t_k-t_{k-1}} |\gamma(t_{k-1}) - \gamma(t)| + \frac{t-t_{k-1}}{t_k-t_{k-1}} |\gamma(t_k) - \gamma(t)| \\ &< \frac{t_k-t}{t_k-t_{k-1}} \delta_1 + \frac{t-t_{k-1}}{t_k-t_{k-1}} \delta_1 = \delta_1 \leq \epsilon_1. \end{aligned}$$

Now, (6.27), (6.31) imply  $|\sigma(t) - \gamma(t)| < |\gamma(t) - z|$  for every  $t \in [a, b]$  and every  $z \in A_w \cup A_\infty$ . Proposition 6.9 implies  $n(\gamma; z) = n(\sigma; z)$  for every  $z \in A_w \cup A_\infty$  and hence

$$\begin{aligned} \sum_{z \in A_w} n(\gamma; z) m(z) - \sum_{z \in A_\infty} n(\gamma; z) m(z) \\ = \sum_{z \in A_w} n(\sigma; z) m(z) - \sum_{z \in A_\infty} n(\sigma; z) m(z). \end{aligned} \quad (6.32)$$

Also, (6.31) implies  $\sigma(t) \in K$  for every  $t \in [a, b]$  and, because of (6.29),  $|f(\sigma(t)) - f(\gamma(t))| < \epsilon_2$  for every  $t \in [a, b]$ . But then (6.28) implies  $|f(\sigma(t)) - f(\gamma(t))| < |f(\gamma(t)) - w|$  for every  $t \in [a, b]$ . Proposition 6.9 again implies

$$n(f(\gamma); w) = n(f(\sigma); w). \quad (6.33)$$

Since the curve  $\sigma$  is piecewise smooth, we have from the first part of the proof that

$$n(f(\sigma); w) = \sum_{z \in A_w} n(\sigma; z) m(z) - \sum_{z \in A_\infty} n(\sigma; z) m(z). \quad (6.34)$$

Now, (6.32), (6.33) and (6.34) imply the equality of the left and the right side of (6.26) for each  $\gamma$  forming  $\Sigma$  and the proof is finished by addition over all such  $\gamma$ .  $\square$

The geometric content of the argument principle is described as follows. *The number of rotations of  $f(\Sigma)$  around  $w$  is equal to the total number of rotations of  $\Sigma$  around the solutions of  $f(z) = w$  minus the total number of rotations of  $\Sigma$  around the poles of  $f$ .* When we count the solutions of  $f(z) = w$  and the poles of  $f$  we take into account their multiplicities. We count  $m(z)$  points at every point  $z \in A_w \cup A_\infty$  which has multiplicity  $m(z)$ .

If  $f$  has no pole in  $\Omega$ , i.e. if  $f$  is holomorphic in  $\Omega$ , then the argument principle says that *the number of rotations of  $f(\Sigma)$  around  $w$  is equal to the total number of rotations of  $\Sigma$  around the solutions of  $f(z) = w$ .* In fact, if  $\Sigma$  is such that for every  $z$  not in the trajectories of the curves forming  $\Sigma$  we have either  $n(\Sigma; z) = 1$  or  $n(\Sigma; z) = 0$ , then *the number of rotations of  $f(\Sigma)$  around  $w$  is equal to the number of solutions of  $f(z) = w$  which are surrounded by  $\Sigma$ .*

**The theorem of Rouché.** *Let  $w \in \mathbb{C}$ . We assume that  $f, g$  are holomorphic in the open set  $\Omega$  and that they are not constant in any connected component of  $\Omega$ . We also consider  $\Sigma$  to be a cycle which is null-homologous in  $\Omega$ . If  $|f(\zeta) - g(\zeta)| < |g(\zeta) - w|$  for every  $\zeta$  in the trajectories of the closed curves forming  $\Sigma$ , then*

$$\sum_{z \in A_{w,f}} n(\Sigma; z) m_f(z) = \sum_{z \in A_{w,g}} n(\Sigma; z) m_g(z),$$

where  $m_f(z)$  and  $m_g(z)$  are the corresponding multiplicities and  $A_{w,f} = \{z \in \Omega \mid f(z) = w\}$ ,  $A_{w,g} = \{z \in \Omega \mid g(z) = w\}$ .

*Proof.* We observe that the condition  $|f(\zeta) - g(\zeta)| < |g(\zeta) - w|$  for every  $\zeta$  in the trajectories of the closed curves forming  $\Sigma$  implies that no element of  $A_{w,f} \cup A_{w,g}$  is in these trajectories. The function  $h = \frac{f-w}{g-w}$  is holomorphic in  $\Omega$  except for the elements of  $A_{w,g}$ , which are either poles or removable singularities of  $h$ . From (6.26) we have

$$n(h(\Sigma); 0) = \sum_{z \in A_{0,h}} n(\Sigma; z) m_h(z) - \sum_{z \in A_{\infty,h}} n(\Sigma; z) m_h(z). \quad (6.35)$$

If  $z \in A_{w,f} \setminus A_{w,g}$ , then  $z \in A_{0,h}$  and  $m_h(z) = m_f(z)$ . Similarly, if  $z \in A_{w,g} \setminus A_{w,f}$ , then  $z \in A_{\infty,h}$  and  $m_h(z) = m_g(z)$ . Finally, if  $z \in A_{w,f} \cap A_{w,g}$ , then we have three cases. If  $m_f(z) > m_g(z)$ , then  $z \in A_{0,h}$  and  $m_h(z) = m_f(z) - m_g(z)$ . If  $m_f(z) < m_g(z)$ , then  $z \in A_{\infty,h}$  and  $m_h(z) = m_g(z) - m_f(z)$ . If  $m_f(z) = m_g(z)$ , then  $z \notin A_{0,h} \cup A_{\infty,h}$  and  $m_h(z) = 0$ . All these imply

$$\begin{aligned} & \sum_{z \in A_{0,h}} n(\Sigma; z) m_h(z) - \sum_{z \in A_{\infty,h}} n(\Sigma; z) m_h(z) \\ &= \sum_{z \in A_{w,f}} n(\Sigma; z) m_f(z) - \sum_{z \in A_{w,g}} n(\Sigma; z) m_g(z) \end{aligned}$$

and from (6.35) we get

$$\sum_{z \in A_{w,f}} n(\Sigma; z) m_f(z) - \sum_{z \in A_{w,g}} n(\Sigma; z) m_g(z) = n(h(\Sigma); 0).$$

Now, our hypothesis says that  $|h(z) - 1| < 1$  for every  $z$  in the trajectories of the curves forming  $\Sigma$ . Therefore, the cycle  $h(\Sigma)$  is in the disc  $D_1(1)$  and hence  $n(h(\Sigma); 0) = 0$ .  $\square$

**Example 6.6.1.** We shall find the number of roots of  $f(z) = z^7 - 2z^5 + 6z^3 - z + 1$  in  $\mathbb{D}$ . We consider  $g(z) = 6z^3$  and we have

$$|f(z) - g(z)| = |z^7 - 2z^5 - z + 1| \leq |z|^7 + 2|z|^5 + |z| + 1 = 5 < 6|z|^3 = |g(z)|$$

for every  $z \in \mathbb{T}$ . Now we apply the theorem of Rouché with  $w = 0$  and  $\Sigma$  consisting of only the curve  $\gamma$  which describes  $\mathbb{T}$  once and in the positive direction. We have  $n(\gamma; z) = 1$  for every  $z \in \mathbb{D}$  and  $n(\gamma; z) = 0$  for every  $z \notin \overline{\mathbb{D}}$ . The only solution of  $g(z) = 0$  in  $\mathbb{D}$  is  $z = 0$  with multiplicity  $m_g(0) = 3$ . Therefore  $\sum_{z \in A_{0,g}} n(\gamma; z) m_g(z) = \sum_{z \in A_{0,g} \cap \mathbb{D}} m_g(z) = 3$ . Moreover,  $\sum_{z \in A_{0,f}} n(\gamma; z) m_f(z) = \sum_{z \in A_{0,f} \cap \mathbb{D}} m_f(z)$ . Now the theorem of Rouché implies that  $\sum_{z \in A_{0,f} \cap \mathbb{D}} m_f(z) = 3$  and hence  $f$  has three roots in  $\mathbb{D}$ .

### Exercises.

**6.6.1.** Let  $f$  be holomorphic in  $D_{z_0}(R)$ , let  $0 < r < R$  and assume that there is no solution of  $f(z) = w$  in  $C_{z_0}(r)$ . If  $k \in \mathbb{N}$ , what is the content of  $\frac{1}{2\pi i} \oint_{C_{z_0}(r)} \frac{f'(z)}{f(z)-w} z^k dz$ ?

**6.6.2.** Let  $f$  be holomorphic in  $\mathbb{D}$  and continuous in  $\overline{\mathbb{D}}$  and let  $|f(z)| < 1$  for every  $z \in \mathbb{T}$ . Prove that the equation  $f(z) = z^n$  has exactly  $n$  solutions in  $\mathbb{D}$ .

**6.6.3.** Find the number of roots of

(i)  $z^4 - 6z + 3$  in  $D_0(1, 2)$ .

(ii)  $z^4 + 8z^3 + 3z^2 + 8z + 3$  in  $\{z \mid \operatorname{Re} z > 0\}$ .

**6.6.4.** Let  $z_1, \dots, z_n \in \mathbb{D}$ . In  $\mathbb{C} \setminus \left\{ \frac{1}{z_1}, \dots, \frac{1}{z_k} \right\}$  we consider the function  $f(z) = \prod_{k=1}^n \frac{z-z_k}{1-\overline{z_k}z}$ .

(i) Prove that  $f(z) \in \mathbb{D}$  for every  $z \in \mathbb{D}$  and that  $f(z) \in \mathbb{T}$  for every  $z \in \mathbb{T}$ .

(ii) Find the index with respect to 0 of the curve with parametric equation  $z = f(e^{it})$ ,  $t \in [0, 2\pi]$ .

(iii) Prove that for every  $w \in \mathbb{D}$  the equation  $f(z) = w$  has exactly  $n$  solutions in  $\mathbb{D}$ .

**6.6.5.** Prove that the set of all meromorphic functions in the region  $\Omega$  is an algebraic field.

**6.6.6.** Let  $f$  be holomorphic in the open set  $\Omega$ . We assume that  $\gamma$  is a closed piecewise smooth curve in  $\Omega$ , that  $\mathbb{C} \setminus \gamma^*$  has only one bounded connected component  $U$  and that  $n(\gamma; z) = 1$  for every  $z \in U$ . We also assume that  $\mathbb{C} \setminus f(\gamma)^*$  has only one bounded connected component  $V$  and that  $n(f(\gamma); w) = N$  for every  $w \in V$ .

(i) If  $f(z) \notin f(\gamma)^*$  for every  $z \in U$ , prove that  $f$  is  $N$ -to-one from  $U$  onto  $V$ .

(ii) If moreover  $N = 1$ , we may consider the inverse function  $f^{-1} : V \rightarrow U$ . Prove that  $f^{-1}(w) = \frac{1}{2\pi i} \oint_{\Sigma} \frac{\zeta f'(\zeta)}{f(\zeta)-w} d\zeta$  for every  $w \in V$ .

**6.6.7.** Let  $f(z) = \sum_{n=0}^{+\infty} a_n z^n$  for  $z \in \mathbb{D}$  and let  $F \subseteq \mathbb{D}$  be compact with  $0 \in F$ . If  $\mu = \inf_{z \in \partial F} |f(z)|$  and  $m$  is the number of roots of  $f$  in  $F$ , prove that  $\mu \leq |a_0| + |a_1| + \dots + |a_m|$ .